


General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

Annual Report
for
NASA Grant NAG5-58
entitled



"A Laboratory Investigation of the Reflective Properties
of Simulated, Optically Thick Clouds"

(NASA-CR-175362) A LABORATORY INVESTIGATION N84-17755
OF THE REFLECTIVE PROPERTIES OF SIMULATED,
OPTICALLY THICK CLOUDS Annual Report, 15
May 1980 - 15 May 1982 (Colorado State Univ.) 15 p HC A02/MF A01 CSCI 04B G3/47 11624
Unclas

Principal Investigator: Thomas B. McKee
Co-Investigator: Stephen K. Cox

Department of Atmospheric Science
Colorado State University
Fort Collins, Colorado 80523

Grant period: 05/15/80 - 05/15/82

This report summarizes the progress toward the specific research goals of the cloud field simulation project which has been made since June 1981. Accomplishments prior to this date were reported in our first semi-annual progress report in February 1981 and at the Fourth Conference on Atmospheric Radiation, Toronto.¹

During the reporting period stated above, progress toward the research goals of CFOS (the Cloud Field Optical Simulator) may be noted in the following areas: (1) improvement in the shape of the desired (visible) spectral response of the measurement, (2) selection of two usable materials for cloud simulation, (3) a means of assigning a "visible" optical depth to the simulated clouds and (4) confirmation that the apparatus is capable of detecting basic finite cloud characteristics. A brief description of the accomplishments in each of these areas follows.

(1) Improved Spectral Bandpass

In order to insure that the measurements were limited to the visible portion of the spectrum an optical filter was placed in front of each detector. Figure 1 shows the relative responses of the system with and without the filter. Although insertion of the filter resulted in a relatively poorer S/N ratio it ensures that the measurements will not include spectrally dependent effects which are visually unobservable.

¹McKee, T. B., J. M. Davis, and S. K. Cox, 1981: Design and verification of a cloud field optical simulator. Preprints Fourth Conference Atmospheric Radiation, Toronto, Amer. Meteor. Soc., pg. 227.

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 1. Relative spectral response of the original and filtered CFOS photodiode.

2. Selection of Suitable Materials for Cloud Simulation

Several materials have been examined in a search for one which adequately simulates the visible reflective properties of optically thick clouds. The criteria which were used to rate the materials were first, the behavior of the radiances reflected into the principal plane by a horizontally semi-infinite cloud, and second, the visual appearance of the clouds simulated from the material. Based on these criteria, two materials were found to be superior to all others tested; they are surgical quality sterilized cotton and decorative billet styrofoam which is marketed by Dow Chemical. Figures 2 and 3 compare measured principal plane reflected radiances of the two materials with theoretical curves for semi-infinite clouds from Monte Carlo radiative transfer calculations. The model was run for two solar zenith angles $\mu_0 = 1.0$ and $\mu_0 = 0.5$ using the C.1 droplet distribution at a wavelength of $0.7 \mu\text{m}$. All radiances have been normalized by the dividing by reflected radiance at the zenith. The materials are almost equivalent in satisfying the first criteria. However, the styrofoam is slightly better in the backscatter direction. The assignment of an equivalent optical depth to the simulation materials is discussed in the next section.

Both materials do well in simulating the visual appearance of clouds. If we add a third criterion, namely the practicality of working with the materials, the styrofoam becomes preferable for a number of reasons. First, the styrofoam material is rigid and will maintain a constant ratio of geometric to optical depth, and the shape of the simulated clouds will not distort due to gravity with rotation of the




Figure 2. A comparison between the calculated and simulated reflected radiances from semi-infinite clouds for solar zenith angles of 0° and 60° using cotton as the cloud simulation material.




Figure 3. A comparison between the calculated and simulated reflected radiances from semi-infinite clouds for solar zenith angles of 0° and 60° using styrofoam as the simulation material.

CFOS apparatus about the simulation zenith. Second, simulated styrofoam clouds are easily exchanged from one cloud scene to another or moved about within a cloud scene. Finally the styrofoam simulated clouds are easily made into regular shapes, i.e. cubes, cylinders, inverted paraboloids etc. for comparison with theoretical results for finite clouds.

3. A Method of Assigning a Visible Optical Depth to the Simulated Clouds

In order to relate the reflective properties of simulated clouds to real water clouds it is necessary to have an accurate estimate of the optical depth of the simulated clouds. One means of doing so is based on the bulk radiative properties of a 'semi-infinite' sheet of the cloud material at 0° zenith. Specifically theory predicts that the ratio of spectral reflectance R_λ to spectral transmittances T_λ of a semi infinite cloud over a non reflecting surface is nearly a linear function of optical depth, see for example Coakley and Chylek (1975) and Stephens (1978). Figure 4 shows a plot of the ratio of R/T generated from the parameterization of Stephens (1978) for water clouds in the visible portion of the spectrum. Also shown is the same ratio from a simple Eddington-model which employed the Henyey-Greenstein approximation for various cases and also a few results of the R/T ratio from Monte Carlo calculations. The linear dependance between R/T and τ the visible optical depth is apparent.

An equivalent visible optical depth (τ_e) may be assigned to the CFOS cloud simulation material by measurement of the ratio of R/T for a sheet of the material with a large ratio of

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 4. A plot of the ratio of reflectance to transmittance derived from various calculations as a function of optical depth.

horizontal to vertical dimension. Once τ_e is established for a large slab of styrofoam several clouds may be 'sculptured' from the slab while maintaining in each the vertical dimension of the original slab. The procedure requires an additional step in the case of cotton since the vertical structure of the original 'slab' (surgical cotton is available in 12" by 60" sheets) can not be maintained while forming clouds of a realistic shape. However, it was found that the mass of the material in a vertical column of unit cross section is also related to the ratio of the R/T in a nearly linear manner; see figure 5. Thus, for a cloud simulated from surgical cotton an approximate visible optical depth may be assigned by weighing the cloud and measuring the cross sectional area to determine the area mass density then using the R/T line as a transfer function to τ_e . The additional measurement is yet another limitation to the suitability of using cotton as the cloud simulation material.

4. Retrieval of Basic Finite Cloud Features

The most recent accomplishment with the CFOS concerns the ability to measure the reflected radiances of finite clouds. Figures 6 and 7 are comparisons between radiances reflected into the principal plane by finite clouds as measured on CFOS and the same quantities predicted by the theoretical Monte Carlo model as described in McKee and Cox (1974). The radiances in each case have been normalized by the radiance measured or calculated at the zenith. The CFOS profiles have been retrieved from measurements over a field of simulated (styrofoam) cubic clouds cut from a slab of material whose R/T ratio corresponds to an equivalent optical

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 5. A plot of the ratio of reflectance to transmittances measured by CFOS for a 'semi-infinite' cotton simulated cloud plotted as a function of area mass density and equivalent visible optical depth (τ_e).

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 6. A comparison between calculated and simulated reflected radiances from finite cubic clouds for a solar zenith angle of 0° using styrofoam for the simulation material.




Figure 7. A comparison between calculated and simulated reflected radiances from finite cubic clouds for a solar zenith angle of 60° using styrofoam for the simulation material.

depth $\tau_e = 76$. The clouds were evenly spaced on a horizontal grid such that the true fraction cloud cover (f) was 0.04. Figures 6 and 7 each show two measured profile retrievals. One assumes that as the nadir angle (θ) increases, the solid angle of cloud field relative to the underlying surface increases as $(1 + \tan \theta)$ resulting in a modified fractional cloud cover $f' = f (1 + \tan \theta)$; see Davis, Cox, and McKee (1979). The second treatment utilizes the exact geometry of the cloud field and the CFOS detectors, counts the clouds in the field of view and calculates the total solid angle subtended by the clouds and their shadows. The radiance measurement (N) is given as

$$N = N_{\text{cld}} \Omega_{\text{cld}} + N_{\text{clr}} \Omega_{\text{clr}} + 0 \Omega_{\text{sdw}}$$

In the above formula the subscripts cld, clr and sdw stand for cloud, clear and shadow, respectively, and Ω represents solid angle. It is assumed that there is no radiance contribution from the shadow regions. The N_{clr} measurements were made over the surface on which the clouds were mounted. The albedo of the surface was 0.02. The second treatment results in a 'noisier' curve because the cloud is either considered completely within or exterior to the field of view. Nevertheless, smoothed versions of the curves are in good agreement. The apparent disagreement between measured and calculated radiances at large nadir in the 0° solar zenith angle case is believed to be due primarily to the basic scattering properties of the material. Similar disagreement is seen in the radiance comparisons for infinite clouds at 0° zenith.

It should be noted that retrieval of the finite cloud reflected radiances, as described above, represents a rigorous test of the CFOS concept. The low surface albedo combined with the extremely small fractional cloud cover result in reflected radiances at the lowest bound of the domain for which CFOS was originally intended. Larger clouds, smaller cloud spacing and more realistic simulated surfaces will all tend to increase the radiance signal. In addition, the above experiment represents the retrieval radiances reflected from only part of the scene. Measurement of total scene radiances are of a more stable nature. Thus, it appears that the CFOS concept is indeed a valid means of exploring the effects of realistic cloud geometries on the radiances reflected by optically thick clouds and cloud fields.

CONTROL POINT IN
OF POOR QUALITY

References

- Coakley, J. A. and P. Chylek, 1975: The two stream approximation in radiative transfer: Including the angle of the incident radiation. J. Atmos. Sci., 32, 409-418.
- Stephens, G. L., 1978: Radiation profiles in extended water clouds. II: Parameterization Schemes. J. Atmos. Sci., 35, 2123-2132.
- McKee, T. B. and S. K. Cox, 1974: Scattering of visible radiation by finite clouds. J. Atmos. Sci., 31, 1885-1892.
- Davis, J. M., S. K. Cox, and T. B. McKee, 1979: Vertical and horizontal distributions of solar absorption in finite clouds. J. Atmos. Sci., 36, 1976-1984.