





Workshop on Biological Integrity of Coral Reefs



Caribbean Coral Reef Institute
Isla Magueyes, La Parguera, Puerto Rico

August 21-22, 2012



Workshop on Biological Integrity of Coral Reefs
August 21-22, 2012
Caribbean Coral Reef Institute
Isla Magueyes, La Parguera, Puerto Rico

by

Patricia Bradley
US EPA
Atlantic Ecology Division
NHEERL, ORD
33 East Quay Road
Key West, FL 33040

Deborah L. Santavy
US EPA
Gulf Ecology Division
NHEERL, ORD
1 Sabine Island Drive
Gulf Breeze, FL 32561

Jeroen Gerritsen
Tetra Tech Inc.
400 Red Brook Boulevard
Suite 200
Owings Mills, MD 21117

Contract No. EP-C-09-001
Work Assignment 3-01
Great Lakes Environmental Center, Inc.

Project Officer:
Shirley Harrison
US EPA
Office of Water
Office of Science and Technology
Washington, DC 20460

Work Assignment Manager:
Susan K. Jackson
US EPA
Office of Water
Office of Science and Technology
Washington, DC 20460

National Health and Environmental Effects Research Laboratory
Office of Research and Development
Washington, DC 20460



Printed on chlorine free 100% recycled paper
with 50% post-consumer fiber using vegetable-based ink.

Notice and Disclaimer

The US Environmental Protection Agency (EPA) through its Office of Research and Development and Office of Water funded and collaborated in the research described here under EP-C-09-001, Work Assignment #3-01, to Great Lakes Environmental Center, Inc. It has been subjected to the Agency's peer and administrative review and has been approved for publication as an EPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

Under authority of the Clean Water Act (CWA), EPA is committed to protecting the biological integrity of the Nation's waters, including marine coastal habitats such as mangroves, seagrasses and coral reefs that lie within the 3-mile territorial waters.

This report summarizes an EPA-sponsored workshop on coral reef biological integrity held at the Caribbean Coral Reef Institute in La Parguera, Puerto Rico, on August 21-22, 2012. The workshop brought together scientists with expertise in coral reef taxonomic groups (e.g., stony corals, fishes, sponges, gorgonians, algae, seagrasses and macroinvertebrates), specializing in community structure, organism condition, ecosystem function and ecosystem connectivity.

The experts evaluated photos and videos for 12 stations collected during EPA coral reef surveys (2010 and 2011) from Puerto Rico coral reefs exhibiting a wide range of conditions. The experts individually rated each station as to observed condition (good, fair or poor) and documented their rationale for the assignment. The group discussed the reef attributes that characterize biological integrity (or the natural condition) for Puerto Rico's coral reefs, which will serve as the baseline condition, since the CWA is grounded in the concept of natural, undisturbed conditions.

The long-term goal is to derive scientifically defensible thresholds for different levels of coral reef condition with a well-defined narrative for each level. Managers will be able to use the narratives to determine which level most appropriately describes the current condition of their coral reefs and which level is the desired condition. From this, managers can set easily communicated, quantitative goals for achieving those conditions. The conceptual model will serve as an interpretative framework to explicitly link science and monitoring information to management and decision-making.

This is a contribution to the EPA Office of Research and Development's Safe and Sustainable Waters Research Program, characterizing the effects of land use on estuarine and coastal resources.

The appropriate citation for this report is:

Bradley P, Santavy DL and Gerritsen J. 2014. Workshop on Biological Integrity of Coral Reefs, August 21-22, 2012, Caribbean Coral Reef Institute, Isla Magueyes, La Parguera, Puerto Rico. US Environmental Protection Agency, Office of Research and Development, Atlantic Ecology Division, Narragansett, RI. EPA/600/R-13/350.

This document can be downloaded from:

http://water.epa.gov/scitech/swguidance/standards/criteria/aqlife/biocriteria/technical_index.cfm

Table of Contents

Figures and Tables	v
Acknowledgements	vii
Executive Summary	viii
Chapter 1. Introduction	
1.1 Coral Reef Ecosystems	1-2
1.2 Puerto Rico's Coral Reef Ecosystems	1-2
1.3 Timelines	1-4
1.3.1 Condition Timeline	1-4
1.3.2 Anthropogenic Activity Timeline	1-7
1.4 Southwestern Puerto Rico	1-9
1.5 The Clean Water Act	1-10
1.6 Why a Coral Reef Ecosystem Conceptual Model is Needed	1-11
1.7 The Framework: The Biological Condition Gradient (BCG)	1-11
1.7.1 How is the BCG Constructed?	1-12
Chapter 2. Approach	
2.1 Video and Photo Evaluations	2-2
2.2 Summary of Ratings	2-5
2.2.1 Best Station, Ranked #1	2-5
2.2.2 Worst Station, Ranked #12	2-6
2.2.3 Stations Rated Fair	2-7
2.2.4 Station Rated Poor	2-8
2.3 Summary of Attributes	2-8
2.4 Reference Condition for Biological Integrity	2-9
2.4.1 Experts' Examples of Reference Condition for Biological Integrity	2-10
2.4.2 Summary Discussion	2-14
2.5 Attributes of a Very Good to Excellent Station	2-14
2.5.1 Three-dimensional Topographic Complexity	2-14
2.5.2 Stony Coral Attributes	2-19
2.5.3 Gorgonian Attributes	2-19
2.5.4 Sponge Attributes	2-20
2.5.5 Fish Attributes	2-20
2.5.6 Large Vertebrate Attributes	2-20
2.5.7 Other Invertebrate Attributes	2-21
2.5.8 Algae Attributes	2-21
2.5.9 Condition	2-21
Chapter 3. Discussion and Next Steps	
3.1 Discussion	3-1
3.2 Next Steps	3-1
3.2.1 Second Workshop	3-1
3.2.2 Species Tolerance Database	3-2
3.2.3 Assembling the Monitoring Data	3-2
3.2.4 Calibrating the BCG	3-3
3.2.5 Economic Valuation of Coastal Ecosystem Services	3-4
3.3 Final Thoughts	3-6

Appendices

A. References	A-1
B. Workshop Participants	B-1
C. Workshop Agenda	C-1
D. Tally Sheet - Rating Condition of Coral Reef Videos (1st)	D-1
E. Tally Sheet - Rating Condition of Coral Reef Videos (2nd)	E-1
F. Notes Sheet - Rating Condition of Coral Reef Videos	F-1
G. Supporting Photos - Rating Condition of Coral Reef Videos	G-1
H. Workshop Glossary	H-1
I. Summary Data Results for BCG Stations	I-1
J. Formulas Used for Calculating Condition Metrics	J-1

Figures

1-1.	The Biological Condition Gradient (BCG)	1-13
2-1.	Map with locations of the 12 EPA stations along the southern coast of Puerto Rico	2-2
2-2.	Recent changes from 2009 to 2012 in <i>Montastraea</i> populations on a coral reef in Curaçao (Watamula) that were caused by a 2010 bleaching event	2-10
2-3.	Two experts suggested Cordelia Banks near Roatan, Honduras, as an example of a reference site for excellent coral condition	2-11
2-4.	Monitoring of massive <i>Acropora cervicornis</i> banks at Cordelia Banks located off a major airport in Roatan, Honduras	2-12
2-5.	Panoramic view of <i>Acropora cervicornis</i> banks at Cordelia Banks, Roatan, Honduras	2-12
2-6.	Cover for the book, <i>Beneath the Waves</i> , by Hector J. Ruiz Torres	2-13
I-1.	Comparison of the density of the major sessile invertebrates assessed in the 12 BCG stations	I-2
I-2.	Comparison of density for fish carnivores vs. herbivores at BCG stations	I-3
I-3.	Comparison of biomass for fish carnivores vs. herbivores at BCG stations	I-3
I-4.	Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 1	I-4
I-5.	Percentages of stony coral species and sponge morphologies for BCG Station 2	I-6
I-6.	Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 3	I-8
I-7.	Percentages of stony coral species and sponge morphologies for BCG Station 4	I-10
I-8.	Percentages of stony coral species and sponge morphologies for BCG Station 5	I-12
I-9.	Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 6	I-14
I-10.	Percentages of stony coral species and gorgonian morphologies for BCG Station 7	I-16
I-11.	Percentages of stony coral species and gorgonian morphologies for BCG Station 8	I-18
I-12.	Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 9	I-20
I-13.	Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 10	I-22
I-14.	Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 11	I-24
I-15.	Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 12	I-26

Tables

1-1.	Number of currently reported species in each of the major marine taxa for Puerto Rico	1-3
1-2.	Complex relationships between stressors exist in southwestern Puerto Rico	1-10
1-3.	Biological and other ecological attributes used to characterize the freshwater streams BCG	1-14
1-4.	Ecological attributes used to characterize the estuarine BCG	1-16
2-1.	Shallow inshore linear reefs used for BCG workshop stations that correspond to US EPA stations sampled along the southern coast of Puerto Rico	2-2

2-2.	Coral reef condition evaluated by the experts	2-4
2-3.	Rankings given by experts after visually evaluating videos and photographs for 12 EPA stations ...	2-4
2-4.	Summary of attributes and their relationships for assessing coral reef condition from station evaluations	2-8
2-5.	Summary of descriptions of four condition categories (very good to poor) based on expert assessments of individual stations	2-15
2-6.	Condition levels and associated attributes	2-17
3-1.	Coral reef ecosystem services and reef attributes	3-5
I-1.	Scleractinian coral summary statistics for BCG stations	I-1
I-2.	Gorgonian summary statistics for BCG stations	I-1
I-3.	Sponge summary statistics for BCG stations	I-2
I-4.	BCG Station 1 data summary for corals and subgroups	I-4
I-5.	Fish species found in BCG Station 1, with density and biomass for 100m ² transect	I-5
I-6.	BCG Station 2 data summary for corals and subgroups	I-6
I-7.	Fish species found in BCG Station 2, with density and biomass for 100m ² transect	I-7
I-8.	BCG Station 3 data summary for corals and subgroups	I-8
I-9.	Fish species found in BCG Station 3, with density and biomass for 100m ² transect	I-9
I-10.	BCG Station 4 data summary for corals and subgroups	I-10
I-11.	Fish species found in BCG Station 4, with density and biomass for 100m ² transect	I-11
I-12.	BCG Station 5 data summary for corals and subgroups	I-12
I-13.	Fish species found in BCG Station 5, with density and biomass for 100m ² transect	I-13
I-14.	BCG Station 6 data summary for corals and subgroups	I-14
I-15.	Fish species found in BCG Station 6, with density and biomass for 100m ² transect	I-15
I-16.	BCG Station 7 data summary for corals and subgroups	I-16
I-17.	Fish species found in BCG Station 7, with density and biomass for 100m ² transect	I-17
I-18.	BCG Station 8 data summary for corals and subgroups	I-18
I-19.	Fish species found in BCG Station 8, with density and biomass for 100m ² transect	I-19
I-20.	BCG Station 9 data summary for corals and subgroups	I-20
I-21.	Fish species found in BCG Station 9, with density and biomass for 100m ² transect	I-21
I-22.	BCG Station 10 data summary for corals and subgroups	I-22
I-23.	Fish species found in BCG Station 10, with density and biomass for 100m ² transect	I-23
I-24.	BCG Station 11 data summary for corals and subgroups	I-24
I-25.	Fish species found in BCG Station 11, with density and biomass for 100m ² transect	I-25
I-26.	BCG Station 12 data summary for corals and subgroups	I-26
I-27.	Fish species found in BCG Station 12, with density and biomass for 100m ² transect	I-27
J-1.	Stony corals included in Western Atlantic and Caribbean assessments with the three-letter identification code and the morphological conversion factor for calculating 3-D surface area	J-2
J-2.	Gorgonian morphological shapes, abbreviations, simulated model, <i>in situ</i> example and regression models to estimate three-dimensional surface area	J-6
J-3.	Sponge morphological shapes, abbreviations, simulated model, <i>in situ</i> example and regression models to estimate surface area	J-8

Acknowledgements

The US Environmental Protection Agency (EPA) Office of Research and Development (ORD) prepared this coral reefs report.

This is ORD tracking number ORD-007028, Atlantic Ecology Division, Narragansett, RI. EPA was supported in the development of this report by the Great Lakes Environmental Center, Inc., through contract EP-C-09-001.

The production of this report would not have been possible without the workshop participants.

Photos from: Charles LoBue, Peggy Harris, Mel Parsons, Alan Humphries, Scott Grossman, Richard Henry, Alina Szmant and Melanie McField.

The report was peer reviewed by Dr. Ku’ulei Rogers (University of Hawaii), Dr. Wendy Wiltse (US EPA, Region 9), Ms. Susan K. Jackson (US EPA, Office of Water), and Dr. Giancarlo Cicchetti and Dr. Marguerite (Peg) Pelletier (US EPA, Office of Research and Development).

We would like to thank the Caribbean Coral Reef Institute, and Drs. Richard Appeldoorn and Francisco Pagan for hosting the workshop and providing the conference facilities on Magueyes Island.

We would also like to thank Mr. Roberto Viquiera (Protectores de Cuencas) for assisting with the logistics, including providing the dormitory facilities in Yauco for visiting scientists.

Executive Summary

EPA's Office of Research and Development (ORD) and Office of Water (OW) hosted a workshop on coral reef biological integrity that brought together scientists with expertise in coral reef taxonomic groups (e.g., stony corals, fishes, sponges, gorgonians, algae, seagrasses and macroinvertebrates), specializing in community structure, organism condition, ecosystem function and ecosystem connectivity. The goals of this first workshop were to:

- Identify key qualitative and quantitative ecological characteristics (reef attributes) that determine the condition of linear coral reefs inhabiting shallow waters (<12 m) in southwestern Puerto Rico.
- Use those reef attributes to recommend categorical condition rankings for establishing a biological condition gradient.
- Ascertain through expert consensus those reef attributes that characterize biological integrity (a natural, fully-functioning system of organisms and communities) for coral reefs.
- Develop a conceptual, narrative model that describes how biological attributes of coral reefs change along a gradient of increasing anthropogenic (human-generated) stress.

The long-term goal is to derive scientifically defensible thresholds for different levels of coral reef condition that can be coupled with management objectives and used to evaluate alternative decision options.

The experts evaluated photos and videos for 12 stations collected in 2010 and 2011 during EPA coral reef surveys from Puerto Rico coral reefs exhibiting a wide range of conditions. The experts individually rated each station as to observed condition (good, fair or poor) and documented their rationale for the assignment. The group discussed the reef attributes that characterize biological integrity (or the natural condition) for Puerto Rico's coral reefs. These attributes will be further developed to characterize the baseline condition, an important concept for achieving Clean Water Act goals.

The attributes and thresholds will be organized into a conceptual, narrative model that describes how biological attributes of coral reefs change along a gradient of increasing anthropogenic stress. By providing the explicit characterization of how attributes of the biological system change as human disturbance increases, decision-makers will be able to use the narratives to determine which level most appropriately describes the current condition of their coral reefs and which level is the desired condition. From this, managers can set easily communicated, quantitative goals for achieving those conditions.

This is the first in a series of facilitated workshops and webinars with this group of coral reef experts.

Chapter 1. Introduction

The approach described in this report will assist in developing a conceptual, narrative model that describes how biological attributes of coral reefs change along a gradient of increasing anthropogenic stress. The framework is expected to serve multiple purposes.

- It will assist decision-makers in understanding the current conditions of the Puerto Rico coral reefs relative to natural, undisturbed conditions, the critical attributes of the coral reefs and how each attribute changes in response to stress. Through this framework, decision-makers can set realistic goals for their coral reefs and establish monitoring (measurement) endpoints that are meaningful based upon the attributes identified by the scientific community.
- It will be used to support the development of an economic survey of Puerto Rico's coral reefs (another project being conducted in collaboration with the National Oceanic and Atmospheric Administration's Office of National Marine Sanctuaries).
- It will inform the Bayesian Belief Network (BBN) and Dynamic Systems Models being developed by the EPA modelers.
- It will contribute to the development of coral reef biological criteria for water quality standards under the Clean Water Act (CWA) for Puerto Rico.

To initiate the process, scientists with expertise in coral reef taxonomy, ecology, and management of stony corals, fishes, sponges, gorgonians, algae, seagrasses, and macroinvertebrates, specializing in community structure, organism condition, ecosystem function and ecosystem connectivity were brought together. These experts participated in the first workshop, held August 21-22, 2012, in La Parguera, Puerto Rico. (See Appendix B for a list of workshop participants.)

The goals of this first workshop were to:

- Identify key qualitative and quantitative ecological characteristics (reef attributes) that determine the condition of linear coral reefs inhabiting shallow waters (< 12 m) in southwestern Puerto Rico.
- Use those reef attributes to recommend categorical condition rankings for establishing a biological condition gradient.
- Ascertain through expert consensus those reef attributes that characterize biological integrity (a natural, fully-functioning system of organisms and communities) for coral reefs.
- Develop a conceptual, narrative model that describes how biological attributes of coral reefs change along a gradient of increasing anthropogenic stress.

The long-term goal is to derive scientifically defensible thresholds for different levels of coral reef condition that can be coupled with management objectives and used to evaluate alternative decision options.

1.1 Coral Reef Ecosystems

Coral reefs are the earth's most biologically diverse marine ecosystems (Sebens 1994; Odum 1997). Scleractinian (stony) corals, octocorals and sponges provide structural habitat that supports harvestable fish species and attracts tourists (Bradley et al. 2008). Stony corals also protect shorelines from erosion by physically blocking current and wave energy (Wilkinson 1996), and coral reefs provide food and income for 500 million people globally (TNC 2006).

Corals are generally found in clear, shallow tropical oceans, and their growth is limited by temperature, salinity, light intensity, water clarity, and other chemical and water quality characteristics (Wells 1957; Brown and Howard 1985; Hubbard 1997; Ogden 1997; Hoegh-Guldberg 1999). Coral reefs are sensitive to relatively small changes in the environment (Richmond 1993) and their lack of resilience to environmental change has led some to regard coral reefs as sentinels of oceanic environmental quality (Hatcher et al. 1987; Andrews and Pickard 1990; Barber et al. 2001).

Healthy stony corals appear to be critical for fish productivity, species richness and fish biomass, all of which have been reported to decrease with a decline in stony coral health (Warren-Rhodes et al. 2003). Additionally, there appears to be a strong positive correlation of habitat complexity to the biodiversity and ecosystem functions of a reef community (Alvarez-Filip et al. 2009), including fish species richness (Walker et al. 2009; Pittman et al. 2007a, b). The rich diversity of coral reefs is partly dependent on the provision of habitable surface area and partly on the variability of that surface area (Principe et al. 2012).

The adjacent habitats of seagrass meadows and mangrove forests are linked with coral reefs to form a complex dynamic mosaic that provides critical nurseries, foraging areas, and refugia for fish and invertebrates (Christensen et al. 2003; Mumby et al. 2004, 2008; Aguilar-Perera and Appeldoorn 2007; McField and Kramer 2007; Meynecke et al. 2008). Mangroves and seagrasses can also trap sediments, nutrients, and pollutants, which can improve the water quality on nearby reefs (Grimsditch and Salm 2006). Many juvenile fishes occupy shallow-water habitats such as mangroves and seagrasses, while the adult forms are found in adjacent coral reefs (Nagelkerken et al. 2000; Adams et al. 2006; Cervený 2006; Dahlgren et al. 2006; Clark et al. 2009; Pittman et al. 2010).

1.2 Puerto Rico's Coral Reef Ecosystems

The US territory of Puerto Rico encompasses the main island of Puerto Rico, two inhabited islands (Culebra and Vieques) and three uninhabited islands (Mona, Monito and Desecheo). Puerto Rico has an estimated coastline of 930 km, a land area of 8,950 km² and fringing coral reefs with a total area of 3,370 km² off the east, south and west coasts (Wilkinson 2004; Burke and Maidens 2004).

The coral reef ecosystem in Puerto Rico is a complex mosaic of interrelated habitats, including mangrove forests, seagrass beds and coral reefs, as well as other coral communities (Garcia-Sais et al. 2008). Ballantine et al. (2008) listed 69 shallow-water (<40 m) scleractinian species, 260 fish species, 46 shallow-water alcyonarian species and 500 species of benthic marine algal flora, excluding cyanobacteria (Table 1-1).

Table 1-1. Number of currently reported species in each of the major marine taxa for Puerto Rico (adapted from Weil 2005).

Taxon	# Species	Source
Algae (diatoms; red, green, blue-green and brown algae)	492	Ballantine and Aponte 1997a, b; Ballantine and Aponte 2002
Mangroves	5	Cerame-Vivas 2001
Seagrasses (<i>Marine Phanerogams</i>)	7	Vicente 1992
Sponges (<i>Phylum Porifera</i>)	61	Wilson 1902; Weil 2005
Corals, anemones, jellyfish (<i>Phylum Cnidaria</i>)	171	Vaughan 1902; Hargitt and Rogers 1902; Almy and Carrion-Torres 1963; Garcia et al. 2003; Weil 2005
Unsegmented worms (<i>Phylum Nemertea</i>)	8	Coe 1902
Bivalves, snails, octopus, mollusks, nudibranchs (<i>Phylum Mollusca</i>)	1,176	Dall and Simpson 1902; Grana 1993; Ortiz 1998; Garcia-Rios 2003
Segmented worms, polychaetes (<i>Phylum Annelida</i>)	129	Treadwell 1902, 1939; Long 1975
Ostracods, crabs, shrimp (<i>Phylum Arthropoda</i>)	342	Benedict 1902; Bigelow 1902; Moore 1902; Rathbun 1902; Menzies and Glynn 1968
Starfish, sea urchins, brittle stars (<i>Phylum Echinodermata</i>)	165	Clark 1902, 1933
Bryozoans (<i>Phylum Ectoprocta</i>)	131	Osburn 1940
Fishes (Superclass Osteichthyes), sharks, rays (Class Chondrichthyes)	677	Dennis 2000; Dennis et al. 2004
Reptiles (turtles, snakes)	5	Rivero 1978
Mammals	18	Beller et al. 1999

While over 60 species of scleractinian corals inhabit the Western Caribbean, reefs in Puerto Rico were historically dominated by the reef-building coral taxa *Montastraea annularis* complex¹, *Agaricia agaricites*, *Montastraea cavernosa*, *Porites astreoides* and *Colpophyllia natans*. Additionally, *Acropora palmata* and *Acropora cervicornis* often formed dense, high-relief monospecific thickets; *A. palmata* in shallow exposed fore-reef habitats and *A. cervicornis* on fore-reefs and in shallow, protected back-reefs (Morelock et al. 2001).

Recent studies in Puerto Rico show that large corals of the genus *Montastraea* are critical for the biodiversity of fish and invertebrates and for maintaining the structure, function, and flow of reef ecosystem services (Beets and Friedlander 1998; Mumby et al. 2008). Mumby et al. (2008) found that one-fourth to one-third of benthic invertebrates and fish occurred in the *Montastraea*-dominated fore-reefs in the Caribbean. *A. palmata* and *A. cervicornis*, which have recently been listed as threatened species in the Caribbean, also significantly contribute to reef growth and development and provide essential fish habitat (NOAA 2012a; Principe et al. 2012).

¹ This report does not adopt the new classifications for the *Montastraea annularis* species complex (*Montastraea annularis*, *Montastraea faveolata* and *Montastraea franksi*) reclassified as the original genus *Orbicella* (Budd et al. 2012).

1.3 Timelines

1.3.1 Condition Timeline

- 1000:** Coral reefs would have been regarded as mostly pristine by current standards with healthy corals, large, well-structured fish and invertebrate communities, with probably only a depletion of some of the larger fauna (Wilkinson 2004).
- 1800:** Large vertebrates such as the green turtle, hawksbill turtle, manatee and Caribbean monk seal were decimated in the central and northern Caribbean Sea (Jackson 1997).
- 1880s:** Early taxonomic studies of reefs on Puerto Rico (e.g., mollusks [Gundlach 1883]; crustaceans [Gundlach 1887]; fishes [Poey 1881]; polyps, worms, fishes and crustaceans [Stahl 1883]; algae [Hauck 1888]; and coral [Vaughn 1902]).
- 1900:** Most coral reefs were healthy and dominated by healthy branching corals, urchins, large schools of game fish, sharks and algal grazers. Waters were clear with low nutrient levels (Wilkinson 2004).
- 1952:** The last confirmed sighting of the Caribbean monk seal was at Serranilla Bank between Jamaica and Nicaragua (Debrot 2000).
- Late 1950s and early 1960s:** Massive fishing pressure began in Puerto Rico (Appeldoorn personal communication). Herbivores and predators were reduced to very small fishes and sea urchins (Jackson 1997).
- 1969:** An intensive and extensive coral bleaching event occurred on coral reefs of southwestern Puerto Rico. The bleaching was probably caused by 38.1 cm of rain during a hurricane that preceded the bleaching (Williams and Bunkley-Williams 1990).
- Late 1970s:** Extensive thickets of *Acropora palmata* were present in 40% of locations surveyed around Puerto Rico; 20% of these reefs had dense *A. palmata* patches and abundant colonies of *A. cervicornis* (Weil et al. 2003).
- Late 1970s and early 1980s:** A white-band disease (WBD) epizootic event caused extensive mass mortality of Acroporid corals throughout their range in the Caribbean with losses up to 95% (Gladfelter 1982; Weil et al. 2003, 2009; Weil and Rogers 2011).
- 1981:** Minor but widespread bleaching caused by elevated sea surface temperatures (SST) occurred in southwestern and western Puerto Rico (Williams and Bunkley-Williams 1989, 1990).
- 1983:** *Diadema antillarum* mass mortality, with 85-100% population declines (Bak et al. 1983; Lessios et al. 1984; Lessios 1988, 2005; Osborne 2000). The *Diadema antillarum* mortality was first observed in Puerto Rico in January 1984 in the coral reefs off La Parguera (Vicente and Goenaga 1984b).

- Late 1980s:** Massive coral bleaching and mortality caused by elevated SST. Extensive partial coral colony mortalities and some total mortalities of coral reef organisms, including death of some 400-500 year-old coral colonies (Velazco-Dominguez et al. 2003; Burke and Maidens 2004). Massive coral bleaching events in Puerto Rico were first reported by Williams et al. 1987 and Goenega et al. 1989.
- 1990:** Severe bleaching in the western north Atlantic caused by elevated SST and doldrum surface waters (from Bermuda, Texas, Florida, throughout the Caribbean, south to Brazil). High mortalities of fire corals, scleractinian corals, gorgonians, sponges and other coral reef organisms (Velazco-Dominguez et al. 2003).
- 1994:** Cumulative impacts of disease, coastal development, coral bleaching and over-fishing have resulted in heavily damaged reefs. The more isolated reefs were in better condition because they were not affected by land-based stressors (Wilkinson 2004).
- 1996:** Caribbean yellow-band disease (YBD) first observed in Puerto Rico with very low prevalence (Bruckner and Bruckner 1997, 2006; Weil et al. 2009). The disease was highly seasonal (summer-fall). YBD affects three species of the former *Montastraea annularis* species-complex (*M. faveolata*, *M. annularis* and *M. franksi*), the most important reef-building corals for this area (Bruckner and Bruckner 2006; Cróquer and Weil 2009; Harvell et al. 2009).
- 1998:** Severe bleaching event in Puerto Rico caused by elevated SST (July–September); 99% of the colonies completely recovered after 9 months; 15% of the colonies bleached again in 1999 and recovered by January 2000 (Velazco-Dominguez et al. 2003).
- 2000:** *Diadema* seem to be making a slow return in many localities in the Caribbean, including La Parguera, PR (Weil et al. 2005).
- 2003:** YBD became chronic and colonies showed disease signs all year (Weil et al. 2009). Surveys of over 100 reefs along the coast and islands found that Acroporid populations continued to decline in some areas from persistent disease, storms and sedimentation coupled with the poor coastal environmental conditions (high turbidity, sub-optimal water quality, etc.) and algal overgrowth.
- 2004:** Most inshore reefs show advanced stages of degradation. *Montastraea annularis* species-complex was the dominant stony coral, but it was virtually absent on reefs with low coral cover. The encrusting octocoral *Erythropodium caribaeorum* occurred at most stations, and zoanthids (particularly the encrusting *Palythoa* species) and sponges were the dominant sessile benthic invertebrates in shallow waters. Macroalgae and turf algae were dominant instead of corals on most intermediate-depth reefs (Garcia-Sais et al. 2008).
- 2005:** A major bleaching event caused by elevated SST in the fall of 2005, followed in 2006 by mass cnidarian mortality, had a dramatic impact on Puerto Rican coral reefs. A total of 82 cnidarian species were impacted by the bleaching, including 52 scleractinians, 13 octocorals, four hydrocorals, four zoanthideans, four actiniarians, three corallimorpharians and two scyphozoans (Garcia-Sais et al. 2006, 2008). The most severe bleaching was observed among

Montastraea annularis species-complex (94%), *Helioseris cucullata* (94%), *Colpophyllia natans* (83%), *Siderastrea siderea* (65%), *Millepora* species (63%), *Mycetophyllia* species (2%), *Diploria* species² (54%), *Agaricia* species (48%) and *Montastraea cavernosa* (46%). Three genera appeared to be less susceptible to bleaching: *Eusmilia fastigiata* (22%), *Meandrina meandrites* (26%) and all *Porites* species (36%). *Millepora alcicornis* was completely bleached at all stations, and most colonies (>65%) had died by December 2005. In August 2006, most corals had regained normal coloration, with the exception of *Montastraea annularis* species colonies, which experienced extensive partial and full colony mortality throughout the region. Total coral cover declined throughout the region by 40-60%. Disease epizootics followed, including a white plague outbreak on the east and southern coasts, and Caribbean yellow-band disease (YBD) that primarily affected the *Montastraea annularis* species-complex that occurred right after the peak of the 2005 bleaching event (Garcia-Sais et al. 2008).

Intense bleaching of octocorals was first noted in late September to early October beginning with *Erythropodium caribaeorum*, followed by *Muricea*, *Briareum* and *Plexaurella* and later by *Pseudoplexaura* and *Pterogorgia* species. Bleaching in scleractinian corals, hydrocorals and the zoanthid *Palythoa caribaeorum*, preceded bleaching of octocorals, suggesting octocorals may have higher tolerance to thermal stress compared to the other major cnidarian taxa. By late November 2005, the majority of the affected octocoral colonies had not died. The exceptions were the bleached colonies of *Muricea*, which had 90% mortality (Prada et al. 2009).

- 2006:** NOAA's National Marine Fisheries Service (NMFS) listed *Acropora palmata* and *Acropora cervicornis* corals as threatened throughout their known range by authority of the Endangered Species Act (ESA). This designation became final in May 2006 (Federal Register 2006).
- 2008:** The Caribbean monk seal was officially declared extinct after an exhaustive five-year search by NOAA NMFS.
- 2014:** NOAA NMFS proposed seven Atlantic/Caribbean corals as endangered: *Acropora cervicornis*, *Acropora palmata*, *Dendrogyra cylindrus*, *Orbicella annularis*, *Orbicella faveolata*, *Orbicella franksi* and *Mycetophyllia ferox* and two species as threatened: *Agaricia lamarcki* and *Dichocoenia stokesii* (Brainard et al. 2011). In their final rule (August 27, 2014), NOAA listed *O. annularis*, *O. faveolata*, *O. franksi*, *D. cylindrus* and *M. ferox* as threatened species and determined that *D. stokesii* and *A. lamarcki* did not warrant listing. NOAA also determined that the listing of *Acropora cervicornis* and *Acropora palmata* as threatened in 2006 is still warranted.

² This report does not adopt the new classifications *Diploria strigosa* and *Diploria clivosa* as the genus *Pseudodiploria* (Budd et al. 2012).

1.3.2 Anthropogenic Activity Timeline

- ~2000-3000 B.C.:** The Ortoiroid people from the Orinoco region in South America settled in Puerto Rico. The Saladoid and Arawak Indians populated the island between 430 BC and 1000 AD. By 1000 AD the Taino culture was dominant (Rouse 1992). These early populations exploited coral reef resources, and there is strong archaeological evidence of major harvesting of fishes, molluscs, manatees and turtles (Wilkinson 2004).
- 1493:** Christopher Columbus landed in Puerto Rico, beginning an intense period of colonization and resource extraction (mainly gold). The early explorers found the indigenous population cultivating, blending, rolling and smoking tobacco. Europeans had never seen tobacco. This discovery marked the start of an international passion for "New World" tobacco and its much sought after byproduct, the cigar.
- 1496-1660:** Tobacco was the major crop. Half the shipping tonnage between Puerto Rico and Europe (mainly Spain) was comprised of tobacco.
- 1508:** Juan Ponce de Leon founded a town (*Guaynía*, meaning "*a place with water*") on the shores of Guánica Bay. The narrow channel and calm waters of Guánica Bay made it a natural refuge for ships sailing the Caribbean Sea.
- Early 1500s:** Sugar was introduced (perhaps when Juan Ponce de Leon began colonizing the island), and many small landowners relied on its export as a source of income.
- 1548:** Hundreds of sugar mills operated by waterpower. The industry was in the hands of small landowners whose enterprises succeeded or failed depending on the price of sugar in the market or the whims of the Spanish Crown.
- 1736:** Coffee plants introduced to Puerto Rico, grown mostly for personal and domestic use.
- Mid-1800s:** French immigrants from the Mediterranean island of Corsica settled around Yauco and became well known as premium coffee exporters to Europe.
- 1867:** (San Narciso), 1899 (San Ciricao), 1928 (San Felipe) and 1932 hurricanes virtually destroyed most coffee plantations and tobacco crops. Many farms never recovered.
- 1873:** First "Centrales" or sugar factories with equipment operated by steam were established. Centrifuges were used to separate the sugar crystals from the molasses.
- 1898:** Puerto Rico was ceded to US as a result of the Spanish American War. US markets opened to Puerto Rico products (tariff free). Sugar cane was the most important cash crop for the territory.
- 1900:** The US enacted the Foraker Act, which removed the previous land ownership cap of 500 acres. Large monoculture farming began (primarily sugar).
- 1901:** The South Porto Rico Sugar Company of New Jersey, USA, began construction of the Central Guánica sugar mill. The Central Guánica organized a company town around the sugar mill that included a hospital, school and housing facilities. This sugar mill was one of the largest in the

Caribbean and was one of the largest in the world until World War I (Ayala 1999; Wikipedia 2013).

1900–1927: Puerto Rico produced around 35 million pounds of tobacco a year. Tobacco represented 38% of the value of commercial crops in 1920 (sugar accounted for 25%). In 1910, 14% of farms reported the cultivation of coffee. 75% of the employed people in Puerto Rico were involved in the sugar industry controlled by US corporations (Miller and Lugo 2009).

Early 1930s: The Roosevelt Administration created the Puerto Rico Emergency Relief Administration, which became the Puerto Rican Reconstruction Administration in 1935. Rural resettlement communities and demonstration farms were established, and coffee and fruit production was reorganized.

1934: Jones-Costigan Act set a quota on the amount of sugar that could be exported to the US tariff free.

1940s: The Puerto Rico Water Resources Authority (PRWRA) initiated the Southwestern Puerto Rico Project (SWPRP). The SWPRP connected five watersheds and a retention pond through construction of dammed reservoirs and an underground aqueduct system that diverted water south into the Guánica Bay/Rio Loco watershed. This increased the watershed drainage area from approximately 57,000 to 97,000 acres.

1941: Land Reform Act was passed, limiting land ownership to 202 hectares (500 acres) or less. Many rural residents were now able to buy 10 hectare (25 acre) parcels, allowing them to grow crops for profit for both export and internal use. This broke up the land monopoly of the large sugar companies (Miller and Lugo 2009).

Late 1940s: The US Department of Agriculture (USDA) ended sugar subsidies for Puerto Rican farmers. Annual production of sugar dropped soon after. Massive numbers of Puerto Ricans migrated to the New York area.

1948: The Industrial Incentives Act (Operation Bootstrap) began to shift Puerto Rico from rural agriculture to more urbanized communities and industrial sources of income (shifts to manufacturing of pharmaceuticals, chemicals, machinery and electronics).

1948: The Guánica fertilizer plant opened, with storage silos and a shipping pier.

1952: Peak sugar production in Puerto Rico.

1953: Puerto Rico passed Act No. 65, authorizing the Lajas Valley Irrigation System.

1954: The Lajas Valley Irrigation (LVI) project, developed under the guidance of experts from the USDA led efforts to remove sugar plantations that had long characterized the region. This project also started small-scale agricultural farming and introduced cultivated fruits, mainly pineapple. The LVI project channelized 200,000 acres of land via 25 miles of a concrete-lined main canal, 60 miles of concrete-lined and unlined lateral canals and the corresponding drainage system. Two hydroelectric dams and two additional water reservoirs were built. Shade-grown coffee was transplanted into sun-exposed areas.

- 1955:** Guánica Lagoon was drained to increase land available for agriculture.
- 1970s:** Government subsidies and support for sun-grown coffee were implemented.
- 1981:** Guánica sugar mill was closed.
- 1980s:** Widespread conversion to sun-grown coffee reduced biodiversity (from loss of canopy habitat) and increased soil erosion from steep and now poorly vegetated, unprotected slopes. Soil washed from hillsides into streams and was trapped in the reservoirs, reducing their water storage capacity (Soler-López 2001) and increasing sediment deposition in Guánica Bay.
- 1999:** Puerto Rico passed an agricultural reserve law in an attempt to reverse the trend of declining agriculture.
- 2000:** The final two sugar factories closed (Coloso and Roig Centrales).
- 2009:** The US Coral Reef Task Force (USCRTF) selected the Guánica Bay watershed as the location for its first multi-agency watershed initiative in an attempt to reduce watershed impacts on coral reefs in the coastal zone.

1.4 Southwestern Puerto Rico

Southwestern Puerto Rico is relatively rural, with low population density compared to northeastern Puerto Rico. There are, however, some population centers: Yauco (population 20,295), San Germán (population 12,055), Guánica (population 9,224) and Sabana Grande (population 8,961). The human presence has created a variety of environmental stresses in this region.

Agricultural growth in southwestern Puerto Rico has resulted in widespread land clearing and modification. Nearly 90% of the area was deforested by the end of the 19th century (Warne et al. 2005), and the largest natural freshwater body in Puerto Rico, the Guánica Lagoon, was drained in 1955 as part of an agricultural development project (Sturm et al. 2012). These modifications have led to increased watershed sediment and nutrient yield, thereby increasing sediment and nutrient discharge to the coastal shelf (Warne et al. 2005).

Municipal growth has increased impervious cover, generation of stormwater runoff and human sewage. Impervious cover increases the loading of nutrients, bacteria, sediments and contaminants such as polycyclic aromatic hydrocarbons (PAHs), heavy metals and other pollutants associated with automobiles. Stormwater accumulates debris, chemicals, dirt and other pollutants, which are untreated and then discharged into coastal rivers and bays. Sewage carries pathogens that can transmit disease to humans and other animals, contains organic matter that can cause odor and nuisance problems and nutrients that can cause eutrophication of receiving water bodies. Much of the rural population in southwestern Puerto Rico relies upon septic systems for their sewage treatment. Too often these are failing, inadequate or improperly maintained. There are also several wastewater treatment plants (WWTP) in the area, which only treat the sewage to eliminate pathogens and solids. Some of these WWTPs are being upgraded to advanced secondary treatment, which provides minimal nutrient reduction.

Communities in the coastal region of southwestern Puerto Rico rely partially on fishing and tourism for their livelihood. Fishing, if not conducted in a sustainable manner, can lead to overexploitation of marine living resources (both target species and the marine system as a whole). Boat anchors, traffic and groundings can adversely impact marine resources. For example, much of the seagrass in the shallow shelf area near La Parguera is vulnerable to damage from boat propellers. Lost fishing gear, such as hooks, lines, nets and lobster traps, can also be damaging to marine resources.

Finally, elevated SSTs are correlated with mass bleaching events (Goreau et al. 1992; Glynn 1988, 1991; Hoegh-Guldberg 1999; McClanahan et al. 2007; Meissner et al. 2012). Sea surface temperatures have been higher during the past three decades than at any other time since reliable observations began in 1880 (NOAA 2012b). Global warming is caused by human activities that emit heat-trapping carbon dioxide and result in increased SSTs. A summary table illustrating some of the major stressors and their sources is shown in Table 1-2.

Table 1-2. Complex relationships between stressors exist in southwestern Puerto Rico.

Source of Stressor	Stressors
Agriculture	Increased sediment, nutrient, pesticide and herbicide loads to aquatic ecosystems
Urban development	Increased sewage (nutrients and pathogens), stormwater runoff (sediment, contaminants)
Fishing	Overexploitation of fish populations; by-catch; damage from fishing gear and boats
Increased global CO ₂ emissions from power generation and transportation	Elevated sea surface temperature and acidification in the marine environment

1.5 The Clean Water Act

The Clean Water Act (CWA) (33 U.S.C. § 1251 et seq. 1972) is the cornerstone for surface water quality protection in the United States. The CWA objective is to restore and maintain the chemical, physical and biological integrity of the Nation's waters. The CWA authorizes EPA to determine "appropriate minimum levels" of protection and to provide oversight to states (states, territories and commonwealths), which are required to establish water quality standards that define the goals (designated uses) and pollution limits (water quality criteria) for all waters within their jurisdictions. States are also required to monitor conditions regularly and submit biannual reports summarizing water quality assessments. Waterbodies that do not meet the water quality criteria are reported as "impaired", triggering a series of management actions to determine the cause of impairment and to restore the condition of the waterbody and its resident biota.

Biological integrity is a long-term objective of the CWA, and water quality standards and criteria can be defined to protect valued aquatic resources, such as coral reef ecosystems. Biological assessments directly measure the condition of the aquatic resource to be protected and the cumulative response of the biological community to all sources of stress. Biological standards (biocriteria) set biological quality goals.

The President's Ocean Action Plan (US Commission on Ocean Policy 2004) required EPA to develop the tools and knowledge necessary to protect coral reefs from land-based pollution using coral reef biological criteria. A comprehensive guide (Bradley et al. 2010) describes the process for using the CWA and biological criteria to enhance coral reef protection efforts.

1.6 Why a Coral Reef Ecosystem Conceptual Model is Needed

Coral reef condition typically degrades as human disturbance increases. Human disturbances threatening coral reefs include polluted runoff from agriculture and land-use practices, over-fishing, ship groundings, coastal development, sewage discharge and climate change. Natural stressors such as tropical storms can also adversely impact coral reefs. Both natural and anthropogenic stressors can cause increases in coral bleaching and diseases. Reefs in the US Caribbean have declined from 50% total coral cover to less than 10% in just 25 years (Wilkinson 2004).

The biological communities of the coral reef reflect overall ecological integrity (i.e., chemical, physical and biological integrity), integrate effects of multiple stressors and provide a measure of aggregate impact (Barbour et al. 1999). Coastal resource managers and coral reef scientists routinely conduct biological assessments to evaluate the condition of coral reefs. This approach integrates the cumulative impacts of chemical, physical and biological stressors on aquatic life. However, while the stated intent of these biological assessments is to support decision-making, they more commonly document the decline of the coral reefs. A missing component in this approach is a scientifically derived process for identifying thresholds that can be coupled with management objectives and used to evaluate alternative decision options. A conceptual model can help to organize information and make sense of system components and their interactions.

1.7 The Framework: The Biological Condition Gradient (BCG)

Beginning in the late 1990s, EPA collaborated with aquatic scientists and managers across the United States to develop and implement the Biological Condition Gradient (BCG) for freshwater streams (Davies and Jackson 2006). The BCG is a conceptual model that describes how biological attributes of aquatic ecosystems (i.e., biological condition) might change along a gradient of increasing anthropogenic stress (e.g., physical, chemical and biological impacts). The BCG was designed to provide a means to map different indicators on a common scale of biological condition to facilitate comparisons between programs and across jurisdictional boundaries in context of the CWA.

Since then, many states have used the BCG to support water quality management, and several states have used the BCG to more precisely define freshwater stream designated aquatic life uses, identify impairment thresholds, monitor status and trends and track progress in restoration and protection (EPA 2011). Additionally, stream BCGs have been developed at the regional and local government scale throughout the US.

Since 2008 EPA has been collaborating with estuarine scientists and managers to adapt the stream BCG framework to more complex estuarine waterbodies (EPA in review). Estuarine BCG pilot work has focused on National Estuary Program (NEP) sites: Narragansett Bay (RI), Casco Bay (ME), Mobile

Bay (AL) and Tampa Bay (FL). NEPs play an important role as conveners of technical, management and public interests. Their ability to create connections among these constituencies makes them a valuable platform to work out the complexities of an estuarine BCG at different scales.

A BCG calibrated with field data can help states more precisely define biological expectations for their designated aquatic life uses, interpret current condition relative to CWA objectives and goals, track biological community responses to management actions and communicate environmental outcomes to the public. The model can serve as a template for organizing field data (biological, chemical, physical, landscape) at an eco-regional, basin, watershed or waterbody scale. It provides a framework for understanding current conditions relative to natural, undisturbed conditions.

In practice, the BCG is used to first identify the critical attributes of an aquatic community and then to describe how each attribute changes in response to stress. Coral reef managers can use the BCG to interpret biological condition along a standardized gradient, regardless of assessment method, and apply that information to different programs.

The BCG model provides a framework to help water quality managers do the following:

- Decide what environmental conditions are desired (goal-setting)—The BCG can provide a framework for organizing data and information and for setting achievable goals for waterbodies relative to natural conditions (e.g., condition comparable or close to undisturbed or minimally disturbed condition).
- Interpret the environmental conditions that exist (monitoring and assessment)—Practitioners can get a more accurate picture of current waterbody conditions.
- Plan for how to achieve the desired conditions and measure effectiveness of restoration—The BCG framework offers water program managers a way to help evaluate the effects of stressors on a waterbody, select management measures by which to alleviate those stresses and measure the effectiveness of management actions (EPA 2011: Case Example 3.16).
- Communicate with stakeholders—When biological and stress information is presented in this framework, it is easier for the public to understand the status of the aquatic resources relative to what high-quality places exist and what might have been lost.

1.7.1 How is the BCG Constructed?

The BCG has been divided into six levels of biological conditions along the stressor-response curve, ranging from observable biological conditions found at no or very low levels of stress (level 1) to those found at high levels of stress (level 6) (Figure 1-1).

The BCG was developed to serve as a scientific framework to synthesize expert knowledge with empirical observations and develop testable hypotheses on the response of aquatic biota to increasing levels of stress.

It is intended to support more consistent interpretations of the response of aquatic biota to stressors and to clearly communicate this information to the public. It is being evaluated and piloted in several regions and states.

Davies and Jackson (2006) provides a description of how 10 attributes of aquatic ecosystems change in response to increasing levels of stressors along the gradient, from level 1 to 6. The attributes include several aspects of community structure, organism condition, ecosystem function, spatial and temporal extent and connectivity (Table 1-3).




Figure 1-1. The Biological Condition Gradient (BCG) (modified from Davies and Jackson 2006).

Table 1-3. Biological and other ecological attributes used to characterize the freshwater streams Biological Condition Gradient (BCG) (Modified from Davies and Jackson 2006).

Attribute	Description
I. Historically documented, sensitive, long-lived, or regionally endemic taxa	Taxa known to have been supported according to historical, museum or archeological records, or taxa with restricted distribution (occurring only in a locale as opposed to a region), often due to unique life history requirements (e.g., Sturgeon, American Eel, Pupfish, Unionid mussel species).
II. Highly sensitive (typically uncommon) taxa	Taxa that are highly sensitive to pollution or anthropogenic disturbance. Tend to occur in low numbers, and many taxa are specialists for habitats and food type. These are the first to disappear with disturbance or pollution (e.g., most stoneflies, Brook Trout [in the east], Brook Lamprey).
III. Intermediate sensitive and common taxa	Common taxa that are ubiquitous and abundant in relatively undisturbed conditions but are sensitive to anthropogenic disturbance/pollution. They have a broader range of tolerance than Attribute II taxa and can be found at reduced density and richness in moderately disturbed stations (e.g., many mayflies, many Darter fish species).
IV. Taxa of intermediate tolerance	Ubiquitous and common taxa that can be found under almost any conditions, from undisturbed to highly stressed stations. They are broadly tolerant but often decline under extreme conditions (e.g., filter-feeding caddisflies, many midges, many Minnow species).
V. Highly tolerant taxa	Taxa that typically are uncommon and of low abundance in undisturbed conditions but that increase in abundance in disturbed stations. Opportunistic species able to exploit resources in disturbed stations. These are the last survivors (e.g., tubificid worms, Black Bullhead).
VI. Non-native or intentionally introduced species	Any species not native to the ecosystem (e.g., Asiatic clam, zebra mussel, Carp, European Brown Trout). Additionally, there are many fish native to one part of North America that have been introduced elsewhere.
VII. Organism condition	Anomalies of the organisms; indicators of individual health (e.g., deformities, lesions, tumors).
VIII. Ecosystem function	Processes performed by ecosystems, including primary and secondary production; respiration; nutrient cycling; decomposition; their proportion/dominance; and what components of the system carry the dominant functions, for example, shift of lakes and estuaries to phytoplankton production and microbial decomposition under disturbance and eutrophication.
IX. Spatial and temporal extent of detrimental effects	The spatial and temporal extent of cumulative adverse effects of stressors; for example, groundwater pumping in Kansas resulting in change of fish composition from fluvial dependent to sunfish.
X. Ecosystem connectivity	Access or linkage (in space/time) to materials, locations and conditions required for maintenance of interacting populations of aquatic life; the opposite of fragmentation. For example, levees restrict connections between flowing water and floodplain nutrient sinks (disrupt function); dams impede fish migration, spawning.

Each attribute provides some information about the biological condition of a waterbody. Combined into a conceptual model like the BCG, the attributes can offer a more complete understanding of current waterbody conditions and also provide a basis for comparison with naturally expected waterbody conditions. All states and tribes that have applied a BCG used the first five attributes that describe the composition and structure of the biotic community on the basis of the tolerance of species to stressors. Where available, states and tribes also included information on the presence or absence of native and non-native species for fish and amphibians, as well as observations of overall health and condition (e.g., size, weight, abnormalities, tumors and diseases).

The last three BCG attributes (ecosystem function, spatial and temporal extent of detrimental effects and ecosystem connectivity) can provide valuable information when evaluating the potential for a waterbody to be protected or restored. Several of EPA's NEPs, in conjunction with EPA ORD, are exploring application of those attributes at a whole-estuary scale (e.g., distribution and connectivity of critical aquatic habitats and associated biota).

Additionally, individual attributes might uniquely respond to a specific stressor or group of associated stressors (biological response signatures) (Yoder and Rankin 1995 a, b; Yoder and Deshon 2003). That information could contribute to the causal analysis of biological impairment, Stressor Identification (SI) and Causal Analysis/Diagnosis Decision Information System (CADDIS) (<http://www.epa.gov/caddis/>).

Currently, applications of the BCG that include development of a BCG-index (BCG-I) and incorporation of the BCG in a state's water quality management have been used only on freshwater streams. More recently, ongoing pilot efforts at several NEPs are extending the BCG concept to assessment and management of estuaries. Efforts to develop an estuarine conceptual model have focused on five attributes (structure, condition, function, connectivity and non-native species [Cicchetti 2010]) at scales ranging from the whole estuary to the single habitat, or biotope (e.g., seagrass beds, salt marshes and clam flats) (Table 1-4). This multi-scale approach is intended to improve restoration and management efforts. At larger scales, managers can prioritize and develop programs to restore the historic balance of critical habitats (biotopes) relative to an undisturbed historic benchmark, while also targeting restoration of all living habitats, to the maximum extent possible. The single habitat scale is assessed using biological assessments, which enjoy an established history within management approaches (Cicchetti and Greening 2011; EPA 2011).

Extending the BCG conceptual model to new waterbodies (coastal waters) and new assemblages (coral reef communities) is a multistep process. A successful process assembles a workgroup of experts on the habitats and assemblages and elicits from these experts: descriptions of the native aquatic assemblages under natural conditions, identifications of the predominant regional stressors and descriptions of degradation levels corresponding to the BCG. Descriptions should include the theoretical foundation and observed assemblage responses to stressors. In addition to expert opinion, the process makes use of empirical monitoring data. During a workshop, experts familiar with local conditions use the data to define the ecological attributes and set narrative statements. The experts determine narrative decision rules for assigning stations to a BCG level on the basis of

the biological information collected at the stations. Further development of quantitative decision rules and a quantitative BCG is more involved and requires a greater time commitment from the expert panel to participate in iterative calibration steps and review of more extensive monitoring data.

Table 1-4. Ecological attributes used to characterize the estuarine Biological Condition Gradient (BCG) (EPA in review).

Attributes	Potential Metrics and Description
Structure and Compositional Complexity	Community or habitat structure and complexity. May also recognize loss of habitats or species due to human activities. Examples include macroinvertebrate or fish indices, phytoplankton or zooplankton community measures, epifaunal measures, biotope mosaics, presence/quality of sensitive or susceptible taxa or biotopes, wetland vegetative indices, etc.
Condition	Measures condition of the waterbody, habitat or species. Also includes measures of resiliency. Examples include harmful algal blooms, disease outbreaks (locally or system-wide), measure of habitat or biotope health, such as seagrass condition or wetland condition, fish pathology or shellfish bed condition.
Function	Measures of energy flow, trophic linkages and material cycling. They may include proxy or snapshot metrics that correlate to functional measures. Examples include photosynthesis/respiration ratios, benthic: pelagic production rates, chlorophyll <i>a</i> concentrations and macroalgal biomass.
Connectivity	Metrics of exchange or migrations of biota between adjacent waterbodies or habitats. Important measures within the area being studied may be strongly affected by factors adjacent to or larger than the immediate study area. Proxies may need to be used as metrics. These may include linkages, fragmentation or hydrological measures.
Non-native Taxa	Metrics of non-native species. May include measures of the impact of invasive species and non-natives. Examples include estimated numbers of species or individuals, biomass measures of natives and non-natives or replacement of native species.

This report communicates the first results to apply the BCG conceptual model to the assessment of the condition of coral reefs. The first stage reported here is a proof-of-concept to introduce the conceptual model to coral reef experts and elicit a preliminary set of narrative decision rules for assigning coral reef stations to levels of the BCG. If the conceptual model passes muster among the experts, it will allow identification of the steps needed to develop and implement more comprehensive quantitative decision models.

Chapter 2. Approach

The coral reef BCG will be developed through a series of facilitated expert workshops. In the workshops, the experts assess the condition of coral reef sites based on biological data collected at the sites (species composition, abundance, health) and assign each site to a condition category (level) of the BCG. The experts' reasoning for making assignments are developed into a set of decision rules, at first qualitative, but through iteration, increasingly quantitative. The expert-derived rules are translated to a quantitative decision algorithm, in this case using mathematical set theory (e.g., Droesen 1996). The decision algorithm allows independent assessments with results comparable to those of the expert panel. Furthermore, the decision rules are documented so that modifications can be made as information and needs change.

The participants for the initial workshop were invited based on their scientific knowledge of the coral reefs and reef organisms of Puerto Rico. As a first step, participants were asked to evaluate and rank coral reef condition from photos, videos and data collected during EPA's 2010 and 2011 coral reef assessment surveys in shallow waters (<12 m deep) of southwestern Puerto Rico. The biological assemblages considered were stony corals, fishes, sponges, gorgonians and benthic macroinvertebrates. Rugosity, a reef-scale indicator of reef complexity, was determined using a chain-transect method that compares the six-foot length of a chain draped along the top of corals and along the bottom of the reef to the length of a taut line across the same linear distance. Participants were asked to share videos and pictures of reefs from the present or past that they believed exhibited full biological integrity.

A unique aspect of this workshop was that participants were reacting to the visual imagery of the reefs and evaluating different levels of coral reef condition. Participants moved from a visual, simple approach to more complex data-driven analysis. The workshop was designed to encourage brainstorming, facilitate discussion and not get mired in EPA terminology or definitions at the beginning of the workshop. Participants examined the visual media, rated the condition of various coral reefs and provided rationale for their ratings. Descriptions of good and bad characteristics relative to ecological condition were captured by facilitated discussions. A preliminary list was generated describing attributes that would characterize a coral reef with high (minimally disturbed or reference) or good condition. A minimally disturbed condition provides a fixed point in time and can help us to avoid problems associated with shifting baselines (Pauly 1995; Stoddard et al. 2006). A firm concept of minimally disturbed anchors ecological condition as a reference and helps us deal with changes (e.g., climate change), which broadly affect conditions that occur after the anchored point. Further, a clear picture of minimally disturbed provides a basis for effective public communication of changes over time and can provide a reference point for certain indices of ecological condition.

The workshop was designed to last three days; however Day 3 was cancelled because of Tropical Storm Isaac. Certain planned activities from Day 3 were shifted into Day 2, and the remainder of workshop information was communicated during webinars.

2.1 Video and Photo Evaluations

On the first day, panelists were asked to view and rate the coral reef condition of 12 EPA stations on the south shore of Puerto Rico (Table 2-1).

Table 2-1. Shallow inshore linear reefs used for BCG workshop stations that correspond to US EPA stations sampled along the southern coast of Puerto Rico.

Station	Year	EPA station	Latitude	Longitude	Depth (ft.)
1	2010	PR_2010_125	17.92486	-66.20363	20
2	2011	PR_2011_15	17.98198	-66.77228	23
3	2011	PR_2011_113	17.95942	-67.03902	21
4	2011	PR_2011_03	17.94420	-66.91638	21
5	2011	PR_2011_19	17.94180	-66.88060	12
6	2010	PR_2010_14	17.93875	-67.10927	27
7	2010	PR_2010_16	17.93922	-67.06197	24
8	2010	PR_2010_108	17.94085	-67.07708	12
9	2010	PR_2010_109	17.95373	-67.05012	16
10	2011	PR_2011_01	17.96380	-67.04980	11
11	2011	PR_2011_46	17.93418	-67.10108	17
12	2011	PR_2011_25	17.93670	-66.88660	17

Stations were limited to shallow (<12 m), near-shore (less than 3 nautical miles from shore) linear reefs as designated by NOAA benthic maps (Kendall et al. 2001), see Figure 2-1.




Figure 2-1. Map with locations of the 12 EPA stations along the southern coast of Puerto Rico.

Videos of eight stations were examined in the morning and four stations in the afternoon. Videos were displayed on four computers looping two stations per video in the morning and one station per video in the afternoon. Each station presented video footage from both panoramic and linear transect views, which began with 15 seconds showing the station number and the type of footage, to allow reviewers time to prepare for each video. A 60-second period between each station allowed panelists time to complete their evaluations before the next station video began. In addition to the video clips (ranging from 30 seconds to 3 minutes long), eight still photographs for each station (Appendix G) were provided in a notebook to supplement the videos and in particular, to capture aspects of the station not represented in the videos (e.g., fish).

The panelists were instructed to draw upon their overall personal experience and expertise to rate the reef condition for each station as either good, fair or poor based on what they viewed in videos and photos. Workshop binders organized by station included a photo diary of key representative photos, two ballots to rate the stations for each session (Appendices D and E) and note sheets (Appendix F) to document the traits or characteristics that panelists used to support their ratings. The panelists were asked to consider all aspects of the reef and specifically instructed to consider the characteristics of the condition of corals, sponges, gorgonians, fish, algae, reef rugosity and topographical heterogeneity. The facilitators suggested that panelists not compare ratings with each other, but panelists were free to discuss and view videos as a group. The panelists were not asked to rate any specific number of stations as good, fair or poor, but had free rein as they circulated around and viewed the video loops at their own pace. The facilitators were available if panelists had questions.

The panelists asked about shifting baseline conditions, and they were encouraged to draw upon their personal experience. Panelists also asked if each assemblage should be rated separately. Facilitators responded that it was all right to consider and document each assemblage, but they must finally select one single rating for each station. When all stations were evaluated, panelists recorded their individual ratings on the ballots and returned them to the facilitators. Note sheets were not collected—they were for each panelist to reference during subsequent discussions.

After ballots were collected from the panelists, the facilitators tabulated all the scores by ranking the stations in order from best condition to worst using a weighted ranking system (Table 2-2). Each good vote was given a value of 10 points, each fair vote 5 points and each poor vote 1 point. Table 2-3 shows the ranking given by each expert after visually evaluating videos and photographs for 12 EPA stations.

The panelists provided feedback on how to improve the process used to evaluate the stations. Many found it challenging to assess reef condition based on the video quality, which was raw footage from inexpensive video-enabled digital cameras. Still photos provided the best way to document fish populations, because many fishes were disturbed during the assessments that occurred before the video was taken. The fish surveyors were the first team to perform visual assessments, followed by

Table 2-2. Coral reef condition evaluated by the experts. Results were obtained from 20 coral reef experts after viewing videos and photos of 12 stations, with the highest score bolded. Overall rank is established by weighted ranking.

Station No.	No. Good	% Good	No. Fair	% Fair	No. Poor	% Poor	Majority Rating	Weighted Score	Overall Rank
1	1	5	9	45	10	50	Poor/Fair	65	8
2	0	0	0	0	20	100	Poor	20	12
3	13	65	7	35	0	0	Good	165	1
4	0	0	2	10	18	90	Poor	28	11
5	0	0	10	50	10	50	Fair/Poor	60	9
6	0	0	6	30	14	70	Poor	44	10
7	4	20	16	80	0	0	Fair	120	4
8	0	0	15	75	5	25	Fair	80	6
9	1	5	12	60	7	35	Fair	77	7
10	5	25	14	70	1	5	Fair	121	3
11	2	10	17	85	1	5	Fair	106	5
12	6	30	12	60	2	10	Fair	122	2

Table 2-3. Rankings given by experts after visually evaluating videos and photographs for 12 EPA stations. Abbreviations: G rated good; F rated fair; P rated poor. Some ratings were intermediate between two classes.

Expert	Station Number											
	1	2	3	4	5	6	7	8	9	10	11	12
Appeldoorn	P	P	G	P	F	P	F	F	F	F	F	F
Ballantine	F	P	G	F	F/P	P	F	F	G	G	G	G
Bauzá	F	P	G	P	F	P	G	F	F	F	F	F
Canals	F	P	F/G	P	P/F	P	F	F	P	F	F	F/G
Cuevas	F	P	G	P	F	P	F	F	P/F	F	F	F
Díaz	F/G	P	F/G	P	F/P	F/P	F/G	F/P	F/P	G	F/P	F
Fisher	P	P	G	P	P	P	G	F	F	G	F	G
Hutchins	P	P	F	P	P	P	F	P	P	F	F	F
McField	P/F	P-	F	P	F	P	F	P/F	P+	F+	F	P+
Miller	F	P	G	P	P/F	P	F	P/F	F	G/F	F	G
Pagan	F/P	P-	G/F	P	P	F	F	F	P	F	F	F
Ramos	F	P	G	P	P	P	F	P	F	P	F	F
Roberson	P	P	F/P	P	P/F	F/P	F	P	F/P	F/P	P/F	F
Ruiz	F	P	G	P	F	F	G	F	F	G	G	F
Sabat	P	P	G	P	P/F	P/F	F	F	P	F/G	F	F
Smith	P	P	F	P	P/F	P/F	F	F	P	F	F	F
Szmant	P	P-	G	P	F	P	F	F	F/P	F	F/G	G
Todd	P	P	G	P	P	P	G	F	F	F	F	G
Vicente	P	P	F	P	F	F	F	F	F	F	F	F
Yoshioka	G/F	P	G	F	F/G	F	F/G	F	F/P	F/G	F	G/F

the surveyors assessing other assemblages, including those photographing and video recording the reef. The photographers began potentially after many fishes had been scared away. The panelists also requested more details on the stations they were evaluating. Panelists felt they needed to know the location of the reef, what specific reef habitat they were viewing, the depth, the wave exposure and other features. They also wanted to know what stressors the station may have experienced. Recommendations for the future included upgrading video camera quality, standardizing the videography approach and providing more information to get a broader perspective of each station and its surrounding reef.

A facilitated discussion followed on the attributes that the panelists had identified to justify their ratings. The stations rated best and worst were considered first, with each panelist's comments captured on flip charts. Panelists were encouraged to edit posted pages of their comments if they felt their thoughts were not accurately captured. All participants were given the opportunity to submit comments on index cards if they wished to provide additional comments privately or anonymously. A summary of the characteristics considered by the panelists in rating the stations is provided in Section 2.2 for the best, worst and several fair stations.

2.2 Summary of Ratings

All of the photos given to the experts are found in Appendix G, ordered by station number.

2.2.1 Best Station, Ranked #1

The experts rated the condition of Station 3 as the best of the 12. It was rated good by 65% of the experts and fair by 35%. No expert rated it as poor. The experts were asked what characteristics or attributes were present that caused them to rate it as the best station. Their responses follow:

- Abundance of *Montastraea annularis* species-complex was high, with low partial mortality of tissue and large colony sizes indicative of older, mature coral colonies.
- Reefs showed high structural complexity, surface heterogeneity and high rugosity or presence of three-dimensional structures, allowing better reef development than would a flat topography.
- Stony coral biodiversity was moderate and included *Colpophyllia natans*, *Siderastrea siderea* and *Porites astreoides*, as expected on near-shore linear reefs in southwestern Puerto Rico.
- The water column showed high clarity with no visible sediment; experts also noted a lack of siltation or films covering the substrate.
- Coralline algae were more abundant than brown algae, and *Dictyota* was rare or absent.
- Gorgonian coverage was high, with most sea fans intact and uninjured. *Diadema* was present. Stony coral colonies had no or few boring clionid sponges.
- Damselfish were seen and the presence of additional fish species contributed to a good rating.

The experts, who rated the station as fair, expressed concern about the uncertainty in identifying fish species because the visual media were not adequate to consider fish size distributions or trophic status. A few grazing fish species were seen, but not enough to alleviate concerns of low grazing potential. Another expert cited the presence of coral disease. Finally, sponge abundance and diversity were low, with an absence of arborescent, vase and barrel morphologies, which are dominant in high quality habitat.

- The experts were also asked what attributes were absent that caused them to not rate the station as excellent. Their responses follow:
- The station had lower than expected diversity of stony corals, fishes and sponges, with little evidence of any recruitment.
- Very few anemones and invertebrates were observed, again indicating low species diversity.
- One expert stated that sponges, because they are efficient filter feeders, might be one of the assemblages most sensitive to chemicals in the water column and could act as an indicator species.

2.2.2 Worst Station, Ranked #12

All the experts agreed that Station 2 was in the worst condition, and rated it poor. This station was characterized by:

- High sedimentation and turbidity in the water column, which appeared as large patches of flocculent material creating low visibility, which the experts judged to represent low water quality.
- The reef colors were drab brown or green.
- Thick goopy sediment (probably of terrigenous origin) covered most of the bottom and organisms living on the bottom. Exposed hard substrate was absent, with no clear surfaces for attachment or recruitment.
- Algal cover was high, with lots of *Dictyota* and cyanobacteria as evidenced by a slimy appearance with a “skuzzy fuzzy” texture.
- Abundance of coralline algae was low.
- The absence of reef relief was coupled with low rugosity and no or very few small and live coral colonies.
- No fish or gorgonians were observed, although a few *Diadema* were noted.
- Sponge morphologies were ropy or encrusting, indicative of poor habitat and water quality.
- Heterotrophic sponges dominated, with a high abundance of filter feeders and no apparent autotrophic sponges. This is a characteristic typical of highly silted areas.

2.2.3 Stations Rated Fair

Station 8 ranked #6. This station was ranked as the most centric station among those with a fair rating. 75% of the experts rated it fair while 25% rated it poor. This station had the highest coverage of *Acropora palmata* in all stations viewed, although overall coral cover was low, with lots of coral rubble. *A. palmata* colonies varied greatly in size, condition and distribution and were present in about 25% of the transect area. Some very mature and large colonies were present in varying condition as evidenced by the amount of healthy coral tissue. Many of the *A. palmata* colonies lacked significant tissue and showed characteristics ranging from signs of tissue recovery, partial tissue mortality from white-band disease, lesions and white denuded tips, perhaps from fish predation. Some standing colonies were completely dead. The coral rubble on the bottom was composed of many broken and dead pieces of *A. palmata* skeletons, but finer sediments were absent. Several experts commented that the clean substrate and unconsolidated rubble showed significant and recent hurricane damage. Although some reef was dead, it showed signs of recovery and resiliency with the persistence of corals.

The clean substrate provided suitable areas available for settlement and recruitment of corals and other sessile invertebrates. Some coralline algae were present but overall algal diversity was low, with some fleshy algae. Much of the substrate appeared as though it had been highly grazed by *Diadema antillarum*. *Palythoa*, often considered an emerging opportunistic species, was prevalent throughout the transect colonizing dead coral skeletons. Several species of fish swam in large schools representing a decent diversity, including evidence of herbivores. The primary sources of rugosity were the *A. palmata* colonies; most other coral colonies showed relatively low relief. Few invertebrates other than those already mentioned were observed, with the exception of some small sea fans and low relief gorgonians.

Station 5 ranked #9. Half of the experts rated it fair and the other half rated it poor. This station shared many attributes with Station 8, but was judged to be in poorer condition because of higher sediment and turbidity, together with lower coral cover and diversity. The transect video showed substantial *A. palmata* coverage, but the colonies appeared to be in poorer condition than those seen in Station 8. One expert described the station as a “beat up Apal zone” (*A. palmata*) with large rubble between colonies. Thicker and larger algal turf patches and more sponges were present in comparison to Station 8. Parrotfish biting scars and scrapes were observed. However, it was noted that sedimentation, turbidity and water quality can vary with year, season and time of day, so the apparent condition could be extremely variable and dependent on when the video was taken.

Station 10 ranked #3. This station was rated as good by 25% of experts, fair by 70% of experts and poor by 5% of experts. Many of the features seen in the previous fair descriptions were also present here. A novel observation was the presence of the boring sea urchin *Echinometra*, which bioerodes coral skeletons; the herbivorous urchin *Diadema* is usually considered an indicator of better condition.

2.2.4 Station Rated Poor

Station 6 ranked #10. This station was rated fair by 30% of experts and poor by 70% of experts. There was evidence of significant bioerosion on the reef surfaces, perhaps due to boring clionids and the encrusting sponge *Chondrilla nucula*. One expert asked if this was a hard bottom station (not a coral reef). Despite the poor condition of Station 6, all experts agreed that Station 2 was in the poorest condition.

2.3 Summary of Attributes

Attributes developed by the experts were assembled into a list. In a facilitated discussion, the experts reached consensus about which direction (increase or decrease) the attribute would go at a station with improving condition, and what types of measurements or sub-attributes would be important. This information is summarized in Table 2-4.

Table 2-4. Summary of attributes and their relationships for assessing coral reef condition from station evaluations.

Attributes of good stations		
Direction with improving condition	Attribute	Sub-attribute/measurements
increase	3D structure	rugosity, cover
increase	stony coral abundance	<i>Montastraea annularis</i> complex, <i>Acropora palmata</i> , <i>Acropora cervicornis</i> , <i>Diploria strigosa</i> , large stony corals
increase	stony coral condition	% live tissue, absence of disease
increase	stony coral diversity	high number of stony coral species
increase	stony coral population structure	large colonies
increase	stony coral recruitment	
decrease	dominance of weedy, tolerant species	<i>Colpophyllia natans</i> , <i>Siderastrea siderea</i> , <i>Porites astreoides</i>
increase	coralline algae	
decrease	zoanthids	<i>Palythoa</i> species
decrease	exotic species	exotic fish, corals
decrease	filter feeders	heterotrophic sponges
increase	fish abundance	
increase	balance in fish population size and structure	
increase	fish biomass	
increase	balance in fish trophic structure	
increase	fish diversity	
decrease	fleshy algae	
increase	gorgonian abundance	

Table 2.4 (continued) Attributes of good stations		
Direction with improving condition	Attribute	Sub-attribute/measurements
increase	gorgonian condition	% live tissue, absence of disease and predators (<i>Cyphoma gibbosum</i>)
increase	gorgonian diversity	
increase	other invertebrates	<i>Diadema antillarum</i> , conch, lobsters, crabs, anemones
increase	sponge abundance	autotrophic sponges
decrease	sponge abundance	heterotrophic sponges
increase	sponge diversity	
increase	substrate condition	clean, no fuzzy algae (cyanobacteria), open space recruitment
increase	water clarity	
decrease	corallivores/bioeroders	bioeroders, <i>Coralliophila</i> (large size), clionids

2.4 Reference Condition for Biological Integrity

The panel agreed that all of the stations were impaired at some level. Many of the experts had been working in Puerto Rico for 30–40 years, while others had recently received their PhDs. The group had a rather lengthy discussion about the historical condition of Puerto Rico’s coral reefs in an attempt to answer: What did the reefs look like before humans came along to stress them?

The term “reference condition” is used by BCG to define biological condition in the absence of human disturbance (Stoddard et al. 2006). The concept of reference condition arose from the objective of the Clean Water Act Section 101: “to restore and maintain the chemical, physical and biological integrity of the nation’s waters”. Biological integrity is defined as “the community of organisms having a species composition, diversity and functional organization comparable to those of natural habitats within a region” (Karr 1991).

Unfortunately, human activities have significantly affected coral reefs. Puerto Rico’s coral reefs were severely degraded long before ecologists began to study them (Jackson 1997). According to Jackson et al. 2001, ecological extinction caused by overfishing preceded all other pervasive human disturbance to coastal ecosystems. Overfishing reduced species populations such as marine reptiles (green turtle, hawksbill turtle [1700s]), mammals (manatee and extinct Caribbean monk seal [1800s]), conch [1980s], fishes (Nassau and goliath groupers [1950s] and reef fishes [1970s]). By the time scientific studies began in the 1950s, herbivores and predators were reduced to very small fishes and sea urchins (Jackson 1997).

So, how can one estimate reference condition when no living human has ever seen a Puerto Rican reef in natural condition? One approach is the reference station approach, where scientists use reefs that are nearly unaffected by anthropogenic disturbance and the related stressors/exposures, or reefs whose present-day good condition is found in conjunction with the best available physical, chemical and biological habitat conditions, as surrogates for natural reefs. However, these

approaches may fail to correctly identify the baseline population sizes, instead representing a shifted baseline as reference condition (Pauly 1995).

Another approach is to apply ecological theory and empirical models to extrapolate reference condition. A relatively recent method is the use of historical ecology, where scientists piece together an understanding of what coral reef ecosystems used to be. The challenge is to use these approaches to adjust our expectations of what a healthy coral reef baseline looks like and use that as reference condition.

2.4.1 Experts' Examples of Reference Condition for Biological Integrity

Workshop panelists were urged to share examples of present or past reefs that they believed to exhibit full biological integrity. Dr. Szmant reported that she had participated in a discussion about shifting baselines with other experienced coral reef scientists. This led to a report (Sale and Szmant 2012) that summarized these scientists' reminiscences on historical reef condition over the last 40 years.

Dr. Szmant also showed photographs of recent changes from 2009 to 2012 in *Montastraea* populations on a coral reef in Curaçao (Watamula) that were caused by a 2010 bleaching event (Figure 2-2). An estimated 50% of previously large healthy coral colonies on Watamula showed partial or complete mortality in less than one year. She also showed photographs of extensive *Acropora cervicornis* beds on Smith Bank (on the south coast of Roatan) near the ones on Cordelia Banks that Dr. McField surveyed (see below). Dr. Szmant has slides of The Buoy and other reef areas in Puerto Rico and the Caribbean from the 1970s that she was willing to share.




Figure 2-2. Recent changes from 2009 to 2012 in *Montastraea* populations on a coral reef in Curaçao (Watamula) that were caused by a 2010 bleaching event (photos: Dr. Alina Szmant).

Dr. McField brought photos from Cordelia Banks on Roatan Island, Honduras (Figures 2-3, 2-4 and 2-5). Cordelia Banks is located near the airport and main port of Roatan, but strong onshore currents likely keep land-based sources of pollution away most of the time. Cordelia Banks is a good candidate for a reference site because it has:

- 52 acres of healthy reef with the highest live coral cover in the Caribbean (up to 73% measured in transects and averaging just over 50%). *Acropora cervicornis*, which is one of the most important reef species for structural reef growth and fish nursery habitat, dominates this reef. Unfortunately, this species and *Acropora palmata* have been reduced by about 98% over the last three decades throughout the Caribbean by disease and bleaching.
- The presence of two important species of sharks – the nurse shark (*Ginglymostoma cirratum*) and the Caribbean reef shark (*Carcharhinus perezii*).
- Spawning aggregation sites for groupers and snappers.



Figure 2-3. Two experts suggested Cordelia Banks near Roatan, Honduras, as an example of a reference site for excellent coral condition. This area contains 52 acres of threatened coral species, high fish abundance and other characteristics important in sustaining healthy reefs (photo: Dr. Melanie McField).



Figure 2-4. Monitoring of massive *Acropora cervicornis* banks at Cordelia Banks located off a major airport in Roatan, Honduras (photo: Dr. Melanie McField).



Figure 2-5. Panoramic view of *Acropora cervicornis* banks at Cordelia Banks, Roatan, Honduras (photo: Dr. Melanie McField).

Dr. McField also brought copies of the 2012 Report Card for the Mesoamerican Reef. Dr. McField's program, Healthy Reefs for Healthy People Initiative, is an international, multi-institutional effort that tracks the health of the Mesoamerican Reef (MAR), the human choices that shape it and the progress to ensure its long-term integrity. The Healthy Reefs for Healthy People Initiative seeks to address two overarching questions:

1. What is a healthy reef and how can we improve our tracking of reef health through a shared vision and common indicators or yardsticks?
2. How can we best convey consistent, scientific information to policymakers, decision-makers and the public, such that the connections between reef health and human health result in effective conservation action at an unprecedented scale?

The Healthy Reefs for Healthy People Initiative has developed a quantifiable, interpretive framework of measurable indicators and criteria to better understand and assess reef health in the MAR region. The Initiative produces report cards on the condition of the MAR resources, using a five-point grading system from very good to critical for key indicators: fish abundance, fleshy macroalgal index, *Diadema* abundance, herbivorous fish abundance, coral mortality, conch abundance, coral recruitment and coral cover. The report card also describes the main threats to the ecosystem and evaluates management actions. More information about the Initiative can be found at: <http://www.healthyreefs.org>

Dr. Weil stated that the average rainfall in Puerto Rico has been increasing steadily since 2000, coinciding with changes in land use. Consequently, rain events have a greater impact on the decline in coral reef condition. He has a draft report on this topic that he could send to other attendees. Dr. Weil also reports that in La Parguera, Puerto Rico, the average winter SST has remained elevated over the last decade. He suggests that water visibility is decreasing, and sedimentation is increasing. Yellow-band disease, which affects the *Montastraea annularis* species-complex, is now chronic all year, whereas it used to be seasonal and limited in distribution. Bleaching events in the area are followed by white plague infection, which leads to increased coral mortality with little or no recovery, and bare coral substrate is colonized by macroalgae.

Mr. Ruiz Torres passed around his recent book *Beneath the Waves*, published by the Sea Grant Program at the University of Puerto Rico (Figure 2-6). It contains nearly a hundred photos documenting the marine environment along the entire coast of Puerto Rico, including algae, fish, crustaceans, mollusks and corals. A description of the location, the depth and the characteristics of the organisms accompany each image. The text is written in English and Spanish.

Dr. Appeldoorn commented that good water flow (medium to high constant speed) is important for high quality reefs. He stated that fish trophic structure is impaired in Puerto Rico because of the low number of apex predators.



Figure 2-6. Cover for the book, *Beneath the Waves*, by Hector J. Ruiz Torres.

Dr. Vicente said that he has over 500 one-hour videos documenting reefs around Puerto Rico and USVI that he is willing to share with the group. He will send an index so experts can request the videos of interest.

2.4.2 Summary Discussion

The experts remarked that gorgonians were present in the videos of the best sites but not in high abundance. At fair condition sites, gorgonians were most abundant, but reduced abundance was seen again at poor coral reef condition (bell-shaped curve). The experts agreed that there is a need to understand the ecology of relationships between these assemblages to predict where we find certain species and abundance of corals and gorgonians, and why they are distributed that way. For example: Are corals replaced by gorgonians when corals die on reefs in lower or poor condition?

2.5 Attributes of a Very Good to Excellent Station

Based on the videos and photographs, the experts identified the attributes of a very good - to excellent station, which would be comparable to BCG Level 2: near natural (minimally disturbed). A summary of the attributes is shown in Table 2-5.

The attributes are reorganized in Table 2-6, into a format that can be more efficiently used during future workshops to facilitate establishing numeric criteria ranges.

2.5.1 Three-dimensional Topographic Complexity

The experts thought that very good - excellent stations would have high rugosity or three-dimensional topographic complexity, including substantial reef built above the bedrock. High topographic complexity is known to be an important attribute (Friedlander and Parrish 1998; Zawada 2011). Coral reefs with high topographic complexity have high species diversity (Talbot 1965; Risk 1972), primary productivity (Barnes 1988) and biomass density (Luckhurst and Luckhurst 1978; Carpenter et al. 1981). These reefs provide refuge from predation (Steele 1999; Idjadi and Edmunds 2006) and supplement larval settlement space (Idjadi and Edmunds 2006). Topographic complexity also provides hydrodynamic effects, determining water flow around, over and through the reef (Munk and Sargent 1954; Monismith 2007; Hearn 2008; Nunes and Pawlak 2008) and enhancing energy dissipation thereby, nutrient uptake and mass-transfer rates (Shashar et al. 1996; Hearn et al. 2001).

Table 2-5. Summary of descriptions of four condition categories (very good to poor) based on expert assessments of individual stations. The descriptions of good to poor condition are comparisons to a very good condition station based on panelists' identifications of aspects missing from expectations for very good stations.

Condition level	Attribute descriptions
Very Good Excellent (approximate BCG Level 1–2)	Physical structure: High rugosity or 3D structure; substantial reef built above bedrock; many irregular surfaces provide habitat for fish; very clear water; no sediment, flocs or films
	Corals: High species diversity including rare; large old colonies (<i>Montastraea</i>) with high tissue coverage; balanced population structure (old and middle-sized colonies, recruits); <i>Acropora</i> thickets present
	Gorgonians: Gorgonians present but subdominant to corals
	Sponges: Large autotrophic and highly sensitive sponges abundant
	Fish: Populations have balanced species abundances, sizes and trophic interactions
	Large vertebrates: Large, long-lived species present and diverse (turtles, eels, sharks)
	Other invertebrates: <i>Diadema</i> , lobster, small crustaceans and polychaetes abundant; some large sensitive anemone species present
	Algae: Crustose coralline algae abundant; turf algae present but cropped and grazed by <i>Diadema</i> and herbivorous fish; low abundance of fleshy algae
Good (approximate BCG Level 3)	Physical structure: Moderate to high rugosity; moderate reef built above bedrock; some irregular cover for fish habitat; water slightly turbid; low sediment, flocs or films on substrate
	Corals: Moderate coral diversity; large old colonies (<i>Montastraea</i>) with some tissue loss; varied population structure (usually old colonies, few middle aged and some recruits); <i>Acropora</i> thickets may be present; rare species absent
	Gorgonians: Gorgonians more abundant than Levels 1–2
	Sponges: Autotrophic species present but highly sensitive species missing
	Fish: Decline of large apex predators (e.g., groupers, snappers) noticeable; small reef fishes more abundant
	Large vertebrates: Large, long-lived species locally extirpated (turtles, eels)
	Other invertebrates: <i>Diadema</i> , lobster, small crustaceans and polychaetes less abundant than Levels 1–2; large sensitive anemone species absent
	Algae: Crustose coralline algae present but fewer than Levels 1–2; turf algae present and longer, more fleshy algae present than Levels 1–2
	Condition: Disease and tumor presence slightly above background level; more colonies have irregular tissue loss

Table 2-5 (continued)

Condition level	Attribute descriptions
Fair (approximate BCG Level 4)	Physical structure: Low rugosity; limited reef built above bedrock; erosion of reef structure obvious; water turbid; more sediment accumulation, flocs and films; <i>Acropora</i> usually gone or present as rubble for recruitment substrate
	Corals: Reduced coral diversity; emergence of tolerant species, few or no living large old colonies (<i>Montastraea</i>); <i>Acropora</i> thickets gone, large remnants mostly dead with long uncropped turf algae
	Gorgonians: Gorgonians more abundant than Levels 1–3, replacing sensitive coral and sponge species
	Sponges: Mostly heterotrophic tolerant species and clionids
	Fish: Absence of small reef fishes (mostly Damselfish remain)
	Large vertebrates: Large, long-lived species locally extirpated (turtles, eels)
	Other invertebrates: <i>Diadema</i> absent; <i>Palythoa</i> overgrowing corals; crustaceans, polychaetes and sensitive anemones conspicuously absent
	Algae: Some coralline algae present but no crustose algae; turf is uncropped, covered in sediment; abundant fleshy algae (e.g., <i>Dictyota</i>) with high diversity
Poor (approximate BCG Level 6)	Physical structure: Very low rugosity; no or little reef built above bedrock; no or low relief for fish habitat; very turbid water; thick sediment film and thick floc covering bottom; no substrate for recruits
	Corals: Absence of colonies, those present are small; only highly tolerant species with little or no live tissue
	Gorgonians: Small and sparse colonies; mostly small sea fans; often diseased
	Sponges: Heterotrophic sponges buried deep in sediment; highly tolerant species
	Fish: No large fishes; only a few tolerant species remain; lack of multiple trophic levels
	Large vertebrates: Usually devoid of vertebrates other than fishes
	Other invertebrates: Few or no reef invertebrates; high abundance of sediment dwelling organisms such as polychaetes and holothurians
	Algae: High cover of fleshy algae (<i>Dictyota</i>); complete absence of crustose coralline algae
	Condition: High incidence of disease and low or no tissue coverage on small colonies of corals, sponges and gorgonians, if present

Table 2-6. Condition levels and associated attributes

Condition level	BCG level	Physical structure	Corals	Gorgonians	Sponges	Fish	Vertebrates	Other invertebrates	Algae/plants	Condition
Very Good - Excellent	1–2	High rugosity or 3D structure; substantial reef built above bedrock; many irregular surfaces provide habitat for fish; very clear water; no sediment, flocs or films	High species diversity including rare; large old colonies (<i>Montastraea</i>) with high tissue coverage; balanced population structure (old and middle-aged colonies, recruits); <i>Acropora</i> thickets present	Gorgonians present but sub-dominant to corals	Large autotrophic and highly sensitive sponge species abundant	Populations have balanced species abundance, sizes and trophic interactions	Large, long-lived species present and diverse (turtles, eels, sharks)	<i>Diadema</i> , lobster, small crustaceans and polychaetes abundant; some large sensitive anemone species	Crustose coralline algae abundant; turf algae present but cropped and grazed by <i>Diadema</i> and herbivorous fish; low abundance fleshy algae	Low prevalence of disease or tumors; mostly live tissue on colonies
Good	3	Moderate to high rugosity; moderate reef built above bedrock; some irregular cover for fish habitat; water slightly turbid; low sediment, flocs or film on substrate	Moderate coral diversity; large old colonies (<i>Montastraea</i>) with some tissue loss; varied population structure (usually old colonies, few middle-aged and some recruitment); <i>Acropora</i> thickets may be present; rare species absent	Gorgonians more abundant than in Levels 1–2	Autotrophic species present but highly sensitive species missing	Decline of large apex predators (e.g., groupers, snappers, etc.) noticeable; small reef fish more abundant than Levels 1–2	Large, long-lived species locally extirpated (e.g., turtles, eels)	<i>Diadema</i> , lobster, small crustaceans and polychaetes less abundant than Levels 1–2; large sensitive anemone species missing	Crustose coralline algae present but less than Levels 1–2; turf algae present and longer; more fleshy algae present	Disease and tumor prevalence slightly above background level; more colonies have irregular tissue loss

Table 2.6 (continued)

Condition level	BCG level	Physical structure	Corals	Gorgonians	Sponges	Fish	Vertebrates	Other invertebrates	Algae/plants	Condition
Fair	4	Low rugosity, limited reef built above bedrock; erosion of reef structure obvious; water turbid; more sediment accumulation, flocs and films; <i>Acropora</i> usually gone or present as rubble for recruitment substrate	Reduced coral diversity; emergence of tolerant species, few or no large old colonies (<i>Montastraea</i>) mostly dead; <i>Acropora</i> thickets gone; large remnants mostly dead with long uncropped turf algae	Gorgonians more abundant than in Levels 1 - 3; replace sensitive coral and sponge species	Mostly heterotrophic sponges with tolerant species and clionids	Absence of small reef fish (mostly Damsel fish)	Large, long-lived species locally extirpated (e.g., turtles, eels)	<i>Diadema</i> absent; <i>Palythoa</i> overgrowing corals, crustaceans, polychaetes and sensitive anemones conspicuously absent	Some coralline algae; turf is uncropped covered in sediment; lots of fleshy algae with high diversity (e.g., <i>Dictyota</i>); possibly covering sessile invertebrates; no turf or coralline algae; complete absence of crustose coralline algae	High incidence of diseased coral, sponges, gorgonians; evidence of high mortality; usually less tissue than dead portions on colonies
Poor	6	Very low rugosity, no or low reef built above bedrock or poor for fish habitat; very turbid water; thick sediment film and high flocs covering bottom; no substrate for recruits	Absence of colonies, those present are small, only highly tolerant species, little or no tissue	Small and sparse colonies, mostly small sea fans, often diseased	Heterotrophic sponges buried deep in sediment; highly tolerant sponge species	No large fish, few tolerant species, lack of multiple trophic levels	Usually devoid of other vertebrates	Low or no reef invertebrates; high abundance of sediment dwelling organisms (e.g., polychaetes, holothurians)	High cover of fleshy algae (<i>Dictyota</i>); possibly covering sessile invertebrates; no turf or coralline algae; complete absence of crustose coralline algae	High incidence disease on small colonies of corals, sponges and gorgonians; if present, low or no tissue coverage

2.5.2 Stony Coral Attributes

For stony corals, attendees decided that very good - excellent stations would have high species diversity that included large colonies of reef-building corals (i.e., *Montastraea*) and those colonies would have high tissue coverage. The *Montastraea annularis* species-complex (*Montastraea annularis*, *Montastraea faveolata* and *Montastraea franksi*) was historically one of the primary reef framework builders of the Caribbean coral reefs, characterizing the “buttress zone” or “annularis zone” in the classical descriptions of Caribbean reefs (Goreau 1959). These corals have declined dramatically throughout their range. The *Montastraea annularis* species-complex is susceptible to bleaching (Oxenford et al. 2008; Brandt 2009; Bruckner and Hill 2009; Wagner et al. 2010), disease (Bruckner and Hill 2009; Miller et al. 2009), sediment (Eakin et al. 1994; Carricart-Ganivet and Merino 2001; Torres and Morelock 2002) and nutrients (Marubini and Davies 1996).

The experts believed that the coral reef population should have a balanced size-class structure, including large and middle-sized colonies as well as new recruits. Ecologists consider population demographics to be vital statistics, particularly those statistics that can impact on present and future population size (Hughes 1996; Edmunds 2013; Edmunds and Elahi 2007). Typically, expanding populations have a large percentage of young individuals, while declining populations have a large percentage of older individuals and stable populations have a relatively even size distribution among age groups.

The experts also concluded that *Acropora palmata* thickets should be present. *A. palmata* was formerly the dominant species in shallow water (3–16 ft. deep) throughout the Caribbean and on the Florida Reef Tract, forming extensive, densely aggregated thickets in areas of heavy surf. These coral colonies prefer the exposed reef crest and fore-reef environments in depths of < 20 ft., although isolated corals may occur to depths of 65 ft. Since 1980, populations have collapsed throughout their range from disease outbreaks, with losses compounded locally by hurricanes, increased predation, bleaching, elevated temperatures and other factors (Ruzicka et al. 2013). This species is also particularly susceptible to damage from sedimentation (NOAA 2013b).

2.5.3 Gorgonian Attributes

There was considerable discussion about the relative distribution of gorgonians and stony corals. The experts decided that very good - excellent stations would have gorgonians present, but the station should be dominated by stony corals. Gorgonians form a major benthic component of Caribbean reefs (Bayer 1973; Brazaeu and Lasker 1989) and can be very abundant in some sites where stony corals apparently are unable to proliferate. Factors controlling the distribution of shallow-water gorgonians include water motion and substrate relief (Kinzie 1973; Yoshioka and Yoshioka 1989a, b) and sediment transport (Yoshioka and Yoshioka 1989b, 2009).

2.5.4 Sponge Attributes

The experts agreed that very good - excellent stations would have high abundances of large autotrophic and highly sensitive sponge species. Most sponges are heterotrophic organisms, obtaining their food from the open water column. However, 35 species of common Caribbean sponges possess photosynthetic endosymbionts (Vicente 1990) that supply food to their hosts (Wilkinson 1983; Thacker and Freeman 2012). This is similar to the relationship between zooxanthellae and their coral hosts. Roberts et al. (2006) found that exposure to shade and siltation significantly reduced the growth and reproductive status of the temperate photosynthetic reef sponge *Cymbastela concentrica*.

2.5.5 Fish Attributes

Workshop participants decided that populations should have balanced distributions of species abundances, sizes and trophic interactions. Caribbean coral reefs can contain as many as 500–700 species of fishes (Lieske and Collins 2001). The mechanisms that lead to these concentrations of fish species on coral reefs have been widely debated over the last 50 years. While many reasons have been proposed there is no scientific consensus on a primary mechanism and it seems likely that a number of factors contribute. These include the rich habitat complexity and diversity inherent in coral reef ecosystems (Luckhurst and Luckhurst 1978; Gladfelter et al. 1980) and the variety and temporal availability of food resources available to coral reef fishes (Randall 1967).

Puerto Rico reef fisheries have shown significant decline since the 1970s, and large reef fishes have virtually disappeared from shallow reefs around Puerto Rico (Garcia-Sais et al. 2008). Fishing may have direct and indirect effects on reef fish trophic structure. Removals of apex predators from the reef complex may result in shifts of species composition (e.g., through trophic and ecological cascades) and for some taxa, increased variability in population dynamics or potential effects on species evolution.

2.5.6 Large Vertebrate Attributes

Several groups of large, long-lived vertebrate species (e.g., sea turtles and manatees) are considered important contributors to Puerto Rican coral reef communities. Other groups (e.g., dolphins, whales, seabirds) spend most of their life cycle in other habitat types but are occasionally seen hunting or feeding in waters around coral reefs. Puerto Rico supports five species of marine turtles, two of which (Hawksbill and Leatherback) are critically endangered. Four sharks (Blacktip Shark, Reef Shark, Tiger Shark and Nurse Shark), eight eels (Brown Garden Eel, Sharptail Eel, Goldspotted Eel, Spotted Snake Eel, Green Moray, Golden Moray, Spotted Moray, Purplemouth Moray) and two rays (Spotted Eagle Ray and Southern Stingray) can also be found on Puerto Rico coral reefs. A recent study (Jackson et al. 2012) found that the biomass of apex predators (sharks, large snappers and groupers) was close to zero in Puerto Rico. The experts decided that large long-lived species should be present and diverse at very good - excellent stations.

2.5.7 Other Invertebrate Attributes

Queen conch, spiny lobsters and some crabs are harvested for food and have been declining throughout the Caribbean for decades (Santavy et al. 2012). Conchs are generally acknowledged to be over-exploited (Appeldoorn and Meyers 1993), and Puerto Rico has established catch limits for these mollusks. Sea urchins (especially *Diadema antillarum*) are important herbivores that were decimated by an epizootic throughout the western Atlantic in the 1980s (Lessios et al. 1984; Lessios 1988, 2005). The population status for the Caribbean spiny lobster stock is unknown (NOAA 2013a).

The experts decided that very good - excellent stations would have abundant *Diadema*, lobster, small crustaceans and polychaetes. They also felt that some large sensitive anemone species should be present.

2.5.8 Algae Attributes

Macroalgae and turf algae compete for space with coral, sponge and other sessile species. Excess nutrients may alter competitive interactions and favor algae over coral. Many fishes and invertebrates are key grazers, helping to maintain algal biomass and prevent algae from overgrowing coral. A number of algal species (e.g., calcareous macroalgae and crustose coralline algae) deposit calcium carbonate during growth and may contribute to reef structural strength. Crustose coralline algae may also facilitate recruitment of stony coral. Algae are primary producers and provide habitat and resources for marine fish and invertebrates but often not to the same degree as coral reef habitat (Santavy et al. 2012). The experts decided that very good - excellent stations would have abundant crustose coralline algae, turf algae would be present but cropped and grazed and fleshy algae would occur in low abundance.

2.5.9 Condition

Bleaching, disease or predation can affect health and condition of stony corals, gorgonians and sponges. An indicator of stony coral/gorgonian health is the amount of live tissue on the organism or colony. However, coral reef fish rarely appear to suffer from tumors or lesions (Panek 2005). The experts decided that there should be a very low prevalence of disease on very good - excellent stations, with mostly live tissue on coral colonies and gorgonians, and low prevalence of tumors on coral reef fish.

Chapter 3. Discussion and Next Steps

3.1 Discussion

EPA convened a group of experts to attempt, for the first time, to develop a coral reef BCG. There was a consensus among the experts that this was an important contribution, because “We have been documenting the demise of coral reefs, instead of taking action to change the direction of their existence”.

A preliminary BCG based on stony corals, fishes, gorgonians, sponges, vertebrates and other invertebrates has been assembled for shallow-water linear reefs of southwestern Puerto Rico. The experts were able to identify four distinct levels of condition: very good – excellent, good, fair and poor. Additional discussion is needed to develop reference condition for biological integrity (e.g., natural level).

Attribute development during the first workshop relied primarily on viewing videos and photos from individual coral reef monitoring sites. This approach resulted in attributes that were largely species-based, with a single notable addition (e.g., organism condition).

EPA anticipates that ecosystem connectivity is also an appropriate attribute to include in a coral reef BCG, since connectivity among coral reefs, mangroves, sea grass beds and lagoons provides a complex and dynamic mosaic that is well documented as a critical ecosystem attribute (Christensen et al. 2003; Cervený 2006; Mumby et al. 2004, 2008; Aguilar-Perera and Appeldoorn 2007; McField and Kramer 2007; Meynecke et al. 2008; Sale et al. 2008; Pittman et al. 2011).

EPA would like to suggest that considering the attributes at multiple scales, similar to the approach being developed for estuarine ecosystems, may also be informative for coral reef ecosystems. The estuarine BCG framework considers structure, function, condition, connectivity and non-native species in waterbodies at multiple scales, using measures such as seagrass health, benthic faunal indices and habitat mosaics (Cicchetti and Pryor 2010; EPA in review). This holistic and integrated approach is intended to improve the understanding and management of the cumulative impacts of multiple stressors in complex waterbodies and should work well for coral reefs.

3.2 Next Steps

3.2.1 Second Workshop

EPA is planning to hold a second workshop in early 2014. At the second workshop EPA hopes to focus more on: 1) different scales and attributes associated with the entire reef ecosystem, 2) tolerance of coral reef species to various anthropogenic stressors and 3) the process for moving towards a quantitative BCG, including development of a data portal to organize and share all of the available data from Puerto Rican coral reef ecosystems. EPA would also like to continue the discussion of reference condition and try to reach consensus on attributes for the reference condition level.

3.2.2 Species Tolerance Database

In preparation for the first workshop, Ms. Bradley and Dr. Santavy began developing spreadsheets of coral reef taxa for each assemblage (e.g., stony corals, octocorals, sponges, other invertebrates, fishes, reptiles, mammals, mangroves, seagrasses and algae) and their characteristics as related to the ten attributes of the BCG, including tolerance levels to various stressors, vulnerabilities, habitat, etc. Thresholds, when known, were documented.

Species lists for the database rows were derived from Miller and Lugo (2009). Some columns are consistent across assemblages (e.g., scientific name, common name, common/rare, tolerance to pollution, tolerance to temperature change, tolerance to wave energy and susceptibility to disease). Other columns are unique to specific assemblages (e.g., for stony corals: maximum colony size, tolerance to acidity, collection or trade; while for fish: juvenile habitat, adult habitat, food preference, solitary/aggregating). Ms. Bradley and Dr. Santavy began to populate the spreadsheets with data, beginning with information from the Humann and DeLoach field guides (Humann and DeLoach 2002a, b; 2003) and Sefton and Webster (1986).

During the first workshop, Ms. Bradley gave a short presentation to introduce the spreadsheets. The group then moved into a brief facilitated discussion. Workshop participants seemed to respond very positively to the concept and were interested in collaboratively working to complete the spreadsheets. The group will discuss how to go about completing the spreadsheets during the second workshop.

3.2.3 Assembling the Monitoring Data

To complete the BCG, the group will utilize pre-existing data collected by others in laboratory and field studies. In Phase 1, EPA is working with EPA and NOAA data for Puerto Rico and USVI (stony corals, fish and benthic invertebrates). The initial biological data set will include fish measurements from several studies conducted by NOAA and EPA. Both groups used the same survey methods for fish, so standardization will easily occur as existing datasets are compiled. The second data set will include stony coral measurements from the same studies by NOAA and EPA. In this case, the methods are very different and will require discussion with the EPA coral reefs BCG team to extract the most meaningful and comparable data for standardized reporting. The third data set will include benthic invertebrates from the same studies by NOAA and EPA. The two survey methods are quite similar, although NOAA counts lobsters, conch and *Diadema*, while EPA also counts crabs and additional urchin species.

For all datasets, EPA will normalize taxonomic naming protocols to the Integrated Taxonomic Information System (ITIS). The standardized data set will contain data in the original format, a crosswalk with translations for converting data and the final standardized format. The data set will also include a field for the organization that generated the data (data owner).

EPA is planning on completing Phase 1 (as described above) prior to the second workshop and plans to use these data during the workshop.

Phase 2 activities will include direct submission into the STORET Data Warehouse (short for STORage and RETrieval). STORET is a repository for water quality, biological and physical data and is used by state and territorial environmental agencies for their water quality data under the CWA. Phase 2 will also include incorporation of additional data and fields, when additional datasets from the coral reef BCG partners are provided (Puerto Rico Department of Natural and Environmental Resources, Puerto Rico Environmental Quality Board, USVI Department of Planning and Environmental Resources, University of Puerto Rico, University of the Virgin Islands, US Geological Survey and National Park Service). The EPA Office of Water has agreed to make the necessary modifications to STORET to include these data. Additionally, metadata for the database in STORET (with URL) will be developed for inclusion in NOAA's Coral Reef Information System (CORIS).

3.2.4 Calibrating the BCG

The group will begin calibrating the BCG by using found (or existing) data to confirm the ecological attributes developed during this first workshop. The experts have determined narrative decision rules for assigning stations to a BCG level on the basis of the photographs and videos collected at multiple stations. Documentation of expert opinion in assigning stations to BCG levels is critical to the process. Next, a decision model will be developed that incorporates those rules and will be tested with independent data sets. The decision model using the tested decision rules will provide a transparent, formal and verifiable method for documenting and validating expert knowledge. A quantitative data analysis program can be developed using those rules.

BCG level descriptions in the conceptual model are qualitative (e.g., high diversity, reduced diversity; Table 2-5) to allow for consistent assignments of stations to levels. It is necessary to formalize and quantify the expert knowledge by codifying level descriptions into a set of quantitative rules (e.g., Droesen 1996). If formalized and quantified, any person (with data) can follow the rules to obtain the same level assignments as the group of experts. This makes the actual decision criteria transparent to stakeholders and potentially applicable to similar types of coral reefs in other areas.

Rules are logic statements that experts use to make their decisions, for example: "If taxon richness is high, then biological condition is high." Rules on attributes can be combined, for example: "If the number of highly sensitive coral taxa (Attribute II) is high, and the number of tolerant colonies (Attribute V) is low, then the assignment is to Level 2." The categories high, moderate, low, etc., are ordinal categories: we know that moderate is greater than low; but the boundaries of the categories are fuzzy and somewhat subjective. In iterations of the process, the expert panel is asked to put quantitative boundaries on the categories they have defined. The objective is to derive combined rules, for example, "If there are more than 10 highly sensitive coral taxa, and the percentage of colonies of tolerant taxa is less than 15%, then assignment is Level 2." The quantitative rules preserve the collective professional judgment of the expert group and set the stage for the development of quantitative models that reliably assign stations to levels without having to reconvene the same expert group. In essence, the rules and the models capture the panel's collective decision criteria.

The decision rule for a single level of the BCG does not usually rest on a single attribute (e.g., highly sensitive taxa) and generally includes other attributes (intermediate sensitive taxa, tolerant taxa, indicator species); such rules are termed Multiple Attribute Decision Rules. After verification with independent data, the quantified rules allow users to consistently assess stations according to the same rules used by the expert panel and allow a computer algorithm, or other persons, to obtain the same level assignments as the original panel. Documentation of the rules and algorithm allow new panels to review and modify the decision rules as necessary.

3.2.5 Economic Valuation of Coastal Ecosystem Services

Despite their open-access nature and many contributions to the public good, coral reefs have often been undervalued in decision-making (Brander et al. 2009). The natural features of a coral reef (including physical structure, water quality, biological organisms and ecological functions) provide many natural benefits to human societies, collectively known as ecosystem goods and services. The economic values of some services (e.g., commercial fishing) are established in markets, while other services provide nonmarket value for local, state/regional and national/international segments of the population (Principe et al. 2012). Most ecosystem service studies have focused on market benefits, which are relatively easy to incorporate in trade-off analyses, but coral reefs also provide numerous nonmarket ecosystem services (e.g., existence value and cultural value) that can be estimated using a variety of methods.

Estimates of the global value of coral reefs range from US \$30 billion per year (Cesar et al. 2003) to US \$377 billion per year (Costanza et al. 1997). However, global estimates are coarse and rarely relevant to local management decisions. Decision contexts differ with reef type and habitat, political climate, stakeholder interests, decision authorities and responsibilities, knowledge, management capacity and expertise. Every decision contains an element of valuation, but values are not always at the forefront of finding optimal decisions (Keeney 1996). Consequences resulting from a decision are often described in terms of value (Hastie 2001). Yet, the values of stakeholders often go ignored before management strategies are implemented. Public and stakeholder values, cares, and priorities should be considered throughout in the focus and design of assessments and management planning and should not be an afterthought in the process.

The BCG effort focuses on how attributes of the coral reef ecosystem change in response to increasing anthropogenic stress. The attributes of the coral reef ecosystem represent the “glue” (Pearce and Moran 1994; Turner et al. 2000) of the properly functioning ecosystem, supporting the growth of reef-building corals for ecosystem services (e.g., shoreline protection, the presence of unique and diverse species to attract tourists, the creation of potentially useful natural products and the maintenance of habitat and nurseries for harvestable fish stocks). The development of concise, rigorous definitions and levels of condition along the human disturbance gradient will provide the fundamental understanding of the factors that affect delivery of ecosystem services.

EPA and NOAA, in partnership with the University of Puerto Rico and the Puerto Rico Sea Grant, are conducting a study to provide the economic valuation of tourism and recreation associated with Puerto Rico's coral reefs to help improve our understanding of the real costs of decisions and management options. Reef-related tourism activities include snorkeling, diving, fishing, viewing wildlife, boating, beach use and surfing. The project will consist of a modified form of the method NOAA used in the Florida Keys National Marine Sanctuary (Leeworthy and Wiley 1996, 1997, 2003; Leeworthy and Bowker 1997; Leeworthy and Vanasse 1999; Park et al. 2002; Bhat 2003; Shirlany et al. 2008), in Southeast Florida (Johns et al. 2001) and in Hawaii (Bishop et al. 2011).

The study is estimating the use and associated market (spending and associated impacts on total output/sales, income and employment) and non-market economic value (consumer's surplus or the net value received by those doing recreation activities on the reef over and above what they pay to undertake the activities) and how those values change with changes in reef attributes. Linking the relationships of how reef attribute values change with changes in the physical/natural levels of those attributes can be used to measure the economic benefits of the changes and thus provide additional performance measures of management actions to protect and restore coral reef ecosystems. Table 3-1 shows coral reef ecosystem services and examples of coral reef attributes that are associated with them.

Table 3-1. Coral reef ecosystem services and reef attributes (adapted from Principe et al. 2012).

Ecosystem service(s)		Natural features (reef attributes)
Final	Intermediate	
Recreational fishing opportunity	Production of benthic and aquatic prey for consumption by recreational fish	Fish diversity and abundance
Recreational diving/snorkeling opportunity	Coral reef formation and maintenance; maintenance of water clarity; production of benthic and aquatic prey for consumption by recreational fish	Coral diversity, abundance and health; fish diversity and abundance; water clarity
Recreational underwater photography opportunity	Coral reef formation and maintenance; maintenance of water clarity; production of benthic and aquatic prey for consumption by recreational fish	Coral diversity, abundance and health; fish diversity and abundance; water clarity
Recreational surfing opportunity	Reef breaks	3-D reef structure
Opportunity to view nature and wildlife	Biological integrity	Biodiversity (birds, marine mammals, turtles)
Opportunity to sunbathe and swim at the beach	Water quality, shoreline protection, sand production	White coralline sands; calm waters
Opportunity to collect objects (beachcombing)	Water quality	Wide sandy beaches, biodiversity, occasional storms

Definitions

- Final ecosystem service – Output of ecological functions or processes that directly contributes to social welfare or has the potential to do so in the future (*sensu* Boyd and Banzhaff 2007).
- Intermediate ecosystem service – Output of ecological functions or processes that indirectly contributes to social welfare or has the potential to do so in the future.
- Natural features – The biological, chemical and physical attributes of an ecosystem or environment.

Estimates of use and value will be made for five regions in Puerto Rico to provide information on the economic value of reefs in various levels of condition, including present condition. Use and economic information can be used in evaluating the economic benefits of investments in protection and restoration of the coral reef ecosystems. Results can be used by both private businesses and government agencies responsible for managing coral reefs in marketing, education and outreach efforts, including Puerto Rico's coral reef management activities in the four coral reef priority areas (Culebra, the Northeast Reserve, Cabo Rojo and Guánica).

3.3 Final Thoughts

The first workshop was a new experience for all involved. While EPA has worked with states and territories to develop BCGs for streams and estuaries, no one has ever attempted to develop a BCG for coral reefs. The experts met the challenge head-on and great progress was made. EPA anticipates that this is just the first of several expert workshops. EPA will host conference calls and webinars as appropriate. EPA has asked that the experts from the first workshop commit to working with us throughout the process. Experts who were not able to attend the first workshop will also be invited to participate in future workshops and webinars.

Appendix A. References

- Adams AJ, Dahlgren CP, Kellison GT, Kendall MS, Layman CA, Ley JA, Nagelkerken I and Serafy JE. 2006. Nursery function of tropical back-reef systems. *Marine Ecology Progress Series* **318**: 287–301.
- Aguilar-Perera A and Appeldoorn RS. 2007. Variation in juvenile fish density along the mangrove-seagrass-coral reef continuum in SW Puerto Rico. *Marine Ecology Progress Series* **348**:139–148.
- Almy CC and Carrion-Torres CC. 1963. Shallow-water stony corals of Puerto Rico. *Caribbean Journal of Science* **2/3**:133–162.
- Alvarez-Filip L, Kulvy N, Gill JA, Isabelle M and Watkinson AR. 2009. Flattening of Caribbean coral reefs: region wide declines in architectural complexity. *Proceedings of the Royal Society B*, **276**:3019–3025.
- Andrews JC and Pickard GL. 1990. The physical oceanography of coral-reef systems. In: *Ecosystems of the World, Volume 25, Coral Reefs*. Z. Dubinsky. (Ed.). New York. pp. 11–48.
- Appeldoorn RS and Meyers S. 1993. Part Z. Puerto Rico and Hispaniola. In: *Marine fishery resources of the Antilles*. FAO Fisheries Technical Paper No. 326 Rome, FAO. pp. 99–158.
- Ayala CJ. 1999. *American sugar kingdom: the plantation economy of the Spanish Caribbean*. UNC Press. ISBN 978-0-8078-4788-6.
- Bak RPM, Carpay MJE and de Ruyter van Steveninck. (Eds). 1983. Density of the sea urchin *Diadema antillarum* before and after mass mortalities on the coral reefs of Curaçao. *Marine Ecology Progress Series* **17**:105–108.
- Ballantine DL and Aponte NE. 1997a. A revised checklist of the benthic marine algae known to Puerto Rico. *Caribbean Journal of Science* **33**:150–179.
- Ballantine DL and Aponte NE. 1997b. Notes on the benthic marine algae of Puerto Rico VI. Additions to the flora. *Botanica Marina* **40**:39–44.
- Ballantine DL and Aponte NE. 2002. *A Checklist of the Benthic Marine Algae Known to Puerto Rico, Second Revision*. Revised 2002, URL: <http://128.32.109.44/fest/PRCL/checklist.html>.
- Ballantine D, Appeldoorn R, Yoshioka P, Weil E, Armstrong R, Garcia J, Otero E, Pagan F, Sherman C, Hernandez-Delgado E, Bruckner A and Lilyestrom C. 2008. Biology and Ecology of Puerto Rican Coral Reefs. In: Riegl BM, Dodge RE. (Eds). *Coral Reefs of the USA*. Springer, pp. 375–406.
- Barber RT, Hilting AK and Hayes ML. 2001. The changing health of coral reefs. *Human and Ecological Risk Assessment* **7**:1255–1270.
- Barbour MT, Gerritsen J, Snyder BD and Stribling JB. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish, Second Edition. US Environmental Protection Agency, Office of Water, Washington, DC. EPA 841-B-99-002.

- Barnes DJ. 1988. Seasonality in community productivity and calcification at Davies Reef, central Great Barrier Reef. *Proceedings of the Sixth International Coral Reef Symposium*. **6**: 521-527.
- Bayer FM. 1973. Colonial organization in octocorals. In: *Animal Colonies: Development and Function Through Time*. Boardman RS, Cheetham AH and Oliver Jr. (Eds.). WA. Wiley and Sons, Inc. pp. 69–93.
- Beets J and Friedland A. 1998. Evaluation of a conservation strategy: a spawning aggregation closure for red hind, *Epinephelus guttatus*, in the U.S. Virgin Islands. *Environmental Biology of Fishes* **55**:91–98.
- Beller W, Casellas MA, Cerame-Vivas MJ, Duffy L, El Koury J, Gelabert PA, Gonzales Liboy JA, Hernandez-Avila M, Maldonado N, Matos CA, Mignucci-Giannonni A, Pantoja-Garcia E, Rigau J, Shelley D, Tacher-Roffe M and N. Zerbi N. 1999. *Puerto Rico and the Sea: An action program for marine affairs*. Report to the governor, PR. Department of Natural Resources, San Juan, PR.
- Benedict JE. 1902. Anomuran collection of Porto Rico. *Bulletin of the United States Fish Commission* **20**:129–148.
- Bhat MG. 2003. Application of non-market valuation to the Florida Keys marine reserve management. *Journal of Environmental Management* **67**:315–325.
- Bigelow RP. 1902. The Stomatopoda of Porto Rico. *Bulletin of the United States Fish Commission* **20**:149–160.
- Bishop RS, Chapman DJ, Kanninen BJ, Krosnick JA, Leeworthy B and Neade NF. 2011. *Total Economic Value for Protecting and Restoring Hawaiian Coral Reef Ecosystems Final Report*. Silver Spring, MD: NOAA Office of National Marine Sanctuaries, Office of Response and Restoration, and Coral Reef Conservation Program. NOAA Technical Memorandum CRCP 16, 405 pp.
- Bradley P, Fisher WS, Bell H, Davis WS, Chan V, LoBue C and Wiltse W. 2008. Development and implementation of coral reef biocriteria in US jurisdictions. *Environmental Monitoring and Assessment* **150**:43–51.
- Bradley P, Fore L, Fisher W and Davis W. 2010. *Coral Reef Biological Criteria: Using the Clean Water Act to Protect a National Treasure*. Narragansett (RI): U.S. Environmental Protection Agency, Office of Research and Development, EPA/600/R-10/054, July 2010.
- Brainard RE, Birkeland C, Eakin CM, McElhany P, Miller MW, Patterson M and Piniak GA. 2011. *Status review report of 82 candidate coral species petitioned under the U.S. Endangered Species Act*. US Department of Commerce. NOAA Technical Memorandum. NOAA-TM-NMFS-PIFSC-27, 530 p. + 1 Appendix.
- Brander LM, Rehdanz K, Tol RSJ and van Beukering PJH. 2009. *The economic impact of ocean acidification on coral reefs*. ESRI Working Paper No. 282.
- Brandt ME. 2009. The effect of species and colony size on the bleaching response of reef-building corals in the Florida Keys during the 2005 mass bleaching event. *Coral Reefs* **28**: 911–924.
- Brazeau DA and Lasker HR. 1989. The reproductive cycle and spawning in a Caribbean gorgonian. *Biological Bulletin* **176**:1–7.

- Brown BE and Howard LS. 1985. Assessing the effects of “stress” on reef coral. *Advances in Marine Biology* **22**:1–63.
- Bruckner AW and Bruckner RJ. 1997. Outbreak of coral disease in Puerto Rico. *Coral Reefs* **16**:260.
- Bruckner AW and Bruckner RJ. 2006. Consequences of yellow band disease (YBD) on *Montastraea annularis* (species complex) populations on remote reefs off Mona Island, Puerto Rico. *Diseases of Aquatic Organisms* **69**:67–73.
- Bruckner AW and Hill RL. 2009. Ten years of change to coral communities off Mona and Desecheo Islands, Puerto Rico, from disease and bleaching. *Diseases of Aquatic Organisms* **87**:19–31.
- Budd AF, Fukami H, Smith ND and Knowlton N. 2012. Taxonomic classification of the reef coral family Mussidae (Cnidaria: Anthozoa: Scleractinia). *Zoological Journal of the Linnean Society* **166**:465–529.
- Burke L and Maidens J. 2004. *Reefs at risk in the Caribbean*. World Resources Institute, Washington, DC. 81 pp.
- Carpenter KE, Miclat RI, Albaladejo VD and Corpuz VT. 1981. The influence of substrate structure on the abundance and diversity of Philippine reef fishes. *Proceedings of the Fourth International Coral Reef Symposium* **2**:497–502.
- Carricart-Ganivet JP and Merino M. 2001. Growth responses of the reef-building coral *Montastraea annularis* along a gradient of continental influence in the southern Gulf of Mexico. *Bulletin of Marine Science* **68**:133–146.
- Cerame-Vivas MJ. (Ed.). 2001. *Ecologia de Puerto Rico*. Publications Puertorriqueñas, xviii. 207 pp.
- Cervený K. 2006. Distribution patterns of reef fishes in southwest Puerto Rico, relative to structural habitat, cross-shelf location, and ontogenetic stage. Department of Marine Sciences, University of Puerto Rico, Mayagüez, Puerto Rico. Masters Thesis. 172 pp.
- Cesar H, Burke L and Pet-Soede L. 2003. *Economics of worldwide coral reef degradation*. World Wildlife Foundation. Veenman Drukkers, Ede. Netherlands. pp. 24.
- Christensen JD, Jeffrey CFG, Caldow C, Monaco ME, Kendall MS and Appeldoorn RS. 2003. Cross-Shelf Habitat Utilization Patterns of reef fishes in southwestern Puerto Rico. *Gulf and Caribbean Research* **14**:9–27.
- Cicchetti G and Greening H. 2011. Estuarine biotope mosaics and habitat management goals: An application in Tampa Bay, FL, USA. *Estuaries and Coasts* **34**:1278–1292.
- Cicchetti G. 2010. Summary of a Technical Workshop held October 28 and 29, 2009: Developing a Biological Condition Gradient for Narragansett Bay. US Environmental Protection Agency, Narragansett, RI. Internal Report.
- Cicchetti G and Pryor M. 2010. Summary of the Estuarine BCG Workgroup November 2008 Workshop, and a Proposed Organizing Framework for Bioassessment of Estuaries. US Environmental Protection Agency, Narragansett, RI. Internal Report. pp. 25.
- Clark HL. 1902. The Echinoderms of Porto Rico. *Bulletin of the United States Fish Commission* **20**:231–263.

- Clark HL. 1933. Handbook of the littoral Echinoderms of Puerto Rico and other West Indian Islands. *New York Academy of Science* **16**:1–147.
- Clark RD, Pittman S, Caldow C, Christensen J, Roque B, Appeldoorn RS and Monaco ME. 2009. Nocturnal fish movement and trophic flow across habitat boundaries in a coral reef ecosystem (SW Puerto Rico). *Caribbean Journal of Science* **45**:282–303.
- Coe WR. 1902. The Nemerteans of Porto Rico. *Bulletin of the United States Fish Commission* **20**:223–229.
- Costanza R, d'Arge R, de Groot R, Farber S, Grasso M, Hannon B, Naeem S, Limburg K, Paruelo J, O'Neill RV, Raskin R, Sutton P and van den Belt M. 1997. The value of the world's ecosystem services and natural capital. *Nature* **387**:253–260.
- Cróquer A and Weil E. 2009. Changes in Caribbean coral disease prevalence after the 2005 bleaching event. *Diseases of Aquatic Organisms* **87**:33–43.
- Dahlgren CP, Kellison T, Adams AJ, Gillanders BM, Kendall MS, Ley JA, Nagelkerken I and Serafy JE. 2006. Marine nurseries and effective juvenile habitats: concepts and applications. *Marine Ecology Progress Series* **312**:291–295.
- Dall WH and Simpson CT. 1902. The Mollusca of Porto Rico. *Bulletin of the United States Fish Commission* **20**:353–524.
- Davies SP and Jackson SK. 2006. The Biological Condition Gradient: a descriptive model for interpreting change in aquatic ecosystems. *Ecological Applications* **16**:1251–1266.
- Debrot A. 2000. A review of records of the extinct W. Indian monk seal. *Marine Mammal Science* **16**:834–837.
- Dennis G. 2000. Annotated checklist of shallow-water marine fishes from the Puerto Rico Plateau including Puerto Rico, Culebra, Vieques, St. Thomas, St. John, Tortola, Virgin Gorda and Anegada. http://www.fcsc.usgs.gov/Marine_Studies/Marine_Puerto_Rico_Plateau/marine_puerto_rico_plateau.html.
- Dennis GD, Hensley DL, Colin PL and Kimmel GG. 2004. New records of marine fishes from the Puerto Rico Plateau. *Caribbean Journal of Science* **40**:70–87.
- Droesen WJ. 1996. Formalisation of ecohydrological expert knowledge applying fuzzy techniques. *Ecological Modeling* **85**:75–81.
- Eakin CM, Feingold JS and Glynn PW. 1994. Oil refinery impacts on coral reef communities in Aruba, N.A. In: Ginsburg RN. *Proceedings of the Colloquium on Global Aspects of Coral Reefs: Health, Hazards and History*. Rosenstiel School of Marine and Atmospheric Science, University of Miami. 1994. pp. 139–145.
- Edmunds PJ. 2013. Decadal-scale changes in the community structure of coral reefs of St. John, US Virgin Islands. *Marine Ecology Progress Series* **489**:107–123.
- Edmunds PJ and Elahi E. 2007. The demographics of a 15-year decline in cover of the Caribbean reef coral *Montastraea annularis*. *Ecological Monographs* **77**: 3–18.
- Federal Register. *Rules and Regulations*. **71**:26852–26861.

- Fisher WS. 2007. *Stony Coral Rapid Bioassessment Protocol*. US Environmental Protection Agency, Office of Research and Development, Washington, D.C. EPA/600/R-06/167, 60 pp.
- Friedlander AM and Parrish JD. 1998. Habitat characteristics affecting fish assemblages on a Hawaiian coral reef. *Journal of Experimental Marine Biology and Ecology* **224**:1–30.
- Garcia JR, Morelock J, Castro R, Goenaga C and Hernandez-Delgado EA. 2003. *Puerto Rican reefs: research synthesis, present threats and management perspectives*. In: *Latin American Coral Reef*. J. Cortés (Ed.). pp. 111–130. Amsterdam: Elsevier Science B.V.
- Garcia-Rios CI (Ed.). 2003. *Los Chitones de Puerto Rico*. Editorial Isla Negra.
- García-Sais J, Appeldoorn R, Battista T, Bauer L, Bruckner A, Caldow C, Carrubba L, Corredor J, Diaz E, Lilyestrom C, García-Moliner G, Hernández-Delgado E, Menza C, Morell J, Pait A, Sabater J, Weil E, Williams E and Williams S. 2008. The State of Coral Reef Ecosystems of Puerto Rico. In: *The state of coral reef ecosystems of the United States and Pacific Freely Associated States: 2008*. Waddell JE and Clarke AM. (Eds). NOAA Technical Memorandum NOS NCCOS 73. 569 pp.
- García-Sais JR, Castro R, Sabater-Clavell J, Esteves R and Carlo M. 2006. Monitoring of coral reef communities from natural reserves in Puerto Rico, 2006: Isla Desecheo, Rincon, Mayagüez Bay, Guánica, Ponce and Isla Caja de Muerto. Final Report submitted to the Department of Natural and Environmental Resources of Puerto Rico. San Juan, PR. 151 pp.
- Gladfelter WB, Ogden JC and Gladfelter EH. 1980. Similarity and diversity among coral reef fish communities: a comparison between Tropical Western Atlantic (Virgin Islands) and Tropical Central Pacific (Marshall Islands) patch reefs. *Ecology* **61**:1156–1168.
- Gladfelter WB. 1982. White-Band Disease in *Acropora palmata*: implications for the structure and growth of shallow reefs. *Bulletin of Marine Science* **32**:639–643.
- Glynn PW. 1988. El-Nino Southern Oscillation 1982–1983. Near shore population, community, and ecosystem responses. *Annual Review of Ecology and Systematics* **19**:309–345.
- Glynn PW. 1991. Coral-reef bleaching in the 1980s and possible connections with global warming. *Trends in Ecology and Evolution* **6**:175–179.
- Goenaga C, Vicente VP and Armstrong RA. 1989. Bleaching Induced Mortalities in Reef Corals from La Parguera, Puerto Rico: A Precursor of Change in the Community Structure of Coral Reefs? *Caribbean Journal of Science* **25**:59–65.
- Goreau TF. 1959. The ecology of Jamaican coral reefs. I. Species composition and zonation. *Ecology* **40**:67–90.
- Goreau TJ, Hayes RH, Clark JW, Basta DJ and Robertson CN. 1992. Elevated Sea Surface Temperatures Correlate with Caribbean Coral Reef Bleaching. In: *A Global Warming Forum: Scientific, Economic, and Legal Overview*. Geyer RA. (Ed.). CRC Press, Boca Raton, Florida USA, Chapter 9, pp. 225–255.
- Grana FA. 1993. *Catalogo de la nomenclatura de los Moluscos de Puerto Rico e Islas Virgenes*. Technical report, Deptmento Recursos Naturales, Puerto Rico.
- Grimsditch GD and Salm RV. 2006. *Coral Reef Resilience and Resistance to Bleaching*. IUCN, Gland, Switzerland. 52pp.

- Gundlach DJ. 1887. Apuntes para la Fauna Puerto-Riqueña (VI). Crustaceos. *Anales de la Sociedad Española de Historia Natural* **16**: 115–133.
- Gundlach J. 1883. Apuntes para la Fauna Puerto-Riqueña (V). Moluscos. *Anales de la Sociedad Española de Historia Natural* **12**:441–484.
- Hargitt CW and Rogers CG. 1902. The Alcyonaria of Puerto Rico. *Bulletin of the United States Fish Commission* **20**:265–287.
- Harvell CD, Altizer S, Cattadori IM, Harrington L and Weil E. 2009. Climate change and wildlife diseases: When does the host matter the most? *Ecology* **90**:912–920.
- Hastie R. 2001. Problems for judgment and decision-making. *Annual Review of Psychology* **52**:653–683.
- Hatcher BG, Imberger J and Smith SV. 1987. Scaling analysis of coral reef systems: An approach to problems of scale. *Coral Reefs* **5**:171–181.
- Hauck F. 1888. Meeresalgen von Puerto-Rico. *Botanische Jahrbücher für Systematik. Pflanzengeschichte und Pflanzengeographie* **9**:457–470.
- Healthy Reefs for Healthy People Initiative. 2012. *Report Card for the Mesoamerican Reef 2012*. 25 pp.
- Hearn CJ, Atkinson MJ and Falter JL. 2001. A physical derivation of nutrient uptakes in coral reefs: Effects of roughness and waves. *Coral Reefs* **20**:347–356.
- Hearn CJ. 2008. *The Dynamics of Coastal Models*. Cambridge University Press, 488 pp.
- Hoegh-Guldberg O. 1999. Climate change, coral bleaching and the future of the world's coral reefs. *Marine and Freshwater Research* **50**:839–866.
- Hubbard DK. 1997. Reefs as dynamic systems. In: *Life and Death of Coral Reefs*. Birkeland C. (Ed.). Chapman and Hall Publishing, New York. pp. 43–67.
- Hughes, TP. 1996. Demographic approaches to community dynamics: a coral reef example. *Ecology* **77**:2256–2260.
- Humann P and DeLoach N. 2002. *Reef Coral Identification*. New World Publications, Inc., Jacksonville, FL. 287 pp.
- Humann P and DeLoach N. 2002. *Reef Fish Identification: Florida, Caribbean, Bahamas*. New World Publications, Inc. 481 pp.
- Humann P and DeLoach N. 2003. *Reef Creature Identification: Florida, Caribbean, Bahamas*. New World Publications, Inc. 420 pp.
- Idjadi JA and Edmunds PJ. 2006. Scleractinian corals as facilitators for other invertebrates on a Caribbean reef. *Marine Ecology Progress Series* **319**:117–127.
- Jackson J, Cramer K, Donovan M, Friedlander A, Hooten A and Lam V. 2012. *Tropical Americas Coral Reef Resilience Workshop*. 29 April–5 May 2012, Tupper Center, Smithsonian Tropical Research Center, Panama City, Republic of Panama.

- Jackson JBC, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, Bradbury RH, Cooke R, Erlandson J, Estes JA, Hughes TP, Kidwell S, Lange CB, Lenihan HS, Pandolfi JM, Peterson CH, Steneck RS, Tegner MJ and Warner RR. 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* **293**:629–637.
- Jackson JBC. 1997. Reefs since Columbus. *Coral Reefs* **16**:S23–S32.
- Johns GM, Leeworthy VR, Bell FW and Bonn MA. 2001. Socioeconomic study of reefs in southeast Florida. Report (Hazen and Sawyer) to Broward County, Palm Beach County, Miami-Dade County (Florida). Florida Fish and Wildlife Commission and the National Oceanic and Atmospheric Administration, 255 pp.
- Karr JR. 1991. Biological Integrity: A Long-Neglected Aspect of Water Quality Resource Management. *Ecological Applications* **1**: 66-84.
- Karr JR and Chu EW. 1999. *Restoring life in running waters: Better biological monitoring*. Island Press, Washington, DC, 206 pp.
- Keeney RL. 1996. Value-focused thinking: Identifying decision opportunities and creating alternatives. *European Journal of Operational Research* **92**:537–549.
- Kendall MS, Monaco ME, Buja KR, Christensen JD, Kruer CR, Finkbeiner M and Warner RA. 2001. *Methods used to map the benthic habitats of Puerto Rico*. NOAA Technical Memorandum NOS NCCOS CCMA 152.
- Kinzie RA III. 1973. The zonation of West Indian gorgonians. *Bulletin of Marine Science* **23**:93–155.
- Leeworthy VR and Bowker JM. 1997. *Nonmarket economic user values of the Florida Keys/Key West*. NOAA, Silver Spring, MD.
- Leeworthy VR and Vanasse P. 1999. *Economic contribution of recreating visitors to the Florida Keys/Key West: updates for years 1996–97 and 1997–98*. NOAA, Silver Spring, MD.
- Leeworthy VR and Wiley PC. 1996. *Importance and satisfaction ratings by recreating visitors to the Florida Keys/Key West*. NOAA, Silver Spring, MD.
- Leeworthy VR and Wiley PC. 1997. *A socioeconomic analysis of the recreation activities of Monroe County residents in the Florida Keys/Key West*. NOAA, Silver Spring, MD.
- Leeworthy VR and Wiley PC. 2003. *Profiles and economic contribution: General visitors to Monroe County, Florida 2001–2001*. NOAA, Silver Spring, MD.
- Lessios H. 2005. *Diadema antillarum* populations in Panama twenty years following mass mortality. *Coral Reefs* **24**:125–127.
- Lessios HA, Robertson DR and Cubit JD. 1984. Spread of *Diadema* mass mortality through the Caribbean. *Science* **226**:335–337.
- Lessios HA. 1988. Mass mortality of *Diadema antillarum* in the Caribbean: What have we learned? *Annual Review of Ecology and Systematics* **19**:371–393.
- Lieske E and Myers R. 2001. *Coral Reef Fishes: Indo-Pacific & Caribbean*. Princeton University Press, Princeton, NJ, USA.

- Long C. 1975. *The Polychaeta from La Parguera, Puerto Rico*. University of Puerto Rico, Department of Marine Sciences.
- Luckhurst BE and Luckhurst K. 1978. Analysis of the influence of substrate variables on coral reef fish communities. *Marine Biology* **49**:317–323.
- Marubini F and Davies PS. 1996. Nitrate increases zooxanthellae population density and reduces skeletogenesis in coral. *Marine Biology* **127**:319–328.
- McClanahan TR, Ateweberhan M, Muhando CA, Maina J and Mohammed MS. 2007. Effects of climate and seawater temperature variation on coral bleaching and mortality. *Ecological Monographs* **77**:503–525.
- McClanahan TR, Weil E, Cortés J, Baird A and Ateweberhan M. 2009. Consequences of coral bleaching for sessile organisms. In M. van Oppen and J. Lough (Eds.) *Coral Bleaching: Patterns, Processes, Causes and Consequences*. *Ecological Studies*. pp: 121–138. Springer-Verlag.
- McField M and Kramer PR. 2007. *Healthy Reefs for Healthy People: A Guide to Indicators of Reef Health and Social Well-being in the Mesoamerican Reef Region*. 208 pp.
- Meissner KJ, Lippmann T and Gupta AS. 2012. Large-scale stress factors affecting coral reefs: Open ocean sea surface temperature and surface seawater aragonite saturation over the next 400 years. *Coral Reefs* **31**: 309–319.
- Menzies RJ and Glynn PW. 1968. The common marine Isopod crustaceans of Puerto Rico. In: *Studies of the Fauna of Curaçao and other Caribbean Islands* **23**:133.
- Meynecke JO, Lee SY and Duke NC. 2008. Linking spatial metrics and fish catch reveals the importance of coastal wetland connectivity to inshore fisheries in Queensland, Australia. *Biological Conservation* **141**:981–996.
- Miller GL and Lugo AE. 2009. *Guide to the ecological systems of Puerto Rico*. General Technical Report. IITF-GTR-35. San Juan, PR: US Department of Agriculture, Forest Service, International Institute of Tropical Forestry. 437 pp.
- Miller J, Muller E, Rogers C, Waara R, Atkinson A, Whelan KRT, Patterson M and Witcher B. 2009. Coral disease following massive bleaching in 2005 causes 60% decline in coral cover on reefs in the US Virgin Islands. *Coral Reefs* **28**:925–937.
- Monismith SG. 2007. Hydrodynamics of coral reefs. *Annual Reviews of Fluid Mechanics* **39**:37–55.
- Moore HF. 1902. Porto Rican Isopoda. *Bulletin of the United States Fish Commission* **20**:161–176.
- Morelock J, Ramirez W, Bruckner A and Carlo M. 2001. Status of coral reefs, southwest Puerto Rico. *Caribbean Journal of Science*, Online Special Publication **4**:57. URL: www.uprm.edu/biology/cjs/reefstatus.htm.
- Mumby PJ, Broad K, Brumbaugh DR, Dahlgren CP, Harborne AR, Hastings A, Holmes KE, Kappel CV, Micheli F and Sanchirico JN. 2008. Coral reef habitats as surrogates of species, ecological functions, and ecosystem services. *Conservation Biology* **22**:941–951.

- Mumby PJ, Edwards AJ, Arias-Gonzalez JE, Lindeman KC, Blackwell PG, Gall A, Gorczynska MI, Harborne AR, Pescod CL, Renken H, Wabnitz CC and Llewellyn G. 2004. Mangroves enhance the biomass of coral reef fish communities in the Caribbean. *Nature* **427**:533–536.
- Munk W and Sargeant M. 1954. *Adjustment of Bikini Atoll to ocean waves*. US Geological Survey Professional Paper 260. pp. 275–280.
- Nagelkerken I, Van der Velde G, Gorissen MW, Meijer GJ, van't Hof T and den Hartog C. 2000. Importance of mangroves, seagrass beds and the shallow coral reef as a nursery for important coral reef fishes, using a visual census technique. *Estuarine, Coastal and Shelf Science* **51**:31–44.
- National Oceanic and Atmospheric Administration (NOAA). 2012a. *Marine Invertebrates and Plants*. URL: www.nmfs.noaa.gov/pr/species/invertebrates/
- National Oceanic and Atmospheric Administration (NOAA). 2012b. *Laboratory for Satellite Altimetry: Sea level rise*. URL: ibis.grdl.noaa.gov/SAT/SeaLevelRise/LSA_SLR_timeseries_global.php.
- National Oceanic and Atmospheric Administration (NOAA). 2013a. *Fish watch, Caribbean Spiny Lobster*. URL: www.fishwatch.gov/seafood_profiles/species/lobster/species_pages/caribbean_spiny_lobster.htm.
- National Oceanic and Atmospheric Administration (NOAA). 2013b. *Elkhorn Coral (Acropora palmata)*. NOAA Fisheries, Office of Protected Resources. URL: www.nmfs.noaa.gov/pr/species/invertebrates/elkhorncoral.htm
- Nunes V and Pawlak G. 2008. Observations of bed roughness of a coral reef. *Journal of Coastal Research* **24**:39–50.
- Odum E. 1997. *Ecology: A bridge between science and society*. Sinauer Associates, Inc., Sunderland, MA. 330 pp.
- Ogden JC. 1997. Ecosystem interactions in the tropical coastal seascape. In: *Life and death of coral reefs*. Birkeland C. (Ed.). Chapman and Hall Publishing. New York. pp. 288–297.
- Ortiz E. 1998. *Los Moluscos recientes de Puerto Rico*. Ph.D. thesis, Department of Marine Sciences, University of Puerto Rico, Mayagüez.
- Osborne PL. 2000. *Tropical Ecosystem and Ecological Concepts*. Cambridge: Cambridge University Press. 464 pp.
- Osburn RC. 1940. Bryozoa of Porto Rico with a resume of the West Indian Bryozoan fauna. *New York Academy of Science* **16**:321–486.
- Oxenford H, Roach R, Brathwaite A, Nurse L, Goodridge R, Hinds F, Baldwin K and Finney C. 2008. Quantitative observations of a major coral bleaching event in Barbados, southeastern Caribbean. *Climatic Change* **87**:435–449.
- Panek FM. 2005. Epizootics and disease of coral reef fish in the tropical western Atlantic and Gulf of Mexico. *Reviews in Fisheries Science* **13**:1–21.
- Park T, Bowker JM and Leeworthy VR. 2002. Valuing snorkeling visits to the Florida Keys with stated and revealed preference models. *Journal of Environmental Management* **65**:301–312.

- Pauly D. 1995. Anecdotes and the shifting base-line syndrome of fisheries. *Trends in Ecology and Evolution* **10**:430.
- Pearce DW and Moran D. 1994. *The Economic Value of Biodiversity*. In association with the Biodiversity Programme, The World Conservation Union. London: Earthscan Publications, 172 pp.
- Pittman SJ, Renchen GF, Clark R, Caldow C, Olsen D and Hill RL. 2011. Applications of estuarine and coastal applications in marine spatial planning. *Treatise on Estuarine and Coastal Science* **1**: 163–205.
- Pittman SJ, Caldow C, Hile SD and Monaco ME. 2007a. Using seascape types to explain the spatial patterns of fish in the mangroves of SW Puerto Rico. *Marine Ecology Progress Series* **348**:273–284.
- Pittman SJ, Christensen JD, Caldow C, Menza C and Monaco ME. 2007b. Predictive mapping of fish species richness across shallow-water seascapes in the Caribbean. *Ecological Modeling* **204**:9–21.
- Pittman SJ, Hile SD, Jeffrey CFG, Caldow C, Kendall MS, Monaco ME, and Hillis-Starr Z. 2008. *Fish assemblages and benthic habitats of Buck Island Reef National Monument (St. Croix, U.S. Virgin Islands) and the surrounding seascape: A characterization of spatial and temporal patterns*. NOAA Technical Memorandum NOS NCCOS 71. Silver Spring, MD. 96 pp.
- Pittman SJ, Hile SD, Jeffrey CFG, Clark R, Woody K, Herlach BD, Caldow C, Monaco ME and Appeldoorn R. 2010. *Coral Reef Ecosystems of Reserva Natural La Parguera (Puerto Rico): Spatial and Temporal Patterns in Fish and Benthic Communities (2001–2007)*. NOAA Technical Memorandum NOS NCCOS 107, Silver Spring, MD. 202 pp.
- Poey F. 1881. Peces. In: Gundlach DJ. Apuntes para la fauna Puerto-Riqueña (III). *Anales de la Sociedad Española de Historia Natural* **10**:317–350.
- Prada C, Weil E and Yoshioka P. 2009. Octocoral bleaching under unusual thermal stress. *Coral Reefs* **29**:41–45.
- Principe P, Bradley P, Yee S, Fisher W, Johnson E, Allen P and Campbell D. 2012. *Quantifying Coral Reef Ecosystem Services*. US Environmental Protection Agency, Office of Research and Development, Research Triangle Park, NC. EPA/600/R-11/206.
- Randall JE. 1967. Food habitats of reef fishes of the West Indies. *Studies of Tropical Oceanography* **5**:665–847.
- Rathbun NJ. 1902. Brachyura and Macrura of Porto Rico. *Bulletin of the United States Fish Commission* **20**:1–127.
- Richmond RH. 1993. Coral reefs: Present problems and future concerns resulting from anthropogenic disturbance. *American Zoologist* **33**:524–536.
- Risk MJ. 1972. Fish diversity on a coral reef in the Virgin Islands. *Atoll Research Bulletin* **153**:1–6.
- Rivero JA. (Ed). 1978. *Los anfibios y reptiles de Puerto Rico*. Editorial Universitaria, Universidad de Puerto Rico.
- Roberts DE, Davis AR and Cummins SP. 2006. Experimental manipulation of shade, silt, nutrients and salinity on the temperate reef sponge *Cymbastela concentrica*. *Marine Ecology Progress Series* **307**:143–154.

- Rouse I. 1992. *The Tainos: Rise and Decline of the People Who Greeted Columbus*. Yale University Press. 232 pp.
- Ruiz Torres H. 2012. *Beneath the Waves*. University of Puerto Rico, Sea Grant. 131 pp.
- Ruzicka RR, Colella MA, Porter JW, Morrison JM, Kidney JA, Brinkhuis V, Lunz KS, Macaulay KA, Bartlett LA, Meyers MK and Colee J. 2013. Temporal changes in benthic assemblages on Florida Keys reefs 11 years after the 1997/1998 El Niño. *Marine Ecology Progress Series* **489**:125–141.
- Sale PF and Szmant AM. (Eds). 2012. *Reef Reminiscences: Ratcheting back the shifted baselines concerning what reefs used to be*. United Nations University Institute for Water, Environment and Health, Hamilton, ON, Canada, 35 pp.
- Sale PF, Jacob P and Kritzer JP. 2008. Connectivity: What it is, how it is measured, and why it is important for management of reef fishes. In: Grober-Dunsmore R and Keller BD. (Eds). Caribbean connectivity: Implications for marine protected area management. *Proceedings of a Special Symposium, 9–11 November 2006, 59th Annual Meeting of the Gulf and Caribbean Fisheries Institute, Belize City, Belize*. Marine Sanctuaries Conservation Series ONMS-08-07. Silver Spring (MD): US Department of Commerce, National Oceanic and Atmospheric Administration, Office of National Marine Sanctuaries, pp. 16–30.
- Santavy DL, Fisher WS, Campbell JG and Quarles RL. 2012. *Field Manual for Coral Reef Assessments*. US Environmental Protection Agency, Office of Research and Development, Gulf Ecology Division, Gulf Breeze, FL. EPA/600/R-12/029. 92 pp.
- Santavy DL, Courtney LA, Fisher WS, Quarles RL and Jordan SJ. 2013. Estimating surface area of sponges and gorgonians as indicators of habitat availability on Caribbean coral reefs. *Hydrobiologia* **707**:1–16.
- Sebens KP. 1994. Biodiversity of coral reefs: What we are losing and why? *American Zoologist* **34**:115–133.
- Sefton N and Webster SV. 1986. *Caribbean Reef invertebrates*. Sea Challengers. Monterey, CA. 112 pp.
- Shashar N, Kinane S, Jokiel PL and Patterson MR. 1996. Hydromechanical boundary layers over a coral reef. *Journal of Experimental Marine Biology and Ecology* **199**:17–28.
- Shivlany M, Leeworthy VR, Murray TJ, Suman DO and Tonioh F. 2008. *Knowledge, attitudes and perceptions of management strategies and regulations of the Florida Keys National Marine Sanctuary by commercial fishers, dive operators, and environmental group members: A baseline characterization and 10-year comparison*. National Oceanic and Atmospheric Administration, Silver Spring, MD.
- Soler-López LR. 2001. Sedimentation Survey Results of the Principal Water Supply Reservoirs of Puerto Rico. In: Sylva WF (Ed.). *Proceedings of the Sixth Caribbean Islands Water Resources Congress, Mayagüez, Puerto Rico, February 22 and 23, 2001*.
- Stahl A. 1883. *Fauna de Puerto Rico, Clasificación sistemática de lo animales que corresponden á esta fauna, y catálogo del gabinete zoológica del Dr. A. Stahl en Bayamon*. San Juan, PR. 249 pp.

- Steele MA. 1999. Effects of shelter and predators on reef fishes. *Journal of Experimental Marine Biology and Ecology* **233**:65–79.
- Stoddard JL, Larsen DP, Hawkins CP, Johnson RK and Norris RH. 2006. Setting expectations for the ecological condition of streams: The concept of reference condition. *Ecological Applications* **16**:1267–1276.
- Sturm P, Viqueira R, Ferguson R and Moore T. 2012. Addressing land based sources of pollution in Guánica, Puerto Rico. In: D. Yellowlees and TP Huges, *Proceedings of the 12th International Coral Reef Symposium, 9-13 July 2012 Cairns, Australia*. URL:<http://www.icrs2012.com/Proceedings.htm>
- Talbot FH. 1965. A description of the coral structure of Tutia reefs (Tanganyika territory, East Africa) and its fish fauna. *Proceedings of the Zoological Society of London* **145**:431–470.
- Thacker RW and Freeman CJ. 2012. Chapter two - Sponge-Microbe Symbioses: Recent Advances and New Directions. In: *Advances in Marine Biology*. Becerro MA, Uriz MJ, Maldonado M and Turon X. (Eds.), Academic Press **62**:57–111.
- The Nature Conservancy (TNC). 2006. *Are Florida's reefs resilient? A guide to the Florida reef resilience program*. Summerland Key, Florida.
- Torres JL and Morelock J. 2002. Effect of terrigenous sediment influx on coral cover and linear extension rates of three Caribbean massive coral species. *Caribbean Journal of Science* **38**:222–229.
- Treadwell AL. 1902. Polychaetous annelids of Puerto Rico. *Bulletin of the United States Fish Commission* **20**:181–210.
- Treadwell AL. 1939. Polychaetous annelids of Puerto Rico and Vicinity. In: *Scientific Survey of Porto Rico and the Virgin Islands* **16**:313. New York Academy of Sciences.
- Turner RK, Brouwer R, Georgiou S and Bateman IJ. 2000. *Ecosystem functions and services: An integrated framework and case study for environmental valuation*. Norwich (UK): University of East Anglia, The Centre for Social and Economic Research on the Global Environment (CSERGE), CSERGE Working Paper GEC 2000-21, 32 pp.
- US Commission on Ocean Policy. 2004. *US Ocean Action Plan*. 41 pp.
- US Environmental Protection Agency (EPA). 2005. *Use of Biological Information to Better Define Designated Aquatic Life Uses in State and Tribal Water Quality Standards: Tiered Aquatic Life Uses*. EPA-822-R-05-001.
- US Environmental Protection Agency (EPA). 2011. *A Primer on Using Biological Assessments to Support Water Quality Management*. EPA 810-R-11-001. Washington, DC.
- US Environmental Protection Agency (EPA). In review. *A BCG Framework for Bioassessment of Estuaries and Coasts*.
- Vaughan TW. 1902. The Stony Corals of Porto Rico. *Bulletin of the United States Fish Commission* **20**:289-320.
- Velazco-Domínguez AT, Weil E and Bruckner A. 2003. *Climate Change and Coral Bleaching in Puerto Rico: Efforts and Challenges*. Presentation given in Oahu, Hawaii.

- Vicente VP. 1990. Response of sponges with autotrophic endosymbionts during the coral-bleaching episode in Puerto Rico. *Coral Reefs* **8**:199–202.
- Vicente VP. 1992. A summary of ecological information on the seagrass beds of Puerto Rico. In: *Coastal Plant Communities of Latin America*, Seliger E. (Ed.). Academic Press, New York, pp. 123–133.
- Vicente VP and Goenaga C. 1984. Mass mortalities of the sea urchin *Diadema antillarum* (Philippi) in Puerto Rico. *CEER- M*-**195**:1–30.
- Wagner DE, Kramer P and Woesik RV. 2010. Species composition, habitat, and water quality influence coral bleaching in southern Florida. *Marine Ecology Progress Series* **408**:65–78.
- Walker BK, Jordan LKB and Spieler RE. 2009. Relationship of reef fish assemblages and topographic complexity on southeastern Florida coral reef habitats. *Journal of Coastal Research* **53**:39–48.
- Warne AG, Webb RMT and Larsen MC. 2005. *Water, sediment, and nutrient discharge characteristics of rivers in Puerto Rico and their potential influence on coral reefs*. USGS Science Investigative Report 2005–5206.
- Warner GF, Smith SR, Jordan-Dahlgren E, Linton DM, Woodley JD, Alcolado P, Bonaire K, Bone D, Buchan KC, Bush P, Cortés J, Croquer A, De Meyer K, Fernandez RG, Fonseca A, Garcia JR, Garcia-Parrado P, Garzón-Ferreira J, Gayle P, Gerace DT, Gerald FX, Gunther J, Guppy R, Juman R, Koltes KH, Knobbe E, Klein E, Laydoo R, Losada F, Menendez G, Mow-Robinson JM, Ostrander G, Oxenford HA, Parker C, Pors LPJJ, Perez D, Ramirez AR, Rodriguez R, Ruiz-Rentaria F, Ryan J, Tschirky JJ and Weil E. 2002. Status and temporal trends at CARICOMP coral reef sites. *Proceedings of the 9th International Coral Reef Symposium, Bali, Indonesia*. pp. 325–330.
- Warren-Rhodes K, Sadovy Y and Cesar HSJ. 2003. Marine ecosystem appropriation in the Indo-Pacific: A case study of the live reef fish food trade. *Ambio* **32**:481–488.
- Weil E, Hernandez-Delgado EA, Bruckner AW, Ortiz AL, Nemeth M and Ruiz H. (2003). Distribution and status of Acroporid coral (*Scleractinia*) populations in Puerto Rico. *Proceedings of the Caribbean Workshop: Potential Application of the US Endangered Species Act (ESA) as a Conservation Strategy*. NOAA-NMFS and NCORE-RSMAS. University of Miami. pp. 71–92.
- Weil E. 2005. Current Status of the Marine Biodiversity of Puerto Rico. In: Miloslavich P and Klein E. (Eds). *Caribbean Marine Biodiversity: The known and unknown*, pp. 85–109. DEStech Publications Inc., Lancaster, PA, USA.
- Weil E, Torres JL and Ashton M. 2005. Population characteristics of the black sea urchin *Diadema antillarum* (Philippi) in La Parguera, Puerto Rico, 17 years after the mass mortality event. *Revista de Biología Tropical* **53**:219–231.
- Weil E, Cróquer A and Urreiztieta I. 2009. Temporal variability and consequences of coral diseases and bleaching in La Parguera, Puerto Rico from 2003-2007. *Caribbean Journal of Science* **45**:221–246.
- Weil E and Rogers CS. 2011. Coral reef disease in the Atlantic-Caribbean. In Z. Dubinski and N. Stambler. (Eds.). *Coral Reefs: An Ecosystem in Transition*. Chapter 27. pp. 465–492. Springer-Verlag.

- Wells JW. 1957. Scleractinia. In: *Treatise on invertebrate paleontology. Part F. Coelentrata*. Moore RC. (Ed.). Geological Society of America, Boulder, Colorado. pp. 328–444.
- Wikipedia. 2013. *Central Guánica*. URL: http://en.wikipedia.org/wiki/Central_Gu%C3%A1nica.
- Wilkinson C (Ed.). 2004. *Status of coral reefs of the world: 2004. Vol.1*. Townsville, Queensland, Australia: Australian Institute of Marine Science. 316pp.
- Wilkinson CR. 1983. Net primary productivity in coral reef sponges. *Science* **219**:410–412.
- Wilkinson CR. 1996. Global change and coral reefs: Impacts on reefs, economies and human cultures. *Global Change Biology* **2**:547–558.
- Williams EH Jr. and Bunkley-Williams L. 1989. Bleaching of Caribbean coral reef symbionts in 1987–1988. *Proceedings of the 6th International Coral Reef Symposium* **3**:313–318.
- Williams EH Jr. and Bunkley-Williams L. 1990. The world-wide coral reef bleaching cycle and related sources of coral mortality. *Atoll Research Bulletin* **335**:1–71.
- Williams EH Jr., Goenega C and Vicente V. 1987. Mass bleaching on Atlantic coral reefs. *Science* **237**:877–878.
- Wilson HV. 1902. The sponges collected in Porto Rico in 1899 by the US Fish Commission 1900. *Bulletin of the United States Fish Commission* **20**:375–411.
- Yoder CO and DeShon JE. 2003. Using biological response signatures within a framework of multiple indicators to assess and diagnose causes and sources of impairments to aquatic assemblages in selected Ohio rivers and streams. In: *Biological Response Signatures: Indicator Patterns Using Aquatic Communities*. Simon TP. (Ed.). CRC Press, Boca Raton, FL. pp. 23–81.
- Yoder CO and Rankin ET. 1995a. Biological criteria program development and implementation in Ohio. In: *Biological assessment and criteria: Tools for water resource planning and decision making*. Davis WS and Simon TP. (Eds.). Lewis Publishers, Boca Raton, FL. pp. 109–144.
- Yoder CO and Rankin ET. 1995b. Biological response signatures and the area of degradation value: New tools for interpreting multimetric data. In: *Biological Assessment and Criteria: Tools for Water Resource Planning and Decision Making*. Davis WS and Simon TP. (Eds.). Lewis Publishers, Boca Raton, FL. pp. 236–286.
- Yoshioka PM. 2009. Sediment transport and the distribution of shallow water gorgonians. *Caribbean Journal of Science* **45**:254–259.
- Yoshioka PM and Yoshioka BB. 1989a. A multispecies, multiscale analysis of spatial pattern and its application to a shallow-water gorgonian community. *Marine Ecological Progress Series* **54**:257–264.
- Yoshioka PM and Yoshioka BB. 1989b. Effects of wave energy, topographic relief and sediment transport on the distribution of shallow-water gorgonians of Puerto Rico. *Coral Reefs* **8**:145–152.
- Zawada DG. 2011. Reef Topographic Complexity. In: *Encyclopedia of Modern Coral Reefs: Structure, Form and Process*. Hopley D. (Ed.). pp. 902–906.

Appendix B. Workshop Participants

Mr. Aaron Hutchins
The Nature Conservancy
3052 Estate Little Princess
Christiansted, St. Croix, VI 00820
340-718-5575
ahutchins@tnc.org

Dr. Alberto Sabat
University of Puerto Rico, Dept. of Biology
P.O. Box 23360
Rio Piedras, PR 00931-3360
787-764-0000 x2113
amsabat@gmail.com

Dr. Alina Szmant
University of North Carolina, Wilmington
Center for Marine Science
5600 Marvin K. Moss Lane
Wilmington, NC 28409
910-962-2362
szmanta@uncw.edu

Ms. Antares Ramos Alvarez
NOAA Office of Ocean and
Coastal Resource Mgmt.
654 Muñoz Avenue, Suite 604
San Juan, PR 00918
787-766-5206 ext. 224
Antares.ramos@noaa.gov

Ms. Brandi Todd
US EPA, Region 6
1445 Ross Ave.
Dallas, TX 75202
214-665-2233
todd.brandi@epa.gov

Dr. David Ballantine
University of Puerto Rico
Department of Marine Sciences
P.O. Box 9000
Mayagüez, PR 00681-9013
david.ballantine@upr.edu

Dr. David Cuevas
US EPA, Region 2
Caribbean Environ. Prot. Div.
City View Plaza II - Suite 7000
#48 RD. 165km 1.2
Guaynabo, PR 00968-8069
787-977-5856
cuevas.david@epa.gov

Dr. Deborah Santavy (Organizer)
US EPA, GED
1 Sabine Island Dr.
Gulf Breeze, FL 32561
850-934-9358
santavy.debbie@epa.gov

Mr. Ernesto Diaz
PR Department of Natural and Environment
Resources (DNER)
Coastal Zone Management Director
P.O. Box 366147
San Juan, PR 00936
787-999-2200 x2729
ediaz@drna.gobierno.pr

Dr. Ernesto Weil
University of Puerto Rico,
Mayagüez Campus
P.O. Box 908
Lajas, PR 00667-0908
787-899-2048 x241, x272
eweil@caribe.net

Dr. Francisco Pagan
Caribbean Coral Reef Institute
University of Puerto Rico
Mayagüez, PR 00681-9013
Francisco.pagan@upr.edu

Mr. Hector Ruiz Torres
University of Puerto Rico
Department of Marine Sciences
Mayagüez, PR 00681-9013
787-691-7410
hectorruizt@me.com

Mr. Jeff Miller
Virgin Islands National Park
1300 Cruz Bay Creek
St. John, VI 00830
340-693-8950 x227
William_J_Miller@nps.gov

Dr. Jeroen Gerritsen (Facilitator)
Aquatic Ecologist
Tetra Tech, Center for Ecological Sciences
400 Red Brook Blvd., Suite 200
Owings Mills, MD 21117
410-356-8993
jeroen.gerritsen@tetratech.com

Dr. Jorge Bauzá
San Juan Bay Estuary Program
P.O. Box 9509
San Juan, PR 00908-9509
787-638-9979
jbauza@estuario.org

Dr. Loretta Roberson
University of Puerto Rico
College of Natural Sciences
Ponce de Leon Ave.
Rio Piedras, PR 00931-3300
787-764-0000 x1-2713
Loretta.Roberson@gmail.com

Dr. Melanie McField
Smithsonian Institution
Director, Healthy Reefs for Healthy People
1755 Coney Dr.
Belize City, Belize, Central America
501-223-4898
mcfield@healthyreefs.org

Mr. Miguel Canals
PR DNER
Guánica State Forest
P.O. Box 1185
Guánica, PR 00653
787-821-5706
menqui@hotmail.com

Ms. Patricia Bradley (Organizer)
US EPA, AED
33 East Quay Road
Key West, FL 33040
305-809-4690
bradley.patricia@epa.gov

Dr. Paul Yoshioka
University of Puerto Rico
Department of Marine Sciences
Mayagüez, PR 00681-9000
Paul.yoshioka@upr.edu

Dr. Richard Appeldoorn
University of Puerto Rico
Department of Marine Sciences
Mayagüez, PR 00681-9013
787-899-2048 x251
Richard.appeldoorn@upr.edu

Mr. Roberto Viqueira
Protectores de Cuencas
Guánica Coordinator
Box 673
Yauco, PR 00698
787-457-8803
rviqueira@hotmail.com

Dr. Tyler Smith
University of the Virgin Islands
#2 John Brewer's Bay
St. Thomas, VI 00802-9990
340-693-1394
tsmith@uvi.edu

Dr. Vance Vicente
Vicente and Associates, Inc.
Garden Hills Pz 1353 19
Guaynabo, PR 00966
787-781-6503
vance@prtc.net

Dr. William Fisher
US EPA, GED
1 Sabine Island Dr.
Gulf Breeze, FL 32561
850-934-9394
Fisher.william@epa.gov

Appendix C. Workshop Agenda

(Workshop was compressed to two days due to Tropical Storm Isaac; times shown below are approximate)

Goal: To develop a conceptual, narrative model that describes how biological attributes of coral reefs change along a gradient of increasing anthropogenic stress.

DAY 1 – Setting the Stage: A Visual Evaluation of Stations

9:00 **Registration**

9:30 **Purpose of the Workshop**

9:45 **Introductions**

Purpose: Who is attending; organization they represent; what scientific expertise?

Desired Outcomes: Relaxed atmosphere, prepare to work as a team.

10:00 **Coral Reef Video Evaluations**

Purpose: Participants individually review coral reef videos, located throughout the 3 rooms in the conference center.

Desired Outcomes: Every participant will have evaluated 8 videos.

12:00 **Lunch**

1:00 **Complete Coral Reef Video Evaluations**

Purpose: Complete final 4 stations.

Desired Outcomes: Participants have rated EPA stations and documented their rationale.

2:00 **Break**

2:15 **Freshwater Stream and Estuarine Attributes**

Presenter: Jeroen Gerritsen (Brief introduction to the attributes developed for freshwater streams and estuaries).

Purpose: Introduce the stream and estuarine attributes.

Desired Outcomes: Understand where others have been and where we hope to go. Further explore the attribute concept.

2:45 **Presentation of Ratings and Discussion of Rationale**

Presenter: Debbie Santavy (ranked stations)

Purpose: Try to reach consensus on station assignments stating the rationale for the decision.

Desired Outcomes: Share rating of stations; document criteria considered during selection and capture on flip charts.

4:30 **Thresholds**

Presenter: Jeroen Gerritsen (management uses of BCG and how to move forward).

Purpose: Establish preliminary thresholds for different levels of conceptual model.

Desired Outcomes: What relative abundance of sessile invertebrates for each level: hard corals, sponges and gorgonians? What else defines each level? Fish, rugosity, other invertebrates? Any inclusion of water quality factors: both qualitative and quantitative.

5:15 **Adjourn**

DAY 2 – Biological Integrity

9:00 **Biological Integrity Discussion**

Presenters: Debbie Santavy (results from coral reef video evaluations and attributes discussion); Experts (share their videos and photos that exhibit full biological integrity of a coral reef); Pat Bradley (reference condition); Pat Bradley (list of coral reef taxa).

Purpose: Discuss biological integrity and reference condition.

10:30 **Break**

10:45 **Reference Condition Discussion**

Desired Outcomes: Preliminary consensus on what the reference station should be. Begin to assemble the attributes.

12:30 **Lunch**

1:30 **Using Data to Rank Stations**

Presenter: Debbie Santavy (overview of EPA data).

Purpose: Focus thinking about the attributes in breakout groups.

Desired Outcomes: Begin thinking about levels of condition and lists of associated attributes.

3:00 **Break**

3:15 **Attributes as Condition Changes**

Purpose: Begin to consider different levels of condition along a human disturbance gradient, using visual and data-derived attributes.

Desired Outcomes: Begin to compile lists of both visual and data-derived attributes that are not station specific, but more overarching characteristics. Perhaps develop levels of attributes from data metrics to begin populating BCG framework.

5:00 **Thank you and next steps**

5:30 **Adjourn**

Appendix D.

Tally Sheet – Rating Condition of Coral Reef Videos (1st)

Name: _____

Ballot

Station No.	Rating (Good, Fair, Poor)	Rationale (indicate 3 most important characteristics considered in ranking)
1		
2		
3		
4		
5		
6		
7		
8		

Appendix E.

Tally Sheet – Rating Condition of Coral Reef Videos (2nd)

Name: _____

Ballot

Station No.	Rating (Good, Fair, Poor)	Rationale (indicate 3 most important characteristics considered in ranking)
9		
10		
11		
12		

Appendix F.

Notes Sheet – Rating Condition of Coral Reef Videos

Notes Sheet - **Station 1**

Rating: **Good** - - - **Fair** - - **Poor**
(Circle your condition rating)





Use this sheet to capture salient points about this station while viewing the video. You also have a photo handout of key photos to assist you. The salient points should provide your rationale for rating condition as good, fair or poor.

Appendix G.






Supporting Photos – Rating Condition of Coral Reef Videos

The following pages show supporting photos for each station (1 page per station). The experts used these as supplemental material to evaluate the videos.


Station 1




Station 2






Station 3






Station 4









Station 5







Station 6








Station 7






Station 8





Station 9



Station 10



Station 11



Station 12



Appendix H. Workshop Glossary

Attribute: Any measurable component of a biological system (Karr and Chu 1999).

Best attainable condition: A condition that is equivalent to the ecological condition of (hypothetical) least disturbed stations where the best possible management practices are in use. This condition can be determined using techniques such as historical reconstruction, best ecological judgment and modeling, restoration experiments, or inference from data distributions.

Biological integrity: The ability of an aquatic ecosystem to support and maintain a balanced, adaptive community of organisms having a species composition, diversity, and functional organization comparable to that of natural habitats within a region.

Human disturbance: Human activity that alters the natural state and can occur at or across many spatial and temporal scales.

Ecosystem-level functions: Processes performed by ecosystems, including, among other things, primary and secondary production, respiration, nutrient cycling, and decomposition (EPA 2005).

Historical condition: The ecological condition at some previous point in history. Conditions reflective of the historic time period may no longer exist in actual ecosystems in an area.

Least disturbed condition: The best available existing conditions with regard to physical, chemical, and biological characteristics or attributes of a waterbody within a class or region. These waters have the least amount of human disturbance in comparison to others within the waterbody class, region or basin. Least disturbed conditions can be readily found but may depart significantly from natural, undisturbed conditions or minimally disturbed conditions. Least disturbed condition may change significantly over time as human disturbances change (EPA 2005).

Minimally disturbed condition: The physical, chemical and biological conditions of a waterbody with very limited or minimal human disturbance in comparison to others within the waterbody class or region. Minimally disturbed conditions can change over time in response to natural processes (EPA 2005).

Non-native species: Any species that is not naturally found in that ecosystem. Species introduced or spread from one region of the US to another outside their normal range are non-native or non-indigenous, as are species introduced from other continents (EPA 2005).

Reference condition: The condition that approximates natural, unimpacted conditions (biological, chemical, physical, etc.) for a waterbody. Reference condition (biological integrity) is best determined by collecting measurements at a number of stations in a similar waterbody class or region under undisturbed or minimally disturbed conditions (by human activity), if they exist. Since undisturbed or minimally disturbed conditions may be difficult or impossible to find, least disturbed conditions combined with historical information, models or other methods, may be used to approximate reference condition as long as the departure from natural or ideal is understood. Reference condition is used as a benchmark to determine how much other water bodies depart from this condition due to human disturbance (EPA 2005).

Reference station: A station selected for comparison with stations being assessed. The type of stations selected and the type of comparative measures used will vary with the purpose of the comparisons. For the purposes of assessing the ecological condition of stations, a reference station is a specific locality on a waterbody that is undisturbed or minimally disturbed and is representative of the expected ecological integrity of other localities on the same waterbody or nearby waterbodies (EPA 2005).

Sensitive-rare taxa: Taxa that naturally occur in low numbers relative to total population density but may make up large relative proportion of richness. May be ubiquitous in occurrence or may be restricted to certain microhabitats, but because of low density recorded occurrence is dependent on sample effort. Often stenothermic (having a narrow range of thermal tolerance) or cold-water obligates, commonly k-strategists (populations maintained at a fairly constant level, slower development, longer life-span), may have specialized food resource needs or feeding strategies. Generally intolerant to significant alteration of the physical or chemical environment; are often the first taxa observed to be lost from a community (EPA 2005).

Sensitive or regionally endemic taxa: Taxa with restricted, geographically isolated distribution patterns (occurring only in a locale as opposed to a region), often due to unique life history requirements. May be long lived, late maturing, low fecundity, limited mobility or require mutualistic relationships with other species. May be listed as threatened, endangered or of special concern species. Predictability of occurrence often low, therefore, requires documented observation. Recorded occurrence may be highly dependent on sample methods, station selection and level of effort (EPA 2005).

Sensitive taxa: Taxa that are intolerant to a given anthropogenic stress, often the first species affected by the specific stressor to which they are "sensitive" and the last to recover following restoration (EPA 2005).

Taxa: A grouping of organisms given a formal taxonomic name such as species, genus, family, etc. (EPA 2005).

Taxa of intermediate tolerance: Taxa that comprise a substantial portion of natural communities, which may increase in number in waters which have moderately increased organic resources and reduced competition, but they are intolerant of excessive pollution loads or habitat alteration. These may be r-strategists (early colonizers with rapid turn-over times; boom/bust population characteristics), eurythermal (having a broad thermal tolerance range), or have generalist or facultative feeding strategies enabling them to utilize more diversified food types. They are readily collected with conventional sample methods (EPA 2005).

Tolerant taxa: Taxa that comprise a low proportion of natural communities. Tolerant taxa often are tolerant of a broader range of environmental conditions and are thus resistant to a variety of pollution or habitat-induced stress. They may increase in number (sometimes greatly) in the absence of competition. They are commonly r-strategists (early colonizers with rapid turn-over times; boom/bust population characteristics), able to colonize when stress conditions occur. Last survivors (EPA 2005).

Appendix I. Summary Data Results for BCG Stations

Table I-1. Scleractinian coral summary statistics for BCG stations (Puerto Rico surveys in 2010 and 2011). Ave. 3D SA is average 3-dimensional surface area cm^2/m^2 . SE=standard error of mean.

BCG Station No.	EPA Station No.	Species Richness	% Total Abundance	No. Colonies	Shannon Diversity Index (H')	Colony Density ($\#/\text{m}^2$)	3D Coral Area (cm^2/m^2)	2D Coral Area (cm^2/m^2)	Rugosity*	SE Rugosity	Rating by Visual Media	Visual Rating
1	125	13	11.61	51	2.040	3.40	6,560	1,533	2.18	0.107	poor/fair	8
2	15	3	2.68	4	1.040	0.16	185	106	1.12	0.019	poor (worst)	12
3	113	10	8.93	79	1.657	3.16	9,518	2,843	1.52	0.107	good (best)	1
4	3	6	5.36	21	1.234	0.84	2,706	565	1.52	0.087	poor	11
5	19	7	6.25	31	1.261	1.24	18,228	3,673	1.54	0.108	fair/poor	9
6	14	10	8.93	54	1.793	3.6	5,359	1,530	1.71	0.106	poor	10
7	16	11	9.82	73	1.971	4.87	14,210	5,089	1.83	0.119	fair	4
8	108	7	6.25	71	1.512	4.73	19,637	4,549	1.48	0.116	fair	6
9	109	8	8.04	87	1.410	5.80	20,080	5,917	1.88	0.134	fair	7
10	1	11	9.82	70	1.784	2.8	11,026	3,698	1.48	0.052	fair	3
11	46	9	8.04	44	1.827	1.16	11,635	2,598	1.12	0.039	fair	5
12	25	8	7.14	95	1.469	3.8	9,199	3,498	1.25	0.086	fair	2

*Rugosity is the linear ratio of 6m divided by the taut linear distance of a 6m chain draped over the tops of corals and along the bottom. Rugosity is a reef-scale indicator of reef contour or surface heterogeneity. See Appendix J for formulas.

Table I-2. Gorgonian summary statistics for BCG stations (Puerto Rico surveys in 2010 and 2011). Ave. 3D SA is average 3-dimensional surface area cm^2/m^2 or per individual. Maximum number of morphologies that can be present at one station is nine.

BCG Station No.	Station No.	Morpho. Richness ^a	No. Individuals	Density $\#/\text{m}^2$	Ave. 3D SA ^b / m^2	Ave. 3D SA ^c /ind
1	125	4	21	4.2	5,154	1,227
2	15	0	0	0	0	0
3	113	7	24	4.8	30,346	6,322
4	3	0	0	0	0	0
5	19	0	0	0	0	0
6	14	6	27	5.4	38,352	7,102
7	16	8	37	7.4	33,342	4,506
8	108	2	6	1.2	86	71
9	109	4	7	1.4	11,229	8,021
10	1	2	10	2	17,649	8,825
11	46	7	52	10.4	26,954	2,592
12	25	8	86	17.2	58,558	3,405

a: Morphological shapes and regression equations for 3D surface estimation of an individual by morphology in Santavy et al., 2012, pp. 36-38.

b: Ave. 3D SA/ m^2 = Σ Gorgonian surface area in transect area/total transect area

c: Ave. 3D SA/ind = Σ Gorgonian surface area in transect area/total # Gorgonians in transect. See Appendix J, Table J-2 for formulas

Table I-3. Sponge summary statistics for BCG stations (Puerto Rico surveys in 2010 and 2011). Ave. 3D SA is average 3-dimensional surface area cm^2/m^2 or per individual. Maximum number of morphologies present at one station is eight.

BCG Station No.	Station No.	Morpho. Richness ^a	No. Individuals	Density $\#/\text{m}^2$	Ave. 3D SA ^b / m^2	Ave. 3D SA ^c /ind
1	125	3	22	4.4	1,615	367
2	15	3	20	4	1,146	286
3	113	1	5	1	48	48
4	3	2	7	1.4	173	124
5	19	1	5	1	208	208
6	14	2	8	1.6	410	256
7	16	0	0	0	0	0
8	108	0	0	0	0	0
9	109	3	7	1.4	1,389	992
10	1	5	33	6.6	10,776	1,633
11	46	3	21	4.2	53,824	12,815
12	25	5	28	5.6	6,213	1,110

a: Morphological shapes and regression equations for 3D surface estimation of an individual by morphology in Santavy et al., 2012, pp. 36-38.

b: Ave. 3D SA/ $\text{m}^2 = \Sigma$ Sponge surface area in transect area/total transect area.

c: Ave. 3D SA/ind = Σ Sponge surface area in transect area/total # Sponges in transect. See Appendix J, Table J-2 for formulas.




Figure I-1. Comparison of the density of the major sessile invertebrates assessed in the 12 BCG stations.




Figure I-2. Comparison of density for fish carnivores vs. herbivores at BCG stations. Number above bar pairs is the ratio of carnivore/herbivore density.




Figure I-3. Comparison of biomass for fish carnivores vs. herbivores at BCG stations. Number above bar pairs is the ratio of carnivore/herbivore biomass.

BCG Station 1
(Field Station 125_2010)
Coral Species Richness: 13
Photo rank and rating: 8, Poor/Fair

Table I-4. BCG Station 1 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	3.40	0.53	0.13	0.13	0.73
Ave. 3D colony skeletal area (cm ²)/colony	2,146	2,462	20,077	417	435
Ave. 3D colony skeletal area (cm ²)/m ²	7,297	1,313	2,677	56	319
Ave. 3D coral tissue area (cm ²)/colony	1,929	2,284	16,446	417	435
Ave. 3D coral tissue area (cm ²)/m ²	6,560	1,218	2,193	56	319
Ave. 2D coral tissue area (cm ²)/m ²	1,533	228	198	54	16




Figure I -4. Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 1.

Table I-5. Fish species found in BCG Station 1, with density and biomass for 100 m² transect.


Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Abudefduf saxatilis</i>	Sergeant Major	2	90
<i>Acanthurus coeruleus</i>	Blue Tang	5	167
<i>Anisotremus surinamensis</i>	Black Margate	1	1,274
<i>Anisotremus virginicus</i>	Porkfish	1	9
<i>Caranx ruber</i>	Bar Jack	1	17
<i>Gramma loreto</i>	Fairy Basslet	3	1
<i>Haemulon flavolineatum</i>	French Grunt	1	40
<i>Halichoeres poeyi</i>	Blackear Wrasse	1	5
<i>Holacanthus bermudensis</i>	Blue Angelfish	1	11
<i>Lutjanus analis</i>	Mutton Snapper	1	637
<i>Lutjanus apodus</i>	Schoolmaster	1	215
<i>Microspathodon chrysurus</i>	Yellowtail Damselfish	5	174
<i>Ocyurus chrysurus</i>	Yellowtail Snapper	8	242
<i>Ophioblennius macclurei</i>	Redlip Blenny	3	12
<i>Pempheris schomburgkii</i>	Glassy Sweeper	1	33
<i>Scarus iseri</i>	Striped Parrotfish	2	15
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	1	40
<i>Sparisoma viride</i>	Stoplight Parrotfish	3	3
<i>Stegastes adustus</i>	Dusky Damselfish	49	510
<i>Thalassoma bifasciatum</i>	Bluehead	55	198

BCG Station 2
(Field Station 15_2011)
Coral Species Richness: 3
Photo rank and rating: 12, Poor (Worst)

Table I-6. BCG Station 2 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	0.16	0	0	0.08	0
Ave. 3D colony skeletal area (cm ²)/colony	1,248	0	0	980	0
Ave. 3D colony skeletal area (cm ²)/m ²	200	0	0	78	0
Ave. 3D coral tissue area (cm ²)/colony	1,157	0	0	822	0
Ave. 3D coral tissue area (cm ²)/m ²	185	0	0	66	0
Ave. 2D coral tissue area (cm ²)/m ²	106	0	0	37	0

Coral Species



Sponge Morphs




Figure I-5. Percentages of stony coral species and sponge morphologies for BCG Station 2.
 No gorgonians were present at BCG Station 2.

Table I-7. Fish species found in BCG Station 2, with density and biomass for 100 m² transect.

Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Acanthurus bahianus</i>	Ocean Surgeonfish	10	457
<i>Anisotremus virginicus</i>	Porkfish	5	220
<i>Canthigaster rostrata</i>	Sharpnose Puffer	1	0
<i>Cephalopholis fulva</i>	Coney	2	68
<i>Chaetodon capistratus</i>	Foureye Butterflyfish	7	105
<i>Elacatinus saucrum</i>	Leopard Goby	6	298
<i>Haemulon flavolineatum</i>	French Grunt	10	404
<i>Haemulon macrostomum</i>	Spanish Grunt	1	40
<i>Halichoeres poeyi</i>	Blackear Wrasse	1	26
<i>Lachnolaimus maximus</i>	Hogfish	1	10
<i>Mulloidichthys martinicus</i>	Yellow Goatfish	2	81
<i>Ocyurus chrysurus</i>	Yellowtail Snapper	7	212
<i>Scarus iseri</i>	Striped Parrotfish	1	33
<i>Stegastes adustus</i>	Dusky Damselfish	8	496
<i>Stegastes diencaeus</i>	Longfin Damselfish	4	127
<i>Stegastes partitus</i>	Bicolor Damselfish	4	2
<i>Stegastes variabilis</i>	Cocoa Damselfish	5	49

BCG Station 3
(Field Station 113_2011)
Coral Species Richness: 10
Photo rank and rating: 1, Good (Best)

Table I-8. BCG Station 3 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	3.16	0.92	0	0.48	1.36
Ave. 3D colony skeletal area (cm ²)/colony	4,537	13,445	0	949	704
Ave. 3D colony skeletal area (cm ²)/m ²	14,337	12,369	0	456	957
Ave. 3D coral tissue area (cm ²)/colony	3,012	8,646	0	626	609
Ave. 3D coral tissue area (cm ²)/m ²	9,518	7,954	0	300	829
Ave. 2D coral tissue area (cm ²)/m ²	2,843	2,212	0	135	326




Figure I-6. Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 3.

Table I-9. Fish species found in BCG Station 3, with density and biomass for 100 m² transect.

Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Acanthurus bahianus</i>	Ocean Surgeonfish	1	46
<i>Acanthurus coeruleus</i>	Blue Tang	9	638
<i>Canthigaster rostrata</i>	Sharpnose Puffer	2	14
<i>Caranx ruber</i>	Bar Jack	2	34
<i>Chaetodon capistratus</i>	Foureyeye Butterflyfish	1	15
<i>Coryphopterus glaucofraenum</i>	Bridled Goby	15	10
<i>Haemulon macrostomum</i>	Spanish Grunt	1	242
<i>Halichoeres bivittatus</i>	Slippery Dick	5	574
<i>Hypoplectrus chlorurus</i>	Yellowtail Hamlet	3	43
<i>Lutjanus apodus</i>	Schoolmaster	2	431
<i>Microspathodon chrysurus</i>	Yellowtail Damselfish	2	324
<i>Scarus iseri</i>	Striped Parrotfish	51	5,749
<i>Serranus tigrinus</i>	Harlequin Bass	1	7
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	5	482
<i>Sparisoma viride</i>	Stoplight Parrotfish	6	979
<i>Stegastes adustus</i>	Dusky Damselfish	30	360
<i>Stegastes partitus</i>	Bicolor Damselfish	4	32
<i>Stegastes planifrons</i>	Threespot Damselfish	2	13
<i>Thalassoma bifasciatum</i>	Bluehead	21	106

BCG Station 4
(Field Station 3_2011)
Coral Species Richness: 6
Photo rank and rating: 11, Poor

Table I-10. BCG Station 4 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	0.84	0.12	0	0	0.04
Ave. 3D colony skeletal area (cm ²)/colony	4,251	17,387	0	0	157
Ave. 3D colony skeletal area (cm ²)/m ²	3,571	2,086	0	0	6
Ave. 3D coral tissue area (cm ²)/colony	3,221	10,681	0	0	141
Ave. 3D coral tissue area (cm ²)/m ²	2,706	1,282	0	0	6
Ave. 2D coral tissue area (cm ²)/m ²	565	315	0	0	4




Figure I-7. Percentages of stony coral species and sponge morphologies for BCG Station 4.
 No gorgonians were present at BCG Station 4.

Table I-11. Fish species found in BCG Station 4, with density and biomass for 100 m² transect.

Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Acanthurus bahianus</i>	Ocean Surgeonfish	1	129
<i>Acanthurus chirurgus</i>	Doctorfish	6	629
<i>Acanthurus coeruleus</i>	Blue Tang	5	1,081
<i>Aulostomus maculatus</i>	Trumpetfish	1	256
<i>Canthigaster rostrata</i>	Sharpnose Puffer	1	7
<i>Chaetodon capistratus</i>	Foureye Butterflyfish	1	15
<i>Decapterus macarellus</i>	Mackerel Scad	2	395
<i>Gymnothorax sp.</i>	Moray Eel sp.	1	3
<i>Haemulon carbonarium</i>	Caesar Grunt	5	1,024
<i>Haemulon chrysargyreum</i>	Smallmouth Grunt	5	475
<i>Haemulon flavolineatum</i>	French Grunt	5	609
<i>Haemulon parra</i>	Sailors Choice	1	404
<i>Halichoeres radiatus</i>	Puddingwife	1	168
<i>Lutjanus apodus</i>	Schoolmaster	9	2,193
<i>Malacanthus plumieri</i>	Sand Tilefish	13	318
<i>Microspathodon chrysurus</i>	Yellowtail Damselfish	8	1,296
<i>Odontoscion dentex</i>	Reef Croaker	1	21
<i>Rypticus saponaceus</i>	Greater Soapfish	1	82
<i>Scarus iseri</i>	Striped Parrotfish	10	1,119
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	3	580
<i>Sparisoma rubripinne</i>	Yellowtail Parrotfish	1	401
<i>Sparisoma viride</i>	Stoplight Parrotfish	1	314
<i>Stegastes adustus</i>	Dusky Damselfish	33	384
<i>Synodus intermedius</i>	Sand Diver	1	339
<i>Thalassoma bifasciatum</i>	Bluehead	3	31

BCG Station 5
(Field Station 19_2011)
Coral Species Richness: 7
Photo rank and rating: 9, Fair/Poor

Table I-12. BCG Station 5 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	1.24	0.2	0.8	0.08	0
Ave. 3D colony skeletal area (cm ²)/colony	23,116	5,030	34,248	520	0
Ave. 3D colony skeletal area (cm ²)/m ²	28,664	1,006	27,399	42	0
Ave. 3D coral tissue area (cm ²)/colony	14,700	2,950	21,764	520	0
Ave. 3D coral tissue area (cm ²)/m ²	18,228	590	17,411	42	0
Ave. 2D coral tissue area (cm ²)/m ²	3,673	71	3,526	16	0




Figure I-8. Percentages of stony coral species and sponge morphologies for BCG Station 5.
 No gorgonians were present at BCG Station 5.

Table I-13. Fish species found in BCG Station 5, with density and biomass for 100 m² transect.

Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Acanthurus bahianus</i>	Ocean Surgeonfish	1	46
<i>Acanthurus coeruleus</i>	Blue Tang	2	72
<i>Anisotremus virginicus</i>	Porkfish	1	128
<i>Aulostomus maculatus</i>	Trumpetfish	3	439
<i>Bodianus rufus</i>	Spanish Hogfish	1	32
<i>Caranx ruber</i>	Bar Jack	1	17
<i>Decapterus macarellus</i>	Mackerel Scad	2	275
<i>Haemulon carbonarium</i>	Caesar Grunt	1	41
<i>Haemulon chrysargyreum</i>	Smallmouth Grunt	6	570
<i>Haemulon flavolineatum</i>	French Grunt	63	6,989
<i>Haemulon macrostomum</i>	Spanish Grunt	1	242
<i>Heteropriacanthus cruentatus</i>	Glasseye Snapper	1	214
<i>Holocentrus adscensionis</i>	Squirrelfish	1	237
<i>Lutjanus apodus</i>	Schoolmaster	2	203
<i>Malacanthus plumieri</i>	Sand Tilefish	13	300
<i>Microspathodon chrysurus</i>	Yellowtail Damselfish	3	486
<i>Mulloidichthys martinicus</i>	Yellow Goatfish	7	283
<i>Myripristis jacobus</i>	Blackbar Soldierfish	1	107
<i>Pempheris schomburgkii</i>	Glassy Sweeper	5	164
<i>Scarus iseri</i>	Striped Parrotfish	3	458
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	2	151
<i>Sparisoma viride</i>	Stoplight Parrotfish	3	521
<i>Stegastes adustus</i>	Dusky Damselfish	14	168
<i>Stegastes partitus</i>	Bicolor Damselfish	1	1
<i>Thalassoma bifasciatum</i>	Bluehead	16	134

BCG Station 6
(Field Station 14_2010)
Coral Species Richness: 10
Photo rank and rating: 10, Poor

Table I-14. BCG Station 6 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	3.6	1.2	0	0.33	1.4
Ave. 3D colony skeletal area (cm ²)/colony	5,431	14,255	0	1,400	466
Ave. 3D colony skeletal area (cm ²)/m ²	19,552	17,106	0	467	653
Ave. 3D coral tissue area (cm ²)/colony	1,489	3,298	0	1,193	443
Ave. 3D coral tissue area (cm ²)/m ²	5,359	3,958	0	398	621
Ave. 2D coral tissue area (cm ²)/m ²	1,530	892	0	141	330




Figure I-9. Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 6.

Table I-15. Fish species found in BCG Station 6, with density and biomass for 100 m² transect.


Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Acanthurus bahianus</i>	Ocean Surgeonfish	1	129
<i>Acanthurus chirurgus</i>	Doctorfish	1	121
<i>Chaetodon capistratus</i>	Foureye Butterflyfish	4	60
<i>Elacatinus genie</i>	Cleaning Goby	1	0
<i>Epinephelus adscensionis</i>	Rock Hind	1	174
<i>Halichoeres garnoti</i>	Yellowhead Wrasse	2	195
<i>Hypoplectrus chlorurus</i>	Yellowtail Hamlet	1	19
<i>Hypoplectrus puella</i>	Barred Hamlet	1	19
<i>Lutjanus apodus</i>	Schoolmaster	1	649
<i>Mulloidichthys martinicus</i>	Yellow Goatfish	2	472
<i>Ocyurus chrysurus</i>	Yellowtail Snapper	1	177
<i>Scarus iseri</i>	Striped Parrotfish	36	632
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	5	466
<i>Sparisoma viride</i>	Stoplight Parrotfish	11	1,606
<i>Stegastes diencaeus</i>	Longfin Damselfish	3	36
<i>Stegastes leucostictus</i>	Beaugregory	4	31
<i>Stegastes partitus</i>	Bicolor Damselfish	7	24
<i>Stegastes planifrons</i>	Threespot Damselfish	12	229
<i>Thalassoma bifasciatum</i>	Bluehead	12	55

BCG Station 7
(Field Station 16_2010)
Coral Species Richness: 11
Photo rank and rating: 4, Fair

Table I-16. BCG Station 7 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	4.87	1.13	0	0.53	1.53
Ave. 3D colony skeletal area (cm ²)/colony	3,529	10,160	0	895	567
Ave. 3D colony skeletal area (cm ²)/m ²	17,173	11,515	0	477	870
Ave. 3D coral tissue area (cm ²)/colony	2,920	9,057	0	790	469
Ave. 3D coral tissue area (cm ²)/m ²	14,210	10,264	0	421	720
Ave. 2D coral tissue area (cm ²)/m ²	5,089	3,537	0	167	303

Coral Species



Gorgonian Morphs




Figure I-10. Percentages of stony coral species and gorgonian morphologies for BCG Station 7.
 No sponges were present at BCG Station 7.

Table I-17. Fish species found in BCG Station 7, with density and biomass for 100 m² transect.


Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Acanthurus coeruleus</i>	Blue Tang	9	414
<i>Chaetodon striatus</i>	Banded Butterflyfish	2	124
<i>Chromis multilineata</i>	Brown Chromis	16	126
<i>Haemulon aurolineatum</i>	Tomtate	3	21
<i>Haemulon flavolineatum</i>	French Grunt	2	81
<i>Holocentrus adscensionis</i>	Squirrelfish	2	474
<i>Lutjanus apodus</i>	Schoolmaster	1	215
<i>Microspathodon chrysurus</i>	Yellowtail Damselfish	13	1,133
<i>Ocyurus chrysurus</i>	Yellowtail Snapper	10	303
<i>Pomacanthus paru</i>	French Angelfish	1	342
<i>Scarus iseri</i>	Striped Parrotfish	16	1,515
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	17	475
<i>Stegastes diencaeus</i>	Longfin Damselfish	8	294
<i>Stegastes leucostictus</i>	Beaugregory	2	89
<i>Stegastes partitus</i>	Bicolor Damselfish	3	31
<i>Thalassoma bifasciatum</i>	Bluehead	36	110

BCG Station 8
(Field Station 108_2010)
Coral Species Richness: 7
Photo rank and rating: 6, Fair

Table I-18. BCG Station 8 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	4.73	0.20	1.47	0.13	1.93
Ave. 3D colony skeletal area (cm ²)/colony	4,938	1,473	14,658	157	258
Ave. 3D colony skeletal area (cm ²)/m ²	23,372	295	21,499	21	499
Ave. 3D coral tissue area (cm ²)/colony	4,149	949	12,274	157	239
Ave. 3D coral tissue area (cm ²)/m ²	19,637	190	18,002	21	462
Ave. 2D coral tissue area (cm ²)/m ²	4,549	52	3,964	10	273

Coral Species



Gorgonian Morphs




Figure I-11. Percentages of stony coral species and gorgonian morphologies for BCG Station 8.
 No sponges were present at BCG Station 8.

Table I-19. Fish species found in BCG Station 8, with density and biomass for 100 m² transect.

Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Abudefduf saxatilis</i>	Sergeant Major	3	99
<i>Acanthurus bahianus</i>	Ocean Surgeonfish	3	386
<i>Acanthurus coeruleus</i>	Blue Tang	31	1,479
<i>Aulostomus maculatus</i>	Trumpetfish	1	130
<i>Haemulon flavolineatum</i>	French Grunt	1	111
<i>Halichoeres bivittatus</i>	Slippery Dick	4	182
<i>Halichoeres maculipinna</i>	Clown Wrasse	6	24
<i>Halichoeres radiatus</i>	Puddingwife	1	0
<i>Lutjanus apodus</i>	Schoolmaster	4	341
<i>Microspathodon chrysurus</i>	Yellowtail Damselfish	28	961
<i>Ophioblennius macclurei</i>	Redlip Blenny	1	4
<i>Pomacanthus paru</i>	French Angelfish	1	156
<i>Scarus iseri</i>	Striped Parrotfish	69	4,592
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	4	332
<i>Sparisoma viride</i>	Stoplight Parrotfish	2	326
<i>Stegastes diencaeus</i>	Longfin Damselfish	54	603
<i>Stegastes partitus</i>	Bicolor Damselfish	6	3
<i>Thalassoma bifasciatum</i>	Bluehead	98	331

BCG Station 9
(Field Station 109_2010)
Coral Species Richness: 9
Photo rank and rating: 7, Fair

Table I-20. BCG Station 9 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	5.80	3.47	0	0.53	0.67
Ave. 3D colony skeletal area (cm ²)/colony	9,091	8,102	0	806	781
Ave. 3D colony skeletal area (cm ²)/m ²	52,731	28,088	0	430	521
Ave. 3D coral tissue area (cm ²)/colony	3,462	4,040	0	734	739
Ave. 3D coral tissue area (cm ²)/m ²	20,080	14,006	0	391	493
Ave. 2D coral tissue area (cm ²)/m ²	5,917	4,015	0	213	298




Figure I-12. Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 9.

Table I-21. Fish species found in BCG Station 9, with density and biomass for 100 m² transect.

Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Abudefduf saxatilis</i>	Sergeant Major	4	263
<i>Acanthurus bahianus</i>	Ocean Surgeonfish	1	129
<i>Acanthurus chirurgus</i>	Doctorfish	10	1,206
<i>Acanthurus coeruleus</i>	Blue Tang	76	11,429
<i>Anisotremus virginicus</i>	Porkfish	3	851
<i>Aulostomus maculatus</i>	Trumpetfish	1	442
<i>Cephalopholis cruentata</i>	Graysby	1	178
<i>Chaetodon striatus</i>	Banded Butterflyfish	2	25
<i>Coryphopterus glaucofraenum</i>	Bridled Goby	10	7
<i>Haemulon carbonarium</i>	Caesar Grunt	3	915
<i>Haemulon plumieri</i>	White Grunt	1	539
<i>Halichoeres bivittatus</i>	Slippery Dick	2	342
<i>Holocentrus adscensionis</i>	Squirrelfish	1	433
<i>Hypoplectrus chlorurus</i>	Yellowtail Hamlet	3	12
<i>Lutjanus apodus</i>	Schoolmaster	2	431
<i>Microspathodon chrysurus</i>	Yellowtail Damselfish	7	402
<i>Mulloidichthys martinicus</i>	Yellow Goatfish	1	236
<i>Scarus iseri</i>	Striped Parrotfish	19	98
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	4	1,008
<i>Sparisoma viride</i>	Stoplight Parrotfish	16	5,024
<i>Stegastes diencaeus</i>	Longfin Damselfish	30	348
<i>Stegastes leucostictus</i>	Beaugregory	2	20
<i>Stegastes partitus</i>	Bicolor Damselfish	18	79
<i>Stegastes planifrons</i>	Threespot Damselfish	2	24
<i>Thalassoma bifasciatum</i>	Bluehead	4	14

BCG Station 10
(Field Station 2011_1)
Coral Species Richness: 11
Photo rank and rating: 3, Fair

Table I-22. BCG Station 10 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	2.8	1.88	0.04	0.08	0.4
Ave. 3D colony skeletal area (cm ²)/colony	7,205	10,095	9,503	3,662	794
Ave. 3D colony skeletal area (cm ²)/m ²	20,174	18,979	380	293	317
Ave. 3D coral tissue area (cm ²)/colony	3,938	5,549	0	1,405	640
Ave. 3D coral tissue area (cm ²)/m ²	11,026	10,432	0	112	256
Ave. 2D coral tissue area (cm ²)/m ²	3,698	3,496	0	50	94




Figure I-13. Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 10.

Table I-23. Fish species found in BCG Station 10, with density and biomass for 100 m² transect.

Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Chaetodon capistratus</i>	Foureye Butterflyfish	2	30
<i>Decapterus macarellus</i>	Mackerel Scad	4	250
<i>Haemulon flavolineatum</i>	French Grunt	22	2,441
<i>Hypoplectrus chlorurus</i>	Yellowtail Hamlet	2	24
<i>Malacanthus plumieri</i>	Sand Tilefish	1	24
<i>Malacoctenus triangulatus</i>	Saddled Blenny	4	1
<i>Odontoscion dentex</i>	Reef Croaker	14	804
<i>Pomacanthus paru</i>	French Angelfish	2	2,161
<i>Scarus iseri</i>	Striped Parrotfish	6	45
<i>Sparisoma viride</i>	Stoplight Parrotfish	15	789
<i>Stegastes partitus</i>	Bicolor Damselfish	3	2
<i>Stegastes planifrons</i>	Threespot Damselfish	96	1,982

BCG Station 11
(Field Station 46_2011)
Coral Species Richness: 9
Photo rank and rating: 5, Fair

Table I-24. BCG Station 11 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	1.76	0.16	0.60	0.20	0.60
Ave. 3D colony skeletal area (cm ²)/colony	8,236	639	21,258	1,210	422
Ave. 3D colony skeletal area (cm ²)/m ²	14,495	102	12,755	242	253
Ave. 3D coral tissue area (cm ²)/colony	6,610	495	16,902	491	363
Ave. 3D coral tissue area (cm ²)/m ²	11,635	79	10,141	98	218
Ave. 2D coral tissue area (cm ²)/m ²	2,598	41	2,187	44	96




Figure I-14. Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 11.

Table I-25. Fish species found in BCG Station 11, with density and biomass for 100 m² transect.

Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Acanthurus bahianus</i>	Ocean Surgeonfish	21	959
<i>Acanthurus coeruleus</i>	Blue Tang	7	831
<i>Aulostomus maculatus</i>	Trumpetfish	1	15
<i>Caranx ruber</i>	Bar Jack	20	134
<i>Cephalopholis cruentata</i>	Graysby	1	82
<i>Chaetodon capistratus</i>	Foureyeye Butterflyfish	10	150
<i>Chaetodon striatus</i>	Banded Butterflyfish	4	247
<i>Haemulon carbonarium</i>	Caesar Grunt	15	1,543
<i>Haemulon chrysargyreum</i>	Smallmouth Grunt	30	1,011
<i>Haemulon flavolineatum</i>	French Grunt	15	606
<i>Halichoeres garnoti</i>	Yellowhead Wrasse	1	82
<i>Halichoeres maculipinna</i>	Clown Wrasse	1	5
<i>Halichoeres radiatus</i>	Puddingwife	4	68
<i>Holocentrus rufus</i>	Longspine Squirrelfish	2	190
<i>Lutjanus apodus</i>	Schoolmaster	5	185
<i>Malacanthus plumieri</i>	Sand Tilefish	15	367
<i>Microspathodon chrysurus</i>	Yellowtail Damselfish	14	1,204
<i>Myripristis jacobus</i>	Blackbar Soldierfish	1	107
<i>Scarus iseri</i>	Striped Parrotfish	46	386
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	2	119
<i>Stegastes adustus</i>	Dusky Damselfish	11	132
<i>Stegastes leucostictus</i>	Beaugregory	1	1
<i>Stegastes partitus</i>	Bicolor Damselfish	10	45
<i>Thalassoma bifasciatum</i>	Bluehead	170	698

BCG Station 12
(Field Station 25_2011)
Coral Species Richness: 8
Photo rank and rating: 2, Fair

Table I-26. BCG Station 12 data summary for corals and subgroups. See Appendix J for formulas used and species abbreviations.

Coral Parameters	All Species	Massive Species	Acroporid Species	<i>Siderastrea siderea</i>	<i>Porites astreoides</i>
Colony density (#/m ²)	3.8	1.56	0	0.36	1.72
Ave. 3D colony skeletal area (cm ²)/colony	3,525	7,563	0	592	740
Ave. 3D colony skeletal area (cm ²)/m ²	13,396	11,799	0	213	1,273
Ave. 3D coral tissue area (cm ²)/colony	2,421	5,037	0	460	619
Ave. 3D coral tissue area (cm ²)/m ²	9,199	7,857	0	166	1,065
Ave. 2D coral tissue area (cm ²)/m ²	3,498	2,928	0	61	472




Figure I-15. Percentages of stony coral species, gorgonian morphologies and sponge morphologies for BCG Station 12.

Table I-27. Fish species found in BCG Station 12, with density and biomass for 100 m² transect.

Fish Species	Common Name	Abundance/ 100 m²	Total Biomass (g/100 m²)
<i>Acanthurus bahianus</i>	Ocean Surgeonfish	2	91
<i>Acanthurus chirurgus</i>	Doctorfish	1	44
<i>Haemulon flavolineatum</i>	French Grunt	3	262
<i>Labrisomus nuchipinnis</i>	Hairy Blenny	1	33
<i>Lutjanus apodus</i>	Schoolmaster	1	101
<i>Myripristis jacobus</i>	Blackbar Soldierfish	1	107
<i>Scarus iseri</i>	Striped Parrotfish	8	85
<i>Sparisoma aurofrenatum</i>	Redband Parrotfish	6	247
<i>Sparisoma viride</i>	Stoplight Parrotfish	4	534
<i>Stegastes adustus</i>	Dusky Damselfish	1	12

Appendix J. Formulas Used for Calculating Condition Metrics

1. Stony Corals

Coral metrics

- Colony surface area
- Coral abundance
- Percent live tissue
- Live colony three-dimensional surface area
- Live colony two-dimensional surface area

Colony condition measurements

Every scleractinian coral within a 25 m² transect area (25 m x 1 m) and greater than 10 cm was identified to species. The maximum height and diameter of each colony was measured in cms, and the percent of living coral tissue on the skeleton of the colony was estimated in 10% increments. The percent tissue (living coral) was estimated for the entire colony in three dimensions, not only from an aerial planar view. The observations and measurements made for each coral colony included: scleractinian taxon, height (cm), maximum diameter (cm) and percent living colony tissue.

Formulas

Colony surface area (CSA) was the total three-dimensional colony surface area (cm²) including both living and dead portions of a single coral colony.

$$CSA = \pi r^2 M$$

$$r = [\text{colony height (cm)} + (\text{colony diameter (cm)}/2)] / 2$$

M = morphological conversion factor (values of 1, 2, 3, or 4 depending on coral species morphology), see Table J-1

Coral abundance (n) was total number of colonies in the entire transect area

% Live tissue (LT) was estimation of percent live tissue on a single coral colony over the entire surface area. It was estimated for every coral colony in transect.

Live colony 3D surface area (LCSA_3D) was a calculated value for the total three-dimensional colony surface area (cm²) of only living tissue on a single coral colony.

$$LCSA_3D = CSA * (LT/100)$$

Live colony 2D surface area (LCSA_2D) was a calculated value for the total planar colony surface area (cm²) of living tissue on a single coral colony as though it were viewed from above. This calculation assumes equal distribution of living tissue on a colony, which was initially recorded for three dimensions rather than two. It approximates percent coral cover used as the standard in many historical assessments.

$$LCSA_2D = \pi [\text{colony diameter (cm)}/2]^2 * (LT/100)$$

Coral metrics calculated for each BCG station

Ave. 3D colony skeletal area (cm²)/colony = $\Sigma \text{CSA}/n$

Ave. 3D colony skeletal area (cm²)/m² = $\Sigma \text{CSA}/\text{area of transect}$

Ave. 3D coral tissue (live) area (cm²)/colony = $\Sigma \text{LCSA_3D}/n$

Ave. 3D coral tissue (live) area (cm²)/m² = $\Sigma \text{LCSA_3D}/\text{area of transect}$

Ave. 2D coral tissue (live) area (cm²)/m² = $\Sigma \text{LCSA_2D}/\text{area of transect}$

Table J-1. Stony corals included in Western Atlantic and Caribbean assessments (as provided by Humann and DeLoach 2002) with the three-letter identification code and the morphological conversion factor for calculating 3D surface area (Santavy et al. 2012).

Genus and Species	ID Code	Conversion Factor
<i>Acropora cervicornis</i>	Acer	4
<i>Acropora palmata</i>	Apal	4
<i>Acropora prolifera</i>	Apro	4
<i>Agaricia agaricites</i>	Aaga	1
<i>Agaricia fragilis</i>	Afra	1
<i>Agaricia humilis</i>	Ahum	1
<i>Agaricia lamarcki</i>	Alam	1
<i>Agaricia tenuifolia</i>	Aten	3
<i>Cladocora arbuscula</i>	Carb	2
<i>Colpophyllia natans</i>	Cnat	2
<i>Dendrogyra cylindrus</i>	Dcyl	3
<i>Dichocoenia stokesii</i>	Dsto	2
<i>Diploria clivosa</i> ¹	Dcli	2
<i>Diploria labyrinthiformis</i>	Dlab	2
<i>Diploria strigosa</i> ¹	Dstr	2
<i>Eusmilia fastigiata</i>	Efas	3
<i>Favia fragum</i>	Ffra	2
<i>Leptoseris cucullata</i>	Lcuc	1
<i>Isophyllastrea rigida</i>	Irig	2
<i>Isophyllia sinuosa</i>	Isin	2
<i>Madracis decactis</i>	Mdec	3
<i>Madracis formosa</i>	Mfor	3
<i>Madracis mirabilis</i>	Mmir	3
<i>Madracis pharensis</i>	Mpha	1
<i>Manicina areolata</i>	Mare	2
<i>Meandrina meandrites</i>	Mmea	2
<i>Millepora complanata</i>	Mcom	3
<i>Montastraea annularis</i> ²	Mann	3
<i>Montastraea cavernosa</i>	Mcav	2
<i>Montastraea faveolata</i> ²	Mfav	2

Table J-1. (continued)

Genus and Species	ID Code	Conversion Factor
<i>Mussa angulosa</i>	Mang	2
<i>Mycetophyllia aliciae</i>	Mali	1
<i>Mycetophyllia danaana</i>	Mdan	1
<i>Mycetophyllia ferox</i>	Mfer	1
<i>Mycetophyllia lamarckiana</i>	Mlam	1
<i>Oculina varicosa</i>	Ovar	3
<i>Porites astreoides</i>	Past	2
<i>Porites colonensis</i>	Pcol	1
<i>Porites divaricata</i>	Pdiv	3
<i>Porites furcata</i>	Pfur	3
<i>Porites porites</i>	Ppor	3
<i>Siderastrea siderea</i>	Ssid	2
<i>Solenastrea bournoni</i>	Sbou	2
<i>Solenastrea hyades</i>	Shya	3
<i>Stephanocoenia intersepta</i>	Sint	2

- 1: This report does not adopt the new classifications for *Diploria strigosa* and *Diploria clivosa* as the original genus *Pseudodiploria* (Budd et al. 2012).
- 2: This report does not adopt the new classification for the *Montastraea annularis* species-complex (*Montastraea annularis*, *Montastraea faveolata* and *Montastraea franksi*) which has been reclassified as the original genus *Orbicella* (Budd et al. 2012).

2. Rugosity

Rugosity measurement and metric

Rugosity was the linear ratio of a 6 m chain length compared to the taunt linear distance in centimeters of a draped chain. Rugosity is a reef-scale indicator of reef contour. It was determined using a chain-transect method that compares the length of a chain draped along the top of corals and along the bottom of the reef to the length of a taut line across the same linear distance using a separate tape measure, laid parallel but not on top of the transect tape. The linked chain was placed such that it follows the relief of hard bottom substrate. The chain was placed on top of any hard substrate encountered, but not on top of gorgonians or sponges since only hard bottom rugosity was being measured.

Formula

Rugosity was the ratio of the overall length of chain draped over the reef contour divided by the straight horizontal distance between the beginning and the end of the chain. Therefore, if 6 m of chain is laid out over a 4 m horizontal distance, the rugosity is $6/4 = 1.5$ for that segment. Rugosity will always be ≥ 1 . Higher values relate to increased rugosity or reef relief. The average rugosity was calculated per transect.

3. Fish

Fish metrics

Abundance at lowest taxonomic level possible

Length in cms

Biomass of fish in g/100 m²

Fish measurements

Fish contained within a 100 m² transect area, 25 m length x 4 m width (height was water depth) were recorded to the lowest taxonomic level possible. All fish greater than 3 cm in size were included in the assessment. Each fish was scored as 5 cm size class increments up to 35 cm for fork length using visual estimation. If a fish was longer than 35 cm, an estimate of the actual fork length was made. The fork length was measured from the fish snout (with mouth closed) to the fork at the base of the tail or caudal fin. Observations and measurements made for each fish included taxon and size class.

Formulas

Abundance (n) was total number of fish in the entire transect area

Density was the number of fish of a single taxon per 100 m²

Biomass (W) was the weight recorded as g/100 m² of a single fish

$$W = \alpha L^{\beta}$$

L = fork length as midpoint between 5 cm increment class (e.g., 10–15 cm class using 12.5 as length in calculation). L > 35 cm uses actual length.

α and β are species specific coefficients obtained from FishBase (www.fishbase.org) for calculating fish biomass (see Appendix A in Santavy et al. 2013). Biomass for species with no published length-weight relationships can be calculated using terms for the closest congener based on morphology.

Fish metrics calculated for each BCG station

$$\text{Total biomass for each taxon per 100 m}^2 = \sum_{i=1}^n W$$

4. Gorgonians

Gorgonian metrics

Density was the number of individuals/m²

Average three-dimensional surface area gorgonian/m²

Average three-dimensional surface area gorgonian/individual

Gorgonian measurements

Every gorgonian ≥ 10 cm (in any dimension) that falls within the quadrat was classified as one of ten gorgonian morphologies (Table J-2). If the base of the gorgonian was in the quadrat, it was considered in the transect area. Colony height (greatest distance from substrate) and maximum diameter (parallel to the substrate) were measured in cms. The observations and measurements made for each individual were gorgonian colony shape, height and maximum diameter. Although gorgonians are prominent reef inhabitants, they are often excluded from monitoring programs. This is partially because they are not widely recognized for their important functional contributions to reef environments, and partially because taxonomic distinctions can be difficult. In this approach, classification was based on morphology, categorized by predetermined shapes, which can be easier to apply than taxonomy and still can influence their ecosystem functions.

Formulas and BCG station metrics

Abundance (n) was total number of gorgonians in the entire transect area


Average three-dimensional surface area of each gorgonian morph/m² =

$$\sum_{i=1}^n \frac{(\text{regression equation for SA in Table J - 2})}{\text{area of transect}}$$

Average three-dimensional surface area of each gorgonian morph/colony =

$$\sum_{i=1}^n \frac{(\text{regression equation for SA in Table J - 2})}{n}$$

Table J-2. Gorgonian morphological shapes, abbreviations, simulated model, *in situ* example and regression models to estimate three-dimensional surface area (Santavy et al. 2013).

Gorgonian Morphology	Species Example	Simulated Model	<i>in situ</i> Example	Surface Area Estimations
Sea Fans	Planar (fan pl) (<i>Gorgonia ventalina</i> , <i>Leptogorgia</i>)			$SA=0.68h^2+0.66d^2-3.61$
	Three-dimensional (3D fan) (<i>Gorgonia flabellum</i>)			$SA=0.0113h^3+106d-1190$
Sea Rods branch and branchlet diameter ≥ 15 – ≤ 30 mm	Unbranched (SR ub) digitate form (<i>Briareum</i>)			$SA=0.341d^3+11.2h-127$
	Branched (SR br) (<i>Plexaura</i>)			$SA=1.46d^2+399$
	Bushy (SR bush) (<i>Eunicea fusca</i>)			$SA=0.0288h^3+939$
	Planar (SR pl) (<i>Eunicea tourneforti</i>)			$SA=76.4d-806$
Sea Whips branch and branchlet diameter ≥ 5 – ≤ 15 mm	Branched (SW br) (<i>Pterogorgia</i>)			$SA=-0.479h^3+3.37h^2-51.3h+354$
	Bushy (SW bush) (<i>Pterogorgia guadalupensi</i>)			$SA=0.0672d^3+1610$
Sea Plumes smallest branch and branchlet diameter usually ≤ 5 mm	(Plume) (<i>Muriceopsis flavida</i> , <i>Pseudopterogorgia</i>)			$SA=4.77h^2-2990$
Encrusting Gorgonians	(<i>Briareum</i> , <i>Erythropodium</i>)			$SA=dw$

Minimum height (h) and diameter (d) of colony size required for use in the equations. h is the maximum colony height measured in cm, d is the maximum colony diameter measured in cm.

5. Sponges

Sponge metrics

Density was the number of individuals/m²

Average three-dimensional surface area of sponges/m²

Average three-dimensional surface area of sponges/individual

Sponge measurements

Every sponge ≥ 10 cm (in any dimension) falling within the quadrat was classified as one of ten sponge morphologies (Table J-3). If the base of sponge was in the quadrat, it was considered in the transect area. Colony height (greatest distance from substrate) and maximum diameter (parallel to the substrate) were recorded in cms. The observations and measurements made for each sponge were colony shape, height and maximum diameter. Although sponges are one of the most prominent sessile invertebrates on coral reefs, they are often overlooked in monitoring programs. This may be in part because sponge taxonomic classification is confounded by high diversity and morphological plasticity. In this approach, classification was based on morphology rather than taxonomy and can influence their ecosystem functions.

Formulas and BCG station metrics

Abundance (n) was total number of sponges in the entire transect area




Average three-dimensional surface area of each sponge morph/m² =

$$\sum_{i=1}^n \frac{(\text{regression equation for SA in Table J-3})}{\text{area of transect}}$$

Average three-dimensional surface area of each sponge morph per individual =

$$\sum_{i=1}^n \frac{(\text{regression equation for SA in Table J-3})}{n}$$

Table J-3. Sponge morphological shapes, abbreviations, simulated model, *in situ* example and regression models to estimate surface area (Santavy et al. 2013).

Sponge Morphology	Species Example	Simulated Model	<i>in situ</i> Example	Surface Area Estimations
Barrel	<i>Xestospongia muta</i> , <i>Verongula reiswigi</i>			$SA=4.31d^2 + 0.827h^2 + 108$
Vase	<i>Callyspongia plicifera</i> , <i>Callyspongia vaginalis</i>			$SA=3.71h^2-161$
Globe	<i>Iricinia strobilina</i> , <i>Sphaciospongia vesparium</i>			$SA=1.88h^2 + 0.0573d^3 + 83.3$
Tube	<i>Aplysina archeri</i> , <i>Aplysina fistularis</i>			$SA=0.493d^3+109$
Mound	<i>Oligoceras hemorrhages</i> , <i>Iricinia felix</i>			$SA=30.0h+18.7d-193$
Rod	<i>Aplysina cauliformis</i> , <i>Niphates erecta</i>			$SA=7.69h+1.83d^3-33.5$
Bushy	<i>Aplysina fulva</i>			$SA=0.462h^2+0.834d^2+19.3$
Branched Ropey	<i>Iotrochota birotulata</i>			$SA=18.8d+7.97h-132$
Encrusting	<i>Amphimedon compressa</i> , <i>Chondrilla caribensis</i>			$SA=dw$

Minimum height (h) and diameter (d) of colony size required for use in the equations. h is the maximum colony height measured in cm, d is the maximum colony diameter measured in cm.



United States
Environmental Protection
Agency

Office of Research and Development
National Health and Environmental
Effects Research Laboratory
Atlantic Ecology Division
Narragansett, RI 02882

Official Business
Penalty for Private use
\$300

EPA/600/R-13/350 | December 2014
www.epa.gov/ord