



Northeast Fisheries Science Center Reference Document 14-10

doi:10.7289/V5JW8BVD

Mortality Determinations for Baleen Whale Stocks along the Gulf of Mexico, United States East Coast, and Atlantic Canadian Provinces, 2008 - 2012

by Allison G. Henry, Timothy V.N. Cole, Lanni Hall, Wayne Ledwell,
David Morin, and Andrew Reid

Mortality Determinations for Baleen Whale Stocks along the Gulf of Mexico, United States East Coast, and Atlantic Canadian Provinces, 2008 - 2012

by Allison G. Henry¹, Timothy V.N. Cole¹, Lanni Hall², Wayne Ledwell³,
David Morin², and Andrew Reid⁴

¹NOAA Fisheries, Northeast Fisheries Science Center, 166 Water Street, Woods Hole, MA 02543

²NOAA Fisheries Service, Greater Atlantic Regional Fisheries Office,
55 Great Republic Drive, Gloucester, MA 01930-2276

³Whale Release and Strandings Group, 244 Tolt Rd., Portugal Cove-St. Philip's,
Newfoundland, CAN A1M 1R2

⁴Marine Animal Response Society, c/o Nova Scotia Museum, 1747 Summer St.,
Halifax, Nova Scotia CAN B3H 3A6

U.S. DEPARTMENT OF COMMERCE

National Oceanic and Atmospheric Administration

National Marine Fisheries Service

Northeast Fisheries Science Center

Woods Hole, Massachusetts

April 2014

Northeast Fisheries Science Center Reference Documents

This series is a secondary scientific series designed to assure the long-term documentation and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review, and most receive copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

All documents issued in this series since April 2001, and several documents issued prior to that date, have been copublished in both paper and electronic versions. To access the electronic version of a document in this series, go to <http://www.nefsc.noaa.gov/nefsc/publications/>. The electronic version is available in PDF format to permit printing of a paper copy directly from the Internet. If you do not have Internet access, or if a desired document is one of the pre-April 2001 documents available only in the paper version, you can obtain a paper copy by contacting the senior Center author of the desired document. Refer to the title page of the document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate (*i.e.*, non-individualized) authorship, then contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

Information Quality Act Compliance: In accordance with section 515 of Public Law 106-554, the Northeast Fisheries Science Center completed both technical and policy reviews for this report. These predissemination reviews are on file at the NEFSC Editorial Office.

This document may be cited as:

Henry AG, Cole TVN, Hall L, Ledwell W, Morin D, Reid A. 2014. Mortality determinations for baleen whale stocks along the Gulf of Mexico, United States east coast, and Atlantic Canadian provinces, 2008 - 2012. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 14-10; 17 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://www.nefsc.noaa.gov/nefsc/publications/>

TABLE OF CONTENTS

Abstract	2
Introduction	2
Methods	2
Results	5
Discussion.....	6
Acknowledgements	7
References Cited	8

LIST OF TABLES

Table 1. Summary of all unique large whale mortalities observed along the Gulf of Mexico Coast, US East Coast, and Atlantic Canadian Provinces, 2008-2012.....	9
Table 2. Confirmed human-caused mortality records of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012.....	10
Table 3. Summary of country of origin for all confirmed human-caused mortalities of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012.....	17

ABSTRACT

The Northeast Fisheries Science Center (NEFSC) developed criteria to evaluate reports of human-caused mortality to baleen whales. The criteria minimize the likelihood of incorrectly assigning whale mortalities to human causes and provide a minimum count of human-caused events. This report describes determinations made for reports received from 2008 - 2012 involving North Atlantic right (*Eubalaena glacialis*), humpback (*Megaptera novaeangliae*), fin (*Balaenoptera physalus*), sei (*B. borealis*), blue (*B. musculus*), minke (*B. acutorostrata*), and Bryde's (*B. edeni*) whales observed in United States waters in the Gulf of Mexico, along the US eastern seaboard and in the Atlantic Canadian provinces. We confirmed a total of 300 mortalities: 60 (20%) caused by human interaction, 14 (5%) because of natural causes, and 236 (75%) which lacked sufficient evidence to determine cause of death. Of the human-caused mortalities, 32 were due to entanglement and 28 were vessel strike. These mortality numbers are minimum counts because of a low probability of detecting events and inadequate documentation to determine cause of death for the majority of events that are detected. Despite the minimum values, the mean annual confirmed human-caused mortality rate exceeds the Potential Biological Removal (PBR) value for 4 of the 7 stocks examined, including North Atlantic right, humpback, sei, and Bryde's whales. The true level of anthropogenic mortality to these stocks is greater than these minimum values, but the amount greater is unknown.

INTRODUCTION

The NOAA National Marine Fisheries Service (NMFS) is required to estimate annual rates of human-caused mortality and serious injury to marine mammal stocks occurring regularly in US waters. The agency is also charged with developing plans to reduce the rate of human-caused mortality and serious injury to strategic stocks to levels below their Potential Biological Removal (PBR). The PBR is the maximum number of animals, not including natural mortalities, which may be removed from a marine mammal stock while allowing that stock to reach or maintain its optimum sustainable population size (Wade and Angliss 1997). A 5-year average rate of human-caused mortality and serious injury is reported for each species in annual marine mammal stock assessment reports (SAR; e.g., Waring et al. 2014). This rate, when compared to a population's PBR, is used to identify stocks for which management actions may be required under the Marine Mammal Protection Act (MMPA Sec. 118).

This report presents the method and results of cause-of-death determination criteria that establish the minimum annual rates of confirmed human-caused mortality for North Atlantic right (*Eubalaena glacialis*), humpback (*Megaptera novaeangliae*), fin (*Balaenoptera physalus*), sei (*B. borealis*), blue (*B. musculus*), minke (*B. acutorostrata*), and Bryde's (*B. edeni*) whale stocks along the Gulf of Mexico, the eastern seaboard of the United States, and the Atlantic Canadian provinces for the period 2008 - 2012. Serious injuries and their causes are reported elsewhere (Cole and Henry, in prep).

METHODS

Members of the US and Canadian regional stranding networks, large whale disentanglement teams, the US and Canadian Coast Guards, and the general public provided

opportunistic marine mammal stranding and human interaction reports to the NMFS Greater Atlantic Regional Fisheries Office (GARFO), Southeast Regional Office (SERO), or the NEFSC. With the exception of minke whales, the incidental takes of large whales recorded by fisheries observer programs were treated as opportunistic reports because of the low number of observed takes. The Regional Offices obtained all available information for each report (photos, necropsy reports, etc.), which was then reviewed by NEFSC and GARFO staff members. Confirmed reports were designated “events,” and for each event the species involved was verified, duplicate records identified, and relevant information from each source consolidated into a single record. Subsequent demographic, health, and sighting history information were obtained, where available, from local population monitoring studies. NEFSC staff reviewed each mortality event and assigned a cause of death following the confirmation criteria listed below. One staff member reviewed all determinations each year to ensure consistency in the application of determination criteria within and across years.

Events from Newfoundland and Labrador involving confirmed transboundary stocks (i.e., stocks that enter US waters during part of the year) were also included. However, humpback events from these regions were not included in tallies because the Newfoundland and Labrador humpback feeding stocks are distinct from the Gulf of Maine stock found in US waters (Palsbøll et al. 2001).

Confirmation Criteria for Identification of Cause of Mortality (listed in order of certainty)

The cause of mortality was considered confirmed if the report included 1 of the following criteria:

1. Photographs or video allowed identification;
2. A marine mammal expert reported the mortality cause as certain; or
3. A report made by a trained observer or volunteer member of a stranding network which was verified by NMFS or stranding network staff.

The cause of mortality was considered confirmed in the following less certain cases:

1. Photographs or video allowed probable identification;
2. A marine mammal expert reported the mortality cause as probable;
3. An inexperienced observer’s report allowed probable identification; or
4. An inexperienced observer’s report was verified by NMFS or stranding network staff.

The cause of mortality was considered unconfirmed if:

1. Photographs or video were of insufficient quality to verify;
2. An inexperienced observer’s report lacked photographs, video, or descriptive detail to confirm;
3. An incomplete examination did not allow for identification; or
4. A carcass was too decomposed to identify.

Human-induced Mortality Determinations

Events were categorized as entanglement mortalities if 1 of the following indications were confirmed to be present on a whale carcass:

1. Fishing line constricted any body part and subdermal hemorrhaging or extensive necrosis was present at point of attachment;
2. An extensive entanglement was evident;
3. An entanglement likely prevented feeding; or
4. A code 2 (fresh dead) whale was pulled up during fishing operations.

Events were categorized as vessel strike mortalities if 1 of the following indications was confirmed to be present on a whale carcass:

1. Large linear laceration(s) was present anywhere on body, as opposed to just dorsally as in Kraus (1990);
2. Large area(s) of subdermal hemorrhaging, hematoma, or edema was evident;
3. Major skeletal fracturing was evident; or
4. A code 2 (fresh dead) carcass was found on the bow of a ship.

Assignment to Country

We assigned entanglement mortalities to either the United States or Canada based on the entangling gear's country of origin. Identification of gear type and country of origin was conducted by GARFO, the Marine Animal Response Society based in Nova Scotia, or the Whale Release and Strandings Group based in Newfoundland. GARFO's results are included in annual Atlantic Large Whale Take Reduction Program (ALWTRP) reports¹. Identified fisheries are categorized in the List of Fisheries (LOF) according to their frequency of incidental mortality or serious injury to marine mammal stocks (50 CFR Parts 216 and 229²).

If gear identification was lacking, a country assignment was made if circumstances clearly indicated in whose waters the event occurred, e.g., apparent duration of the entanglement and distance from the US/Canadian border (Hague Line). If there was a reasonable chance that the event may have occurred in either country's waters, the country assignment was listed as unknown. Events placed in this category were typically mortalities either first detected near the US/Canadian border, mortalities involving severely decomposed carcasses that potentially drifted across jurisdictions, or mortalities stemming from chronic entanglement injuries.

Vessel collision mortalities were assigned to a country according to the location of the carcass. This differs from the country assignment of entanglements because vessel collision mortalities are more likely to be instantaneous (Kraus 1990; Moore et al. 2004). The country assignment was listed as unknown if there was an equal chance that the event occurred in either country's waters.

¹ See <http://www.greateratlantic.fisheries.noaa.gov/protected/whaletrp/reports/index.html>

² See <http://www.nmfs.noaa.gov/pr/pdfs/fr/fr60-45086.pdf>

RESULTS

From 2008 through 2012, a total of 300 mortalities was documented, of which 60 (20%) were confirmed to be the result of human interactions, 15 (5%) were due to natural causes, and 225 (75%) had insufficient evidence to determine cause of death. Of the human-caused mortalities, 32 were due to entanglement and 28 were due to vessel strike. Table 1 gives the tallies of mortalities and the cause of death by stock for the period. Table 2 provides details, by stock, of each confirmed human interaction event that resulted in mortality. The LOF column indicates entanglement events that would be used in categorizing any US fishery responsible in the LOF if the gear or fishery could be identified. There were 22 entanglement events that warranted inclusion in the LOF during the period. In 4 of these cases the gear type was identified, but the specific fishery was not.

Over the 5 year period, there were 18 verified mortalities of North Atlantic right whales. Of these, 6 were due to entanglements, 2 due to vessel strikes, 4 due to natural causes, and 6 for which the cause was undetermined.

Humpbacks had the highest number of mortalities, with 93 carcasses detected. Of these mortalities, 8 were due to entanglement, 7 due to vessel strike, 3 due to natural causes, and cause of death was undetermined for the remaining 75 events. We assumed all humpback events occurring in or near US and southeast Canadian waters involved the Gulf of Maine stock unless a whale was confirmed to be from another stock. Humpback events from Labrador and Newfoundland were assumed to not involve the Gulf of Maine stock and are therefore not included in the tallies.

There were 32 verified fin whale mortalities: 3 were due to entanglement, 9 due to vessel strikes, 4 due to natural causes, and 16 were from undetermined causes.

Of the 7 events involving sei whale mortalities, 1 was attributed to entanglement, 2 to vessel strikes, and cause of death was undetermined for the remaining 4 events.

Minke whales were involved in 91 mortality events, of which 14 were due to entanglements, 6 due to vessel strikes, 4 due to natural causes, and cause of death was undetermined for the remaining 67 events.

Bryde's whales had 2 documented mortalities. One was the result of a vessel strike. The cause of the other death could not be determined.

Blue whales had 2 documented mortalities. The cause of death could not be determined for either case.

In 55 of the 300 confirmed unique large whale mortality events during 2008 - 2012, species identification was not possible. In 2 of the 55 events, the similarity in body shape and size between fin and sei whales prevented us from distinguishing which of these 2 species were involved. In another 12 events, the whales could only be identified as balaenopteridae based on the presence of ventral pleats. The taxonomic identity of the whales involved in the remaining 41 events could not be assigned to genus. Of these mortalities involving unidentified species, 1 was attributed to vessel strike. The cause of death was not determined for the remaining 54 mortalities.

Mortality tallies by country are presented in Table 3 and include a total of 10 Canadian entanglements (6 identified by gear and 4 assigned by location of first detection) and 10 US entanglements (4 identified by gear and 6 assigned by location of first detection). The remaining 12 entanglement events that resulted in mortality could not be assigned to either country's waters with certainty. Twenty-six of 28 confirmed collision events leading to death were first detected in

US waters. One event was found close to the US and Canadian boundary, so the location of the interaction could not be established. The remaining event involved an international commercial vessel carrying a fresh, unidentified whale carcass on its bow. It could not be determined if the strike occurred in US or international waters.

DISCUSSION

Our criteria attempt to encompass all event scenarios and minimize the likelihood of incorrectly assigning whale mortalities to human causes. The resulting values provide a minimum count of confirmed human-caused mortality for baleen whale stocks along the Gulf of Mexico, US east coast, and the Atlantic Canadian provinces. These values do not include observed serious injuries presumed to result in mortality (Cole and Henry, in prep). Despite the minimum values, the annual confirmed human-caused mortality rates exceed PBR for 4 of the 7 stocks examined, including North Atlantic right, humpback, sei, and Bryde's whales, with PBRs of 0.9, 2.7, 0.5 and 0.1, respectively (Waring et al, in prep).

Differentiating injuries that cause mortalities from preexisting injuries or postmortem damage is problematic but can be accomplished through necropsy or, in many cases, parsimonious evaluation of available evidence. For example, fishing line constrictions on a whale carcass can be considered circumstantial evidence of premortem entanglement, as these constrictions are likely the result of force applied by an active animal. Large linear (nonwrapping) lacerations can be considered an indication of a premortem vessel collision since only whales at depth would be exposed to the propellers of a ship; floating carcasses would likely be pushed aside by the ship's bow wave (Knowlton et al. 1995).

However, carcasses detected at sea often cannot be examined sufficiently for either internal or even external indications of anthropogenic injury. Most notably, vessel collision mortalities frequently lack external evidence and may not be detected unless a necropsy is conducted. Necropsies frequently identify subdermal hemorrhaging or hematomas, the result of blunt trauma and the circulation of blood at the time of injury.

Given the likelihood that not all entanglement and vessel strike mortalities are detected, that the criteria applied here are designed to minimize the likelihood of incorrectly assigning whale mortalities to human causes, and that observed serious injuries are not considered here, the numbers in this report represent minimum values for confirmed human-caused mortality to baleen whale stocks in US waters of the Gulf of Mexico, the US eastern seaboard, and in the Atlantic Canadian provinces. The true level of anthropogenic impact to these stocks is greater than these minimum values, but the amount greater is unknown.

ACKNOWLEDGEMENTS

We are especially grateful to the Gulf of Mexico, US, and Canadian Maritime Provinces and Newfoundland stranding and entanglement networks, whose members searched for and examined whales both live and dead. It is a difficult, dirty, and ceaseless job that deserves special recognition. The United States Coast Guard was instrumental in conveying sightings reported by mariners, investigating carcasses at sea, and assisting in disentanglement efforts. We are also grateful to the staffs of the Center for Coastal Studies (CCS), New England Aquarium (NEA), Whale Center of New England (WCNE), NOAA aerial survey teams, Sea to Shore Alliance, the states of Florida and Georgia, Northeast Fisheries Observer Program, Marine Animal Response Society, Whale Release and Stranding Group, and many others for providing the sightings that have allowed this work to be conducted. Betty Lentell, Kathryn Roy, and William Greer assisted in verifying fishery observer program records. NEA, CCS and WCNE provided sighting histories and demographic information. Jooke Robbins (CCS), Laura Ganley (CCS), and Katie Jackson (Florida Fish and Wildlife Conservation Commission) were particularly helpful with sighting, photo, and health histories. Members of the Atlantic Scientific Review Group have provided numerous helpful comments on the protocols described here. We also thank the anonymous reviewers of earlier drafts of this report.

REFERENCES CITED

- Cole TVN, Henry AG. 2013. Serious injury determinations for baleen whale stocks along the Gulf of Mexico, United States East Coast and Atlantic Canadian Provinces, 2007-2011. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 13-24; 14 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://nefsc.noaa.gov/publications/>
- Cole TVN, Henry AG. In prep. Serious injury determinations for baleen whale stocks along the Gulf of Mexico, United States East Coast and Atlantic Canadian Provinces, 2008-2012.
- Henry AG, Cole TVN, Hall L, Ledwell W, Reid A. 2013. Mortality determinations for baleen whale stocks along the Gulf of Mexico, United States east coast and Atlantic Canadian provinces, 2007-2011. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 13-18; 15 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://nefsc.noaa.gov/publications/>
- Knowlton AR, Korsmeyer FT, Kerwin JE, Wu H, Hynes B. 1995. The hydrodynamic effects of large vessels on right whales. Boston (MA): New England Aquarium. Final report to NOAA Fisheries for Contract No. 40EANFF400534.
- Kraus SD. 1990. Rates and potential causes of mortality in North Atlantic right whales (*Eubalaena glacialis*). Mar Mamm Sci. 6(4):278-291.
- Moore MJ, Knowlton AR, Kraus SD, McLellan WA, Bonde RK. 2004. Morphometry, gross morphology and available histopathology in North Atlantic right whale (*Eubalaena glacialis*) mortalities (1970–2002). J Cetacean Res Manage. 6(3):199-214.
- Palsbøll PJ, Allen J, Anderson TH, Bérubé M, Clapham PJ, Feddersen TP, Friday N, Hammond P, Jørgensen H, Katona SK, Larsen AH, Larsen F, Lien J, Mattila DK, Nygaard FB, Robbins J, Sponer R, Sears R, Sigurjónsson J, Smith TD, Stevick PT, Vikingsson G, Øien N. 2001. Stock structure and composition of the North Atlantic humpback whale, *Megaptera novaeangliae*. Cambridge (UK): International Whaling Commission. Presented to International Whaling Commission Scientific Committee. SC/53/NAH11.
- Wade PR, Angliss RP. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop, April 3-5, 1996, Seattle, Washington. Seattle (WA): NMFS Alaska Fisheries Science Center. US Dept of Commer, NOAA Tech. Memo, NMFS-OPR-12.
- Waring GT, Josephson E, Maze-Foley K, Rosel PE, editors. 2014. U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments -- 2013. NOAA Tech Memo NMFS NE 228; 464 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://www.nefsc.noaa.gov/nefsc/publications/>
- Waring GT, Josephson E, Maze-Foley K, Rosel PE, eds. In prep. U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments – 2014.

Table 1. Summary of all unique large whale mortalities observed along the Gulf of Mexico Coast, US East Coast, and Atlantic Canadian Provinces, 2008-2012. Determinations of human-caused mortality follow the criteria established by the NEFSC.

Species	Western North Atlantic right whale (<i>Eubalaena glacialis</i>)	Gulf of Maine humpback whale (<i>Megaptera novaeangliae</i>)	Western North Atlantic fin whale (<i>Balaenoptera physalus</i>)	Nova Scotian sei whale (<i>B.borealis</i>)	Western North Atlantic blue whale (<i>B. musculus</i>)	Canadian East Coast minke whale (<i>B. acutorostrata</i>)	Northern Gulf of Mexico Bryde's whale (<i>B. edeni</i>)	Unidentified fin/sei whale	Unidentified balaenopterid ^a	Unidentified whale spp.	Totals
Total confirmed mortalities	18 (3, 4, 4, 4, 3)	93 (28, 20, 19, 12, 14)	32 (5, 5, 6, 8, 8)	7 (3, 2, 0, 1, 1)	2 (0, 0, 1, 1, 0)	91 (16, 9, 15, 25, 26)	2 (0, 1, 0, 0, 1)	2 (0, 0, 2, 0, 0)	12 (0, 1, 5, 4, 2)	41 (6, 6, 10, 14, 5)	300
Confirmed entanglement mortalities	6 (0, 0, 3, 1, 2)	8 (2, 2, 4, 0, 0)	3 (0, 0, 0, 3, 0)	1 (1, 0, 0, 0, 0)	0	14 (4, 0, 0, 4, 6)	0	0	0	0	32
Confirmed vessel strike mortalities	2 (0, 0, 1, 1, 0)	7 (1, 0, 3, 3, 0)	9 (1, 1, 2, 1, 4)	2 (0, 1, 0, 1, 0)	0	6 (0, 1, 1, 3, 1)	1 (0, 1, 0, 0, 0)	0	0	1 (0, 0, 1, 0, 0)	28
Confirmed mortalities, NOT vessel strike or entanglement	4 (3, 1, 0, 0, 0)	3 (0, 0, 1, 0, 2)	4 (1, 1, 1, 0, 1)	0	0	4 (0, 1, 1, 1, 1)	0	0	0	0	15
Confirmed mortalities, IITD ^b	6 (0, 3, 0, 2, 1)	75 (25, 18, 11, 9, 12)	16 (3, 3, 3, 4, 3)	4 (2, 1, 0, 0, 1)	2 (0, 0, 1, 1, 0)	67 (12, 7, 13, 17, 18)	1 (0, 0, 0, 0, 1)	2 (0, 0, 2, 0, 0)	12 (0, 1, 5, 4, 2)	40 (6, 6, 9, 14, 5)	225
Annual Human-Caused Mortality Rate (EN, VS)	1.6 (1.2, 0.4)	3 (1.6, 1.4)	2.4 (0.6, 1.8)	0.6 (0.2, 0.4)	0	4 (2.8, 1.2)	0.2 (0, 0.2)	0	0	0.2 (0, 0.2)	12 (6.4, 5.6)

^a Described as having throat grooves (rorqual pleats).

^b IITD = insufficient information to determine cause of death or if the injury was serious and likely lethal.

Table 2. Confirmed human-caused mortality records of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012.

Date	Individual	Catalog #	General Location	Cause of Fate	Country of Origin ^a	Country Conf Code ^b	Gear Type ^c	Comments	LOF ^d
Western North Atlantic right whale (<i>Eubalaena glacialis</i>)									
27-Jun-10	Tips	1124	off Cape May, NJ	EN	XU	CE	NR	Evidence of constricting rostrum, mouth & pectoral wraps w/ associated hemorrhage & bone damage	1
02-Jul-10	U	-	off Great Wass Island, ME	VS	XU	CE	-	2 large lacerations from dorsal to ventral surface	-
12-Aug-10	Trident	1113	Digby Neck, NS	EN	XC	CE	NP	Evidence of entanglement w/ associated hemorrhaging around right pectoral	1
25-Dec-10 ^e	Bayla	3911	off Jacksonville, FL	EN	XU	CE	GU	Embedded line in mouth and pectoral; severe health decline; proximate COD - entanglement, ultimate COD - shark predation	1
16-Mar-11	U	-	Cape Romain, SC	EN	XU	CE	GU	Multiple wraps embedded in right pectoral bones	1
27-Mar-11	-	1308	Nags Head, NC	VS	US	AE	-	Fractured right skull	-
19-Jul-12	U	-	Clam Bay, NS	EN	XC	CE	NR	Multiple constricting wraps on peduncle; COD - peracute underwater entrapment	1
18-Dec-12	-	4193	off Palm Coast, FL	EN	US	GI	PT	Constricting and embedded wraps with associated hemorrhaging at peduncle, mouthline, tongue, oral rete, rostrum and pectoral; malnourished	1

Table 2, continued. Confirmed human-caused mortality records of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012.

Individual	Catalog #	General Location	Cause of Fate	Country of Origin ^a	Country Conf Code ^b	Gear Type ^c	Comments	LOF ^d
Gulf of Maine humpback whale (<i>Megaptera novaeangliae</i>)								
30-May-08	U	Georges Bank	EN	XU	CE	NR	Constricting body wraps, one wrap under lower jaw; open wound on right pectoral	1
09-Jun-08	U	Georges Bank	EN	US	GI	PT	Constricting body wrap	1
04-Nov-08	U	Assateague Island, MD	VS	US	AE	-	Cranial fractures w/ associated hemorrhaging	-
08-Feb-09	U	Cape Fear, NC	EN	XU	CE	NP	Evidence of entanglement at mouthline, peduncle, & pectoral w/ associated hemorrhaging; emaciated	1
16-Feb-09	U	Nags Head, NC	EN	XU	CE	NP	Evidence of entanglement involving anchoring or heavily weighted gear w/ associated hemorrhaging	1
13-Mar-10	U	Ocean City, MD	VS	US	AE	-	Skull fractures w/ associated hemorrhaging	-
08-May-10	U	Narragansett, RI	EN	CN	GI	GN	Evidence of constricting gear w/ associated hemorrhaging; fluid filled lungs	-
15-May-10 ^e	U	Cape Hatteras Inlet, NC	EN	XU	CE	NP	Necrotic infected wounds at fluke insertion; chronic abrasions on head; proximate COD - entanglement, ultimate COD - euthanasia	1
28-May-10	U	Edgartown, MA	EN	XU	CE	GU	Evidence of entanglement w/ associated bruising & edema	1
10-Jun-10	U	Jones Beach State Park, NY	VS	US	AE	-	Extensive hemorrhage & edema on right dorsal lateral surface	-

Table 2, continued. Confirmed human-caused mortality records of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012.

Individual	Catalog #	General Location	Cause of Fate	Country of Origin ^a	Country Conf Code ^b	Gear Type ^c	Comments	LOF ^d
Gulf of Maine humpback whale (<i>Megaptera novaeangliae</i>)								
04-Jul-10	U	off Assateague, MD	VS	US	AE	-	Extensive hemorrhage & edema to left lateral area	-
27-Nov-10	U	Bay of Fundy	EN	XC	CE	NR	Evidence of constricting wraps on fluke, peduncle, & pectoral	1
07-Mar-11 ^e	U	Thorofare Bay, Core Sound, NC	VS	US	AE	-	8 deep lacerations across back; robust with anemia & pale musculature indicates exsanguination; proximate COD - vessel strike, ultimate COD - euthanasia	-
05-May-11	U	Little Compton, RI	VS	US	AE	-	Hemorrhaging at left jaw associated w/ blunt trauma; evidence of healing entanglement injuries	-
27-May-11	U	Barnegat Inlet, NJ	VS	US	AE	-	5 broken vertebral processes along left side w/ associated hemorrhaging	-
Western North Atlantic fin whale (<i>Balaenoptera physalus</i>)								
02-Jul-08	U	Barnegat Inlet, NJ	VS	US	AE	-	Vertebral fractures w/ associated hemorrhaging; hemorrhaging around ball joint of right pectoral	-
01-Oct-09	U	Port Elizabeth, NJ	VS	US	AE	-	Fresh carcass w/ broken pectoral, hematomas, & abrasions	-
18-Mar-10	U	off Bethany Beach, DE	VS	US	AE	-	Fractured skull w/ associated hemorrhaging; abrasion mid-dorsal consistent w/ being folded over the bow of a ship	-

Table 2, continued. Confirmed human-caused mortality records of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012.

Individual	Catalog #	General Location	Cause of Fate	Country of Origin ^a	Country Conf Code ^b	Gear Type ^c	Comments	LOF ^d
Western North Atlantic fin whale (<i>Balaenoptera physalus</i>)								
03-Sep-10	U	Cape Henlopen State Park, DE	VS	US	AE	-	Large laceration & vertebral fractures w/ associated hemorrhaging	-
01-Jan-11	U	off Portland, ME	EN	XU	CE	NP	Fresh carcass w/ evidence of constricting gear	1
05-Jun-11	U	off Long Beach, NJ	VS	US	AE	-	Extensive hemorrhage & soft tissue damage to the dorsal & right lateral thoracic region	-
24-Jul-11	U	Cheticamp, NS	EN	CN	AE	NP	Fresh carcass w/ evidence of extensive entanglement	-
21-Sep-11	U	off Atlantic City, NJ	EN	US	AE	NP	Fresh carcass w/ evidence of extensive entanglement	1
23-Jan-12	U	Ocean City, NJ	VS	US	AE	-	Hemorrhaging along right, midlateral surface; fish in stomach indicated feeding	-
19-Feb-12	U	Norfolk, VA	VS	US	AE	-	Deep laceration on head; skeletal fractures of rostrum and vertebrae; extensive hemorrhaging	-
10-Aug-12	U	Hampton Bays, NY	VS	US	AE	-	Extensive bruising along right lateral and ventral aspects	-
07-Oct-12	U	Boston Harbor, MA	VS	US	AE	-	Deep mid-line impression with associated hemorrhaging consistent with being folded across bow of ship	-

Table 2, continued. Confirmed human-caused mortality records of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012.

Individual	Catalog #	General Location	Cause of Fate	Country of Origin ^a	Country Conf Code ^b	Gear Type ^c	Comments	LOF ^d
Nova Scotian sei whale (<i>Balaenoptera borealis</i>)								
29-Jun-08	U	Slack's Cove, NB	EN	CN	AE	NP	Extensive entanglement evident	-
19-May-09	U	off Rehobeth Beach, DE	VS	US	AE	-	Posterior portion of skull & right mandible fractured; hemorrhaging dorsal to left pectoral	-
26-Mar-11	U	Virginia Beach, VA	VS	US	AE	-	Jaw, scapula, rib & vertebral fractures along right side w/ associated hemorrhaging	-
Canadian East Coast minke whale (<i>Balaenoptera acutorostrata</i>)								
14-Jun-08	U	Orleans, MA	EN	US	AE	NP	Braided line impressions wrapped body in 3 places & left a deep, hemorrhaged laceration across the rostrum & blowholes; hemorrhaged abrasions present on roof of mouth; wet, blood-filled lungs indicate drowning	1
23-Jul-08	U	Kelligrews, NL	EN	CN	AE	GU	Constricting wraps of gear on caudal peduncle	-
26-Jul-08	U	Conception Bay, NL	EN	CN	GI	GN	Constricting wraps of gear through mouth & around tail	-
21-Aug-08 ^t	U	off Richibucto Cape, NB	EN	CN	AE	NR	Evidence of constricting body wraps	-
20-May-09	U	off Point Pleasant, NJ	VS	US	AE	-	Large hemorrhage at right pectoral	-

Table 2, continued. Confirmed human-caused mortality records of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012.

Individual	Catalog #	General Location	Cause of Fate	Country of Origin ^a	Country Conf Code ^b	Gear Type ^c	Comments	LOF ^d
Canadian East Coast minke whale (<i>Balaenoptera acutorostrata</i>)								
09-Jul-10	U	Fire Island, NY	VS	US	AE	-	3-4 large dorsal lacerations associated w/ fractured ribs	-
06-May-11	U	off Martha's Vineyard, MA	EN	US	GI	PT	Anchored in gear; embedded line at fluke; evidence of entanglement w/ associated hemorrhaging at mouth corners & insertion of pectorals	1
04-Aug-11	U	off Sandy Hook, NJ	VS	US	AE	-	4 propeller lacerations across dorsal surface; fractured ribs w/ associated hemorrhaging	-
26-Aug-11	U	off Sandy Hook, NJ	EN	US	AE	NP	Fresh carcass w/ evidence of extensive entanglement	1
29-Aug-11	U	Moriches, NY	VS	US	AE	-	Extensive hemorrhage & edema along dorsal & both lateral surfaces	-
06-Oct-11	U	off Matinicus Island, ME	EN	US	GI	PT	Fresh carcass anchored in gear	1
07-Dec-11	U	Carolina Beach, NC	VS	US	AE	-	Healed deep & superficial propeller lacerations; internal lesions associated w/ deep lacerations indicative of peritonitis & infection	-
19-Dec-11	U	Bay of Fundy	EN	CN	GI	PT	Live entanglement; recovered dead in gear the following day; constricting peduncle wraps	-
16-Mar-12	U	Ipswich, MA	EN	US	AE	NP	Evidence of extensive, constricting gear with associated hemorrhaging	1

Table 2, continued. Confirmed human-caused mortality records of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012.

Individual	Catalog #	General Location	Cause of Fate	Country of Origin ^a	Country Conf Code ^b	Gear Type ^c	Comments	LOF ^d
Canadian East Coast minke whale (<i>Balaenoptera acutorostrata</i>)								
23-Jun-12	U	Container Terminal Port, Newark, NJ	VS	US	AE		Fresh carcass on bow of ship; deep laceration across ventral surface; COD - disembowelment and hypovolemic shock	-
26-Jun-12	U	off Renew's Rock, NL	EN	CN	GI	PT	Fresh carcass with constricting gear around peduncle	-
30-Jun-12	U	off Campbell Cove, Naufrage, PEI	EN	CN	GI	PT	Fresh carcass anchored in gear	-
01-Jul-12	U	East Point, Northern Lake Harbor, PEI	EN	CN	GI	PT	Constricting gear with associated hemorrhaging; COD - drowning	-
05-Aug-12	U	Chatham, MA	EN	US	AE	NR	Multiple constricting wraps through and around mouth and on fluke blades; COD - acute underwater entrapment	1
04-Oct-12	U	off Cliff Island, ME	EN	US	AE	NR	Evidence of constricting gear at mouthline, across ventral pleats, and at peduncle	1
Northern Gulf of Mexico Bryde's whale (<i>Balaenoptera edeni</i>)								
04-Oct-09	U	Tampa, FL	VS	US	AE		Vertebral separation; lung damage; subdermal contusions	-

Notes:

a. CN=Canada, US=United States, XC=Unk 1st sight in CN, XU=Unk 1st sight in US

b. GI=gear identified, AE=acute/very recent event or confirmed location, CE=chronic/prolonged event without indication of initial location

c. GN=gillnet, GU=gear unidentifiable, NP=none present, NR=none recovered/received, PT=pot/trap

d. Consider for List of Fisheries inclusion (1=yes)

e. Previously reported as a Serious Injury (Cole and Henry 2013)

f. Previously reported as two separate events: 1 live injury on 8/21/10 (Cole and Henry 2013) and 1 mortality on 8/25/08 (Henry et al 2013). Events were confirmed to be the same animal

Table 3. Summary of country of origin for all confirmed human-caused mortalities of baleen whale stocks along the Gulf of Mexico, US East Coast, and Atlantic Canadian Provinces, 2008-2012

Event Location	Event Type	Western North Atlantic right whale (<i>Eubalaena glacialis</i>)	Gulf of Maine humpback whale (<i>Megaptera novaeangliae</i>)	Western North Atlantic fin whale (<i>Balaenoptera physalus</i>)	Nova Scotian sei whale (<i>B.borealis</i>)	Western North Atlantic blue whale (<i>B. musculus</i>)	Canadian East Coast minke whale (<i>B. acutorostrata</i>)	Northern Gulf of Mexico Bryde's whale (<i>B. edeni</i>)	Unidentified fin/sei whale	Unidentified balaenopterid ^a	Unidentified whale spp.	Totals
US waters	Entanglement	1	1	1	0	0	7	0	0	0	0	10
	Vessel Strike	1	7	9	2	0	6	1	0	0	0	26
Canadian waters	Entanglement	0	1	1	1	0	7	0	0	0	0	10
	Vessel Strike	0	0	0	0	0	0	0	0	0	0	0
Unassigned waters	Entanglement	5	6	1	0	0	0	0	0	0	0	12
	Vessel Strike	1	0	0	0	0	0	0	0	0	1	2

^a Described as having throat grooves (rorqualn pleats).

Procedures for Issuing Manuscripts in the *Northeast Fisheries Science Center Reference Document (CRD) Series*

Clearance

All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If any author is not a federal employee, he/she will be required to sign an "NEFSC Release-of-Copyright Form." If your manuscript includes material from another work which has been copyrighted, then you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

For more information, NEFSC authors should see the NEFSC's online publication policy manual, "Manuscript/abstract/webpage preparation, review, and dissemination: NEFSC author's guide to policy, process, and procedure," located in the Publications/Manuscript Review section of the NEFSC intranet page.

Organization

Manuscripts must have an abstract and table of contents, and (if applicable) lists of figures and tables. As much as possible, use traditional scientific manuscript organization for sections: "Introduction," "Study Area" and/or "Experimental Apparatus," "Methods," "Results," "Discussion," "Conclusions," "Acknowledgments," and "Literature/References Cited."

Style

The CRD series is obligated to conform with the style contained in the current edition of the United States Government Printing Office Style Manual. That style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the CSE Style Manual. Manuscripts should be prepared to conform with these style manuals.

The CRD series uses the American Fisheries Society's guides to names of fishes, mollusks, and decapod

crustaceans, the Society for Marine Mammalogy's guide to names of marine mammals, the Biosciences Information Service's guide to serial title abbreviations, and the ISO's (International Standardization Organization) guide to statistical terms.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of cited works. Personal communications must include date, full name, and full mailing address of the contact.

Preparation

Once your document has cleared the review process, the Editorial Office will contact you with publication needs – for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as files on zip disks or CDs, email attachments, or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

Production and Distribution

The Editorial Office will perform a copy-edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once both the PDF (print) and Web versions of the CRD are ready, the Editorial Office will contact you to review both versions and submit corrections or changes before the document is posted online.

A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.

Research Communications Branch
Northeast Fisheries Science Center
National Marine Fisheries Service, NOAA
166 Water St.
Woods Hole, MA 02543-1026

**MEDIA
MAIL**

Publications and Reports of the Northeast Fisheries Science Center

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of living marine resources for the benefit of the nation through their science-based conservation and management and promotion of the health of their environment." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (*e.g.*, anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media. Currently, there are three such media:

NOAA Technical Memorandum NMFS-NE -- This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document -- This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review and most issues receive copy editing.

Resource Survey Report (formerly *Fishermen's Report*) -- This information report is a regularly-issued, quick-turnaround report on the distribution and relative abundance of selected living marine resources as derived from each of the NEFSC's periodic research vessel surveys of the Northeast's continental shelf. This report undergoes internal review, but receives no technical or copy editing.

TO OBTAIN A COPY of a *NOAA Technical Memorandum NMFS-NE* or a *Northeast Fisheries Science Center Reference Document*, either contact the NEFSC Editorial Office (166 Water St., Woods Hole, MA 02543-1026; 508-495-2350) or consult the NEFSC webpage on "Reports and Publications" (<http://www.nefsc.noaa.gov/nefsc/publications/>). To access *Resource Survey Report*, consult the Ecosystem Surveys Branch webpage (<http://www.nefsc.noaa.gov/femad/ecosurvey/mainpage/>).

ANY USE OF TRADE OR BRAND NAMES IN ANY NEFSC PUBLICATION OR REPORT DOES NOT IMPLY ENDORSEMENT.