

NASA/TM—1998-208508



Theoretical and Experimental Determination of the Proton Affinity of $(\text{CF}_3\text{CH}_2)_2\text{O}$

Michael J. Zehe
Lewis Research Center, Cleveland, Ohio

David W. Ball
Cleveland State University, Cleveland, Ohio

The NASA STI Program Office . . . in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.

The NASA STI Program Office is operated by Langley Research Center, the Lead Center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:

- **TECHNICAL PUBLICATION.** Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA's counterpart of peer-reviewed formal professional papers but has less stringent limitations on manuscript length and extent of graphic presentations.
- **TECHNICAL MEMORANDUM.** Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- **CONTRACTOR REPORT.** Scientific and technical findings by NASA-sponsored contractors and grantees.

- **CONFERENCE PUBLICATION.** Collected papers from scientific and technical conferences, symposia, seminars, or other meetings sponsored or cosponsored by NASA.
- **SPECIAL PUBLICATION.** Scientific, technical, or historical information from NASA programs, projects, and missions, often concerned with subjects having substantial public interest.
- **TECHNICAL TRANSLATION.** English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized data bases, organizing and publishing research results . . . even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at <http://www.sti.nasa.gov>
- E-mail your question via the Internet to help@sti.nasa.gov
- Fax your question to the NASA Access Help Desk at (301) 621-0134
- Telephone the NASA Access Help Desk at (301) 621-0390
- Write to:
NASA Access Help Desk
NASA Center for Aerospace Information
7121 Standard Drive
Hanover, MD 21076



Theoretical and Experimental Determination of the Proton Affinity of $(\text{CF}_3\text{CH}_2)_2\text{O}$

Michael J. Zehe
Lewis Research Center, Cleveland, Ohio

David W. Ball
Cleveland State University, Cleveland, Ohio

National Aeronautics and
Space Administration

Lewis Research Center

Trade names or manufacturers' names are used in this report for identification only. This usage does not constitute an official endorsement, either expressed or implied, by the National Aeronautics and Space Administration.

Available from

NASA Center for Aerospace Information
7121 Standard Drive
Hanover, MD 21076
Price Code: A03

National Technical Information Service
5287 Port Royal Road
Springfield, VA 22100
Price Code: A03

Theoretical and Experimental Determination of the Proton Affinity of $(\text{CF}_3\text{CH}_2)_2\text{O}$

Michael J. Zehe

National Aeronautics and Space Administration

Lewis Research Center

Cleveland, Ohio 44135

and

David W. Ball

Department of Chemistry

Cleveland State University

Cleveland, Ohio 44115

ABSTRACT

We report the experimental determination of the proton affinity of the molecule $(\text{CF}_3\text{CH}_2)_2\text{O}$ using chemical ionization mass spectrometry, and we compare it to the theoretical value obtained for protonation at the oxygen atom using the calculational methodology (MP2/6-31G**//MP2/3-21G). The proton affinity for this molecule as measured by bracketing experiments is between 724 kJ/mole and 741 kJ/mole. Ab initio (MP2/6-31G**//MP2/3-21G) calculations yield a value of about 729 kJ/mole, in agreement with the chemical ionization experiments. The results of these and related calculations suggest that the (MP2/6-31G**//MP2/3-21G) methodology is acceptable for estimating the proton affinities of partially- and fully-fluorinated methyl and ethyl ethers. We submit that any conclusions about the chemistry of fluoroether polymer lubricants based on their basicity can also be predicted reliably with such calculations.

INTRODUCTION

The polymeric perfluoroalkyl ethers have been the subject of intense interest in the technical community recently, because of their widespread use as liquid lubricants and their chemical behavior under long-term tribological use (Refs. 1 and 2). Of continuing interest is the chemical degradation which can occur when these fluids interact with metal or metal oxides surfaces in bearings or sliding contacts. There is a great deal of evidence that this degradation involves acid-base chemistry between weakly acidic surface sites and basic sites on the lubricant chain (Refs. 3 and 4).

The proton affinity (PA) of simple fluoroether molecules has been used to characterize their reactivity to surface acid sites. The primary experimental methods used for measurement of gaseous proton affinity use ion cyclotron resonance (ICR) spectroscopy for measurement of the proton transfer to another base, or chemical ionization (CI) measurements to “bracket” the compound between two bases of known PA. These measurements are experimentally difficult to perform and interpret (Ref. 5). Their accuracy depends upon the presence of chemical equilibrium between the species, a certainty in the local temperature in the measurement cell, and the assumption that the rotational and vibrational components of ΔS_T are zero or known. In addition, there may be uncertainty as to the site of protonation on the molecule. Despite these difficulties, there exists a scale of relative proton affinities which is well-established and well-accepted (Ref. 6).

An alternative to the measurement of proton affinities of small molecules is their determination through the use of high-level computational chemistry. Modern computational chemical methods can yield very accurate values for the proton affinity of molecules composed of first-row elements (Ref. 7). Delhalle et al. (Refs. 8 and 9) performed Hartree-Fock calculations for a number of small perfluoroether molecules protonated at the oxygen atoms and at fluorine; they reported that, relative to protonation at oxygen, protonation at an α -fluorine gave a substantially higher predicted PA. Orgel, Ball, and Zehe (Ref. 10) included electron correlation and performed calculations for a number of symmetrical partially and fully-fluorinated dimethyl ethers. Their results showed very good agreement with experimental values of the PA's of dimethyl and diethyl ether, but the lack of experimental data for fluoroether PA's prevented comparison.

In this communication, we report the experimental determination of the proton affinity of the molecule $(CF_3CH_2)_2O$ using CI mass spectrometry, and we compare it to the theoretical value obtained for protonation at the oxygen atom using the calculational methodology (MP2/6-31G**//MP2/3-21G). This methodology has

been used by the current authors (Refs. 10 and 11) to predict the PA of $(\text{CH}_3)_2\text{O}$ and $(\text{C}_2\text{H}_5)_2\text{O}$ within the acceptance limits of the more computationally expensive G2 theory, and this work represents in part a test of its applicability to fluorinated molecules.

EXPERIMENTAL

Chemical ionization mass spectrometry of $(\text{CF}_3\text{CH}_2)_2\text{O}$ was performed on a Finnigan TSQ-45 gas chromatograph/mass spectrometer using its standard CI sample volume with a filament as its electron source. The temperature of the sample volume was 100 °C. After the reagent gas (H_2O , CH_2O , or CH_3OH) was introduced and a stable discharge was established, the $(\text{CF}_3\text{CH}_2)_2\text{O}$ was introduced and the mass spectrum was measured and evaluated for the presence of $(\text{CF}_3\text{CH}_2)_2\text{OH}^+$. The observation of this ion indicates that the PA of $(\text{CF}_3\text{CH}_2)_2\text{O}$ is higher than that of the reagent gas.

CALCULATIONS

Ab initio calculations were performed using the GAUSSIAN 92 program (Ref. 12) on an IBM RS 6000 workstation or a Cray C-90 supercomputer. Geometry optimizations were first performed using the 3-21G basis set with an MP2 correction, and the resulting geometry was used for MP2 single point energy calculations with the 6-31G(d,p) basis set (designated MP2/6-31G**//MP2/3-21G). Initial geometries were chosen as the all-trans configurations (Refs. 13 and 14). The ether was assumed to protonate at the oxygen atom. Thermal energies were simultaneously computed, and the ΔH_{298} for the protonation reaction was determined using:

$$\Delta H_{298} = \Delta E_0 + \Delta E_{\text{therm}} + \Delta(\text{PV}) \quad (1)$$

where ΔE_0 is the difference in the total electronic energies of the species at 0 K, and ΔE_{therm} includes contributions from zero-point vibrational energy differences, thermal vibrational energy differences, rotational energy differences, and thermal translational energy differences. $\Delta(\text{PV})$ is the standard conversion from internal energy to enthalpy and for all proton affinity reactions equals $-RT$ ($= -2.48$ kJ/mol). All vibrational frequencies were scaled by a factor of 0.95 for thermochemical calculations.

RESULTS AND DISCUSSION

The molecular ion for $(\text{CF}_3\text{CH}_2)_2\text{O}$ appears at 182 amu, with the protonated form at 183 amu. The strongest peak in the proton CI spectra appears at 113 amu and is due to loss of CF_3H from the protonated ether. Also present is a strong peak at 163 amu, due to loss of HF from the protonated molecule. The intensity of the M+1 peak at 183 amu is <3 percent of the strongest peak at mass 113. This illustrates one of the challenges of measuring the mass spectra of protonated fluorocarbons: the facility with which stable fluorine species split off from the energized, protonated molecule.

The proton affinities of H_2O , formaldehyde and methanol are, respectively, 723.8, 741.4, and 773.6 kJ/mole (Ref. 6). CI mass spectra using an $\text{H}_2\text{O}/\text{H}_3\text{O}^+$ discharge resulted in a strong peak at mass 183, indicating efficient ionization of the base molecule by H_3O^+ . For $(\text{CF}_3\text{CH}_2)_2\text{O}$ in either a formaldehyde or methanol discharge, however, the mass 183 peak was absent. Thus these experiments show that the proton affinity of bis (2,2,2-trifluoroethylether) (g) is between that of H_2O (g) and CH_2O (g), or between about 724 and 741 kJ/mole.

All calculations converged on stationary points that were true minima as indicated by vibrational frequency calculations that yielded positive frequencies only. Conformational studies to verify that the final geometries were absolute minima were not performed, but it is our judgement that conformational contributions to the proton affinity are slight. Figures 1 and 2 show the structures of the optimized molecule and its protonated form. A summary of the structural parameters is given in Tables I and II; we plan to publish a more detailed structural analysis in a forthcoming paper (Ref. 11).

The neutral molecule has C_{2v} symmetry, with a coplanar arrangement of all five ether backbone atoms and two of the terminal fluorine atoms. Protonation to $(\text{CF}_3\text{CH}_2)_2\text{OH}^+$ changes the C-O-C bond distances and angles only slightly, but now the terminal carbon atoms have rotated out of the C-O-C plane by 42 degrees, and the CF_3 groups have themselves rotated by a few degrees. The result is an ion with C_s symmetry, with an (O)H—F distance of about 2.12Å. This distance is well within the limits that define hydrogen bonding, (Ref. 15) and we suggest that intramolecular hydrogen bonding is contributing to the overall stability of the protonated ether. It is difficult to tell the extent to which the intramolecular hydrogen bonding is contributing to the overall stability, but it may be substantial, based upon common H-bond energies of 10 to 35 kJ/mole (Ref. 15).

Table III lists a summary of the calculated single-point energies of the unprotonated and protonated ethers using different levels of theory, with an indication of the predicted proton affinity. The MP2/6-31G**//MP2/3-21G calculations predict a proton affinity of about 729 kJ/mole, in agreement with the chemical ionization experiments. We conclude that calculations at this level can be considered reliable for predicting the proton affinities of related compounds.

Earlier work (Ref. 10 and 11) showed that MP2/6-31G**//MP2/3-21G calculations did a very good job of predicting proton affinities of dimethyl ether and its fluorinated derivatives. In several cases, the calculational results approached the level of performance expected for the G2 level of theory (Ref. 16). We suggest, therefore, that the current methodology is acceptable for estimating the proton affinities of partially- and fully-fluorinated methyl (Ref. 10) and ethyl (Ref. 11) ethers. We further submit that any conclusions about the chemistry of fluoroether polymer lubricants based on their basicity can also be predicted reliably with such calculations. Specifically, the approximately 80 to 100 kJ/mole (excluding intramolecular H-bonding) decrease in the proton affinity which results when diethyl ether is fully fluorinated, indicates that caution must be exercised when attributing perfluoroether lubricant decomposition to acid-base chemistry at oxygen.

REFERENCES

1. P. Herrera-Fierro, W.R. Jones, S.V. Pepper, *J. Vac. Sci. Tech. A* **1993**, 11, 354.
2. J. Pacansky, M. Miller, W. Hutton, V. Liu, A. Scheiner, *J. Am. Chem. Soc.* **1991**, 113, 329.
3. M.J. Zehe, O. Faut, *STLE Trans.* **1990**, 33, 634.
4. P. Kasai, *Macromolecules* 1992, 25, 6791.
5. M. Meot-Ner, L.W. Sieck, *JACS* 1991, 113, 4460.
6. S.G. Lias, J.F. Liebman, R.D. Levin, *J. Phys. Chem. Ref. Data* **1984**, 13, 695.
7. B.J. Smith, L. Radom, *JACS* 1993, 115, 4885.
8. M. Fontaine, J. Delhalle, M. Defranceschi, G. Lecayon, *Int. J. Quant. Chem.* 1993, 46, 171.
9. M. Fontaine, J. Delhalle, M. Defranceschi, J.M.M. Bourin, *J. Mol. Struct.* 1993, 300, 10.
10. V.B. Orgel, D.W. Ball, M.J. Zehe, *Journal of Molecular Structure(Theochem)* **1997**, 417, 195.

11. M.J. Zehe, D. Ball, "Optimized Structures and Proton Affinities of Fluorinated Diethyl Ethers: An Ab Initio Study," manuscript in preparation.
12. GAUSSIAN 92. Revisions B and D.2. M.J. Frisch, G.W. Trucks, M. Head-Gordon, P.M. W. Gill, M.W. Wong, J. B. Foresman, B.G. Johnson, H. B. Schlegel, M.A. Robb, E.S. Replogle, R. Gomperts, J.L. Andres, K. Raghavachari, J.S. Binkley, C. Gonzales, R.L. Martin, D.J. Fox, D.J. Defrees, J. Baker, J.J.P. Stewart, and J. A. Pople, Gaussian, Inc., Pittsburgh, PA, 1992.
13. G.D. Smith, R.L. Jaffe, D.Y. Yoon, *Macromolecules* **1994**, 27(12), 3166.
14. G.D. Smith, R.L. Jaffe, D.Y. Yoon, *Macromolecules* **1995**, 28(17), 5804.
15. Pimentel, George C., "The Hydrogen Bond," Reinhold Pub. Corp., New York, 1960.
16. L.A. Curtiss, K. Raghvachari, G.W. Trucks, J.A. Pople, *J. Chem. Phys.* **94** (1991), 7221.




Figure 1.—MP2/3-21G Optimized Geometry for $(\text{CF}_3\text{CH}_2)_2\text{O}$ (C_{2v}).

TABLE I.—MP2/3-21G OPTIMIZED
STRUCTURAL PARAMETERS
FOR $(\text{CF}_3\text{CH}_2)_2\text{O}$ (C_{2v})

Bond	Value
C1-O-C2	116.0°
C-O	1.46 Å
C1-C3	1.51 Å
C3-C1-O-C2	145.5°
C3-F1	1.38 Å
C3-F2	1.37 Å
F1-C3-C1-O	180°




Figure 2.—MP2/3-21G Optimized Geometry for $(\text{CF}_3\text{CH}_2)_2\text{OH}^+$ (C_s).

TABLE II.—MP2/3-21G OPTIMIZED
STRUCTURAL PARAMETERS FOR
 $(\text{CF}_3\text{CH}_2)_2\text{OH}^+$ (C_s)

Bond	Value
C1-O-C2	121.9°
C-O	1.52 Å
C1-C3	1.53 Å
C3-C1-O-C2	169.1°
O-H	1.01 Å
H-O-C1-C3	35.6°
F-H (min)	1.91 Å

TABLE III.—CALCULATED ENERGIES (HARTREES) AND ESTIMATED PROTON
AFFINITY FOR (CF₃CH₂)₂O⁺ BASED ON THE STRUCTURE FROM MP2/3-21G
OPTIMIZATIONS

	HF/3-21G	6-31G//MP2/3-21G	6-31G*//MP2/3-21G	6-31G**//MP2/3-21G
(CF ₃ CH ₂) ₂ O	−820.82442	−826.21700	−827.00468	−827.03589
(CF ₃ CH ₂) ₂ OH ⁺	−821.11574	−826.49085	−827.28370	−827.32412
PA, kJ/mole	736.6	690.7	704.3	728.5

^aCalculation includes changes in energy due to thermal vibrational and rotational energy and ΔPV; see text for details.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE December 1998		3. REPORT TYPE AND DATES COVERED Technical Memorandum
4. TITLE AND SUBTITLE Theoretical and Experimental Determination of the Proton Affinity of (CF ₃ CH ₂) ₂ O			5. FUNDING NUMBERS WU-523-22-13-00	
6. AUTHOR(S) Michael J. Zehe and David W. Ball				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Lewis Research Center Cleveland, Ohio 44135-3191			8. PERFORMING ORGANIZATION REPORT NUMBER E-11358	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, DC 20546-0001			10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA TM-1998-208508	
11. SUPPLEMENTARY NOTES Michael J. Zehe, NASA Lewis Research Center, and David W. Ball, Cleveland State University, Department of Chemistry, Cleveland, Ohio 44115. Responsible person, Michael J. Zehe, organization code 5140, (216) 433-5833.				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified - Unlimited Subject Categories: 05, 19, 07 and 25 This publication is available from the NASA Center for AeroSpace Information, (301) 621-0390.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) We report the experimental determination of the proton affinity of the molecule (CF ₃ CH ₂) ₂ O using chemical ionization mass spectrometry, and we compare it to the theoretical value obtained for protonation at the oxygen atom using the calculational methodology (MP2/6-31G**//MP2/3-21G). The proton affinity for this molecule as measured by bracketing experiments was between 724 kJ/mole and 741 kJ/mole. Ab initio (MP2/6-31G**//MP2/3-21G) calculations yield a value of about 729 kJ/mole, in agreement with the chemical ionization experiments. The results of these and related calculations suggest that the (MP2/6-31G**//MP2/3-21G) methodology is acceptable for estimating the proton affinities of partially- and fully-fluorinated methyl and ethyl ethers. We submit that any conclusions about the chemistry of fluoroether polymer lubricants based on their basicity can also be predicted reliably with such calculations.				
14. SUBJECT TERMS Quantum chemical; Lubricants; Modelling; Semiempirical; Lubrication; Perfluoroethers			15. NUMBER OF PAGES 15	
			16. PRICE CODE A03	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT	