


STUDIES IN
TERTIARY
STRATIGRAPHY OF
THE CALIFORNIA
COAST RANGES



GEOLOGICAL SURVEY
PROFESSIONAL PAPER 1213


**COVER PHOTOGRAPH of benthic foraminifers from
the Bear Creek area, Santa Cruz Mountains**

Studies in Tertiary Stratigraphy of the California Coast Ranges

Edited by EARL E. BRABB

GEOLOGICAL SURVEY PROFESSIONAL PAPER 1213

*A series of closely related biostratigraphic studies
that define the character, age, and correlation
of microfossils from Tertiary rocks of the
California Coast Ranges, as a basis for correlating
with international stratigraphic standards*



UNITED STATES DEPARTMENT OF THE INTERIOR

JAMES G. WATT, *Secretary*

GEOLOGICAL SURVEY

Dallas L. Peck, *Director*

Library of Congress Cataloging in Publication Data

Studies in Tertiary stratigraphy of the California Coast Ranges.

(Geological Survey professional paper ; 1213)

Bibliography: p. 78-82

Supt. of Docs. no.: I 19.16:1213

1. Geology, stratigraphic--Tertiary. 2. Stratigraphic correlation--
Coast Ranges. 3. Geology--California. I. Brabb, Earl E., 1929-
II. Series: United States. Geological Survey. Professional Paper
1213.

QE691.S78

551.7'8'097941

81-17864
AACR2

CONTENTS

	Page		Page
Introduction, by Earl E. Brabb	1	Stratigraphy—Continued	
Regional setting	2	Devils Den area	41
Paleogene benthic foraminiferal biostratigraphy and		Mount Diablo area	42
paleobathymetry of the central Coast Ranges of		San Jose area	42
California, by W. A. Berggren and Jane Aubert	4	Sveadal area	44
Abstract	4	Age	44
Introduction	5	Systematic paleontology	45
Sample material	5	Locality descriptions	47
Lodo Gulch area	5	San Jose area	47
Foraminiferal biostratigraphy	6	Sveadal area	48
Paleobathymetry	12	Mount Diablo area	48
Devils Den (aqueduct section) area	13	Devils Den area	48
Foraminiferal biostratigraphy	13	Eocene to Miocene calcareous plankton from the Santa Cruz	
Paleobathymetry	13	Mountains and northern Santa Lucia Range,	
Santa Cruz Mountains	13	California, by Richard Z. Poore and David Bukry	48
Smith Grade-Empire Grade area	13	Abstract	48
Lompico area	19	Introduction	49
Summary	19	Año Nuevo section	49
Lower Tertiary nannoplankton biostratigraphy in the central		Zayante Creek section	51
Coast Ranges of California, by A. D. Warren	22	San Lorenzo River section	52
Abstract	22	Church Creek area	53
Introduction	22	Discussion	58
Devils Den (aqueduct) area	22	Taxonomic notes	59
Lodo Gulch area	29	Calcareous nannofossils	60
Media Agua Creek area	32	Locality data	60
Comparison of the three areas	33	Reference to illustrations of identified coccolith species	60
The Paleogene of California—summary	36	Upper Eocene to lower Miocene benthic foraminifers from	
Calcareous nannofossil biostratigraphy of the Paleogene of		the Santa Cruz Mountains area, California,	
the Santa Cruz Mountains, California, by Bilal U. Haq	38	by Kristin McDougall	61
Definition of the Lodo Formation, by Earl E. Brabb	39	Abstract	61
Abstract	39	Introduction	61
Discussion	39	San Lorenzo River section	62
Large foraminifers of Eocene age from the Coast Ranges		Zayante Creek section	69
of California, by Alphonse Blondeau and E. E. Brabb	41	Año Nuevo section	70
Abstract	41	Summary	72
Introduction	41	Reference list of taxa	74
Stratigraphy	41	References cited	78
		Index	83

ILLUSTRATIONS

[Plates follow index]

- PLATE 1-5. Benthic foraminifers from the California Coast Ranges
6. Large foraminifers from California
 7. Planktic foraminifers from San Lorenzo Formation
 8. Planktic foraminifers from Año Nuevo and San Lorenzo River sections
 9. Calcareous nannofossils from the Santa Cruz Mountains
 - 10-17. Benthic foraminifers from Santa Cruz Mountains

	Page
FIGURE 1. Map showing natural provinces of California and areas sampled	2
2. Generalized paleogeographic map of California during early Tertiary	3
3. Map showing provinces of California and localities where microfossils were collected	6
4-8. Diagrams showing:	
4. Foraminiferal biostratigraphy of Lodo Formation at its type of locality in Lodo Gulch area	8
5. Foraminiferal biostratigraphy in Devils Den (aqueduct section) area	14
6. Foraminiferal biostratigraphy of Locatelli Formation, Smith Grade-Empire Grade area	18
7. Foraminiferal biostratigraphy of Butano Sandstone in Lompico area	20
8. Late Paleocene and early Eocene bathymetric history of Lodo Gulch and Devils Den areas	21
9. Index map showing nannofossil localities	22
10-15. Diagrams showing:	
10. Relation between nannoplankton and planktic foraminiferal zones	23
11. Biostratigraphy of aqueduct section, Devils Den area	26
12. Biostratigraphy of Lodo Gulch area	32
13. Biostratigraphy of Media Agua Creek area	34
14. Correlations of benthic foraminiferal stages to nannoplankton zones and subzones, Devils Den, Lodo Gulch, and Media Agua Creek areas	35
15. Correlation of Paleogene provincial benthic foraminiferal stages and calcareous microplankton zonations	37
16. Geologic map showing Lodo Gulch sections	40
17. Map of San Francisco Bay region showing localities of large foraminifers	43
18. Geologic maps of southern San Jose area showing fossil localities	44
19. Geologic map and stratigraphic column for Sveadal area	45
20. Index map showing location of sections	49
21. Generalized stratigraphic column of Año Nuevo section	50
22. Diagram showing ranges of calcareous plankton in Año Nuevo section	52
23. Stratigraphic column for north limb of Zayante Creek section	54
24. Stratigraphic column for San Lorenzo River section	55
25. Diagram showing ranges of plankton and zone assignments for San Lorenzo River section	57
26. Diagram showing lithostratigraphy and biostratigraphy of San Lorenzo River section	58
27. Map showing areas examined for benthic foraminifers	62
28-34. Diagrams showing:	
28. Stratigraphic relation of samples and benthic foraminiferal stages	63
29. Age interpretations of San Lorenzo River section	66
30. Stratigraphic distribution of diagnostic benthic foraminifers in San Lorenzo section	67
31. Stratigraphic distribution of diagnostic benthic foraminifers in Zayante Creek section	70
32. Stratigraphic distribution of diagnostic benthic foraminifers in the Año Nuevo section	70
33. Correlation of measured sections with benthic foraminiferal zones and stages	72
34. Ranges of benthic foraminiferal species	73

TABLES

	Page
TABLE 1. Benthic foraminifers in the type of Lodo Formation, Fresno County, California	10
2. Benthic foraminifers in the Devils Den (aqueduct) area	16
3. Nannoplankton from Devils Den and Lodo Gulch	30
4. Nannoplankton from the Paleogene of the Santa Cruz Mountains	38
5. Large foraminifers from the California Coast Ranges	42
6. Coccoliths and resultant zone assignments for the Año Nuevo section	51
7. Planktic foraminifers in samples from Año Nuevo section	51
8. Coccoliths and zone assignments for the Zayante Creek section	53
9. Coccoliths and zone assignments for the San Lorenzo River section	56
10. Planktic foraminifers and zone assignments for samples in the San Lorenzo River and Kings Creek sections	59
11. Planktic foraminifers and zone assignments for Church Creek Formation, Santa Lucia Range	59
12. Benthic foraminifers from the San Lorenzo River section	64
13. Key to field numbers and sample numbers	67
14. Benthic foraminifers from Zayante Creek section	69
15. Benthic foraminifers from the Año Nuevo area	71

STUDIES IN TERTIARY STRATIGRAPHY OF THE CALIFORNIA COAST RANGES

INTRODUCTION

By EARL E. BRABB

The correlation of rocks of Paleogene age in California with those in Europe has had a long and complex history that can only be highlighted here. Kleinpell (1938, p. 168-181), in his classic work defining Miocene benthic¹ foraminiferal stages of California, attempted to correlate faunas of California with those of western Europe and elsewhere. He pointed out that rocks usually considered lower Miocene in California are probably correlative with those considered Oligocene in Europe. Schenck and Childs (1942) correlated the Vaqueros Sandstone of California with the middle and upper Oligocene of Europe and elsewhere, based on the stratigraphic occurrence of *Lepidocyclina*. Their correlation was discussed extensively by a number of paleontologists in the same report. Some favored a Miocene age based on similarities of the Vaqueros molluscan fauna with the Burdigalian faunas of Europe, and on the apparent Miocene age of vertebrates from beds below the Vaqueros. Others argued that some of the mollusks in the Vaqueros were similar to those considered Oligocene in Trinidad, and that the percentage of living mollusks in the Vaqueros Sandstone (1-2 percent) was most similar to faunas of Oligocene age. These differences in opinion proved to be irreconcilable in the attempt by the U.S. Committee on Stratigraphy to provide a standard correlation chart for marine Cenozoic formations of western North America (Weaver and others, 1944). Two standards had to be provided, one based largely on mollusks, echinoids, and corals, and the other on benthic foraminifers. This dual classification has persisted almost to the present.

¹In this report, *benthic* refers to the seafloor habitat, and *benthonic* refers to the foraminiferal stages based on benthic organisms. Likewise, *planktic* organisms, that is, free floaters and weak swimmers, are distinguished from the *planktonic* zones based on plankton.

The validity of some California Tertiary stages based largely on benthic foraminifers has been questioned in recent years. Pierce (1972) and Barron (1976) believed that the Delmontian Stage of Kleinpell (1938), for example, is coeval with the lower and middle part of his Mohnian Stage. Steineck and Gibson (1971), Gibson and Steineck (1972), Schmidt (1975), Bandy (1972) and Bukry, Brabb, and Vedder (1977) stated their belief that probably all of the California Paleocene and Eocene stages of Mallory (1959) are time-transgressive when compared to nannoplankton and planktonic foraminifer zonations. Hornaday and Philips (1972), on the other hand, challenged some of these opinions.

In order to further the study of the relation between California Paleogene stages based on benthic foraminifers with zonations based on planktic foraminifers and nannoplankton, nine paleontologists were invited to examine the faunas from several Coast Range sections measured by Brabb, Clark, and Throckmorton (1977). The results of their investigation were presented orally at a meeting of the International Subcommission on Paleogene Stratigraphy in Menlo Park, California on October 28, 1977. The talks were preceded by three days of field trips to the measured sections and were followed by a microscope workshop to debate the identification and age of the various faunas. The paleontologists were then encouraged to submit papers for this volume, and all have graciously complied.

Acknowledgments.—Professor Ch. Pomerol of the University of Paris, currently chairman of the International Subcommission on Paleogene Stratigraphy, provided the leadership and inspiration for a series of biostratigraphic meetings that began in Paris in 1968, continued in Germany in 1969 and in the Caribbean in 1973, and culminated in the 1977 California meeting.

REGIONAL SETTING

The Paleogene faunas described in this report are from the California Coast Ranges (fig. 1). Most are from areas west of the San Andreas fault in the Santa Cruz Mountains of west-central California; a few are from areas east of the San Andreas fault from Mount Diablo to Devils Den. The Paleogene rocks

west of the San Andreas fault rest either directly on crystalline metasedimentary and granitic rocks of the Salinian block or on sedimentary strata of Cretaceous age. East of the San Andreas fault, the Paleogene rocks commonly rest on Jurassic and Cretaceous strata of the Great Valley sequence, which in turn are faulted against a complex tectonic assemblage of Mesozoic sedimentary, metamorphic, and




FIGURE 1.—Natural provinces of California, major geologic units and general location of areas sampled (modified from Hinds, 1952). MD, Mount Diablo; SJ, San Jose; SCM, Santa Cruz Mountains; LG, Lodo Gulch; DD, Devils Den.



FIGURE 2.—Generalized restoration paleogeographic map of California during early Tertiary, based on restoration of 305 km of late Cenozoic right-lateral slip along the modern San Andreas fault (from Nilsen and Clark, 1975). The present location of the Garlock fault and the following geographic features are included for orientation purposes: SAC, Sacramento; SF, San Francisco; BAK, Bakersfield; MON, Monterey; LA, Los Angeles; SD, San Diego; DD, Devils Den; LG, Lodo Gulch; MD, Mount Diablo; SJ, San Jose; SCM, Santa Cruz Mountains.

igneous rocks that are commonly referred to by petroleum geologists as Franciscan "basement." The relation of Franciscan "basement" rocks to the granitic basement rocks along which they are juxtaposed by the San Andreas fault has long been one of the most intriguing and complex problems of California geology. A series of lectures edited by Nilsen (1977) gives a synopsis of the problems and some working hypotheses for the tectonic evolution of California during Mesozoic and Cenozoic time. The most significant concept for the purpose of this report is that the San Andreas and other northwest-trending strike-slip faults have displaced the Paleogene sequences hundreds of kilometers with respect to each other. The Santa Cruz Mountains, for example, were thought by Nilsen and Clark (1975) to have been opposite the Devils Den area, 250 km to the south, during the early Tertiary.

The paleogeography of central California during the Paleogene is not well established. Most published analyses are not based on a firm biostratigraphic framework, so that rocks of middle Eocene age in one area may be compared with rocks of late Eocene age in another. There is also a problem with displacement of the Paleogene sequences along secondary lateral faults. Although extensive lateral displacement along the San Andreas has been considered

true since the classic report by Hill and Dibblee (1953), many new lateral faults have been discovered that make paleogeographic reconstructions more complex. Nevertheless, much progress has been made, especially by a few geologists who have attempted to apply modern sedimentological concepts to understanding Paleogene depositional patterns. Nilsen (1977) and Dibblee (1977) provided extensive bibliographies listing many of these studies, and they also provided the synthesis that is summarized below.

A generalized paleogeographic map of California during the early Tertiary, based on about 300 km of late Cenozoic right-lateral slip (fig. 2), shows that during the Paleogene the region between Devils Den and Mount Diablo was characterized by uplands in the Sierra Nevada, a linear basin in the area of the Great Valley in which a westward-thickening sequence of marine strata was deposited, islands on the continental borderland that contributed sediments in local areas, and a trench and subduction zone in what is now the Pacific Ocean. The La Honda basin, where all October 1977 field trips took place, was part of a continental borderland, mostly at lower bathyal and abyssal depths with open connections to the ocean.

PALEOGENE BENTHIC FORAMINIFERAL BIOSTRATIGRAPHY AND PALEOBATHYMETRY OF THE CENTRAL RANGES OF CALIFORNIA

By W. A. BERGGREN¹ and JANE AUBERT²

ABSTRACT

Paleogene foraminiferal assemblages of California have been examined from sections in the Lodo Gulch area, Fresno County; the Devils Den (aqueduct) area, Kern County; and the Smith and Empire Grade and Lompico areas, Santa Cruz County.

Benthic foraminifera indicate that the Lodo Formation in its type area was deposited during a single sedimentary cycle (lasting about 10 million years) at depths ranging from lower neritic

(somewhat shallower than 200 m) near its base and in its upper part to middle and upper bathyal (about 600 m) in its lower part. The Gredal Shale Member of the Kreyenhagen Formation in the Devils Den (aqueduct) section contains a rich planktic foraminiferal fauna throughout and shows no evidence in its upper part of the neritic fauna present in the upper part of the type Lodo Formation. Middle bathyal depths (below 600 m) are suggested for the lower part of the Gredal and upper bathyal depths (above 600 m) for its upper part. The marked shallowing in these sections in Zone P9 is denoted by the local disappearance or sporadic occurrence of bathyal taxa (such as *Nuttallides truempyi*) with more extensive stratigraphic ranges elsewhere and coincides closely with a major eustatic sea-level fall recently delineated in seismic stratigraphy.

¹Woods Hole Oceanographic Institution, Woods Hole, Mass., and Brown University, Providence, R. I.

²Société Nationale ELF-Aquitaine (Production) Centre Micoulau, 6400 Pau, France

The faunal assemblages from the Santa Cruz Mountains sections consist of predominantly benthic agglutinated foraminifers (with minor calcareous benthic and planktic foraminifers) which are characteristic of flysch deposits formed at the distal margins of turbidite fans in water depths of 1-2 km. Whereas the calcareous benthic assemblages of the Lodo Formation and Gredal Shale Member in the Devils Den area contain numerous cosmopolitan elements, the agglutinated flysch faunas of these units and those in the Santa Cruz Mountains appear to be less cosmopolitan, that is, they do not appear to be closely related to Paleogene flysch faunas of the Carpathian Mountains of Europe or the North Atlantic.

INTRODUCTION

California contains one of the best developed and representative Paleogene sequences in North America. Numerous studies have been devoted to the description of the foraminiferal faunas in this region (Martin, 1943; Israelsky, 1951, 1955; Mallory, 1959, 1970; Sullivan, 1962; Schmidt, 1970, 1975; Stine-meyer, 1976).

In connection with the Field Conference on the Paleogene sponsored by the International Commission on Paleogene Stratigraphy (October 1977) we were requested to make a preliminary study of prepared samples from four sections that represent a cross section of the California early Paleogene. The sections included: (1) the Lodo Formation at its type locality in the Lodo Gulch area, Fresno County; (2) the Lodo Formation, Avenal Sandstone, and Gredal Shale and lower part of the Point of Rocks Sandstone Members of the Kreyenhagen Formation in the Devils Den (aqueduct section) area, Kern County; (3) the Locatelli Formation in the Smith Grade-Empire Grade area in the Santa Cruz Mountains, Santa Cruz County; and (4) the Butano Sandstone in the Lompico area in the Santa Cruz Mountains, Santa Cruz County (fig. 3). The last two areas were visited during the course of the field conference.

Within the context of this preliminary analysis we have not been able to recognize and apply the California Paleogene foraminiferal stages of Mallory (1959). Mallory's stages are not time-stratigraphic units at all in the sense prescribed by the American Code of Stratigraphic Nomenclature or the International Stratigraphic Guide. These stages are better characterized as loosely defined assemblages of benthic foraminifers (faunules) and, as such, of ecologic rather than chrono-stratigraphic significance. Indeed, the California provincial stages are demonstrably time-transgressive (Steineck and Gibson, 1971; Poore, 1976; Bukry and others, 1977; see also Warren, 1977).

Details of the stratigraphic sections from which our samples have been collected are presented in

Brabb, Clark, and Throckmorton (1977).

Acknowledgments.—This paper was presented, in considerably abbreviated form, at the Field Conference on the Paleogene of California held at Menlo Park, Calif. in late October-early November 1977.

We thank Earl E. Brabb and Joseph C. Clark (Indiana, Penn.) for providing the sample material upon which the investigation is based. Richard Z. Poore provided the Merle Israelsky foraminiferal collection from the type Lodo Formation on loan, and this proved extremely helpful to us. We thank Francis Saffon for the scanning electron micrographs prepared on the Cambridge Mark II instrument in the laboratories of Societe National Elf-Aquitaine (Production), Pau, France.

This investigation is part of a joint project with the Woods Hole Oceanographic Institution (sponsored by Chevron, Exxon, Marathon, Mobil, and Shell Oil Companies) and Societe National Elf-Aquitaine (Production) devoted to Cenozoic benthic foraminiferal biostratigraphy and ecology. We wish to thank B. U. Haq and R. C. Tjalsma of Woods Hole for discussions on Paleogene stratigraphy and for reviewing the manuscript.

SAMPLE MATERIALS

Nineteen samples, collected by Earl Brabb and prepared by the U.S. Geological Survey, have been examined from the type section of the Lodo Formation along Lodo Gulch (section TS, fig. 16 of Brabb, this volume). These samples were supplemented by a cursory examination of the Israelsky (1951) collection of 110 samples from two reference sections less than 1000 m south of Lodo Gulch. Fourteen samples collected by Brabb have been examined from the Lodo Formation and the Gredal Shale, and the Point of Rocks Sandstone Members of the Kreyenhagen Formation exposed in the Devils Den (aqueduct section) area, some 100 km southeast of the type Lodo.

Fourteen samples collected by Earl Brabb and Joseph C. Clark have been examined from the terrigenous arkosic sandstone and siltstone of the Locatelli Formation and Butano Sandstone in the Smith Grade-Empire Grade and Lompico areas of the Santa Cruz Mountains.

LODO GULCH AREA

The type section of the Lodo Formation at Lodo Gulch (fig. 4) is in the Tumey Hills, Fresno County, on the west side of the San Joaquin Valley. It is one of the most important reference sections for lower Paleogene stratigraphy of California and has been the subject of several studies on foraminifers (Martin,




FIGURE 3.—Physiographic provinces of California and localities where microfossil assemblages were collected. 1, Lodo Gulch; 2, Devils Den; 3, Smith Grade-Empire Grade area; 4, Lompico.

1943; Israelsky, 1951, 1955; Mallory, 1959; Schmidt, 1970) and calcareous nannoplankton (Bramlette and Sullivan, 1961). At its type section the Lodo is approximately 360 m thick and is of late Paleocene and early Eocene (planktic foraminiferal Zone P4-P9) age. It is unconformably underlain by the Cretaceous part of the Moreno Shale and is overlain by the Domengine Sandstone of middle Eocene age. Molluscan faunas collected near the base of the Lodo Formation some 700 m northwest of its type section are correlated by Smith (1975) with the Martinez Stage of the provincial molluscan standard of Weaver and others (1944), and with mollusks from the Bracheux Sandstone of Paleocene age in the northern Paris Basin. A potassium-argon date of 58.5 m.y. (Funnell, 1964, p. 188, item 113) from glauconite in the basal beds of the Lodo Formation provides a minimum age on the *Planorotalites pseudomenardii* and *Heliolithus riedeli* Zones, within which the glauconite is situated (Bramlette and Sullivan, 1961; Schmidt, 1970). Additional details on the geologic and stratigraphic setting of the type Lodo Formation may be found in Israelsky (1951) and Schmidt (1970), and its relation to other Paleogene sections in central California is discussed by Schmidt (1975).

FORAMINIFERAL BIOSTRATIGRAPHY

The type Lodo Formation has been zoned with the aid of planktic foraminifers by Schmidt (1970), and our examinations have essentially confirmed his findings, including the probable presence of a short hiatus (or disconformity) between the Paleocene and Eocene about 50 m above the base of the type section at the Lodo Gulch (fig. 4). We did not find definitive evidence for the presence of Zone P6. The boundary between Zones P8 and P9 has been determined on the basis of the initial appearance of *Subbotina inaequispira*, *S. frontosa*, *S. linaperta* and *Acarinina densa* over the interval of samples 70-72 (Israelsky) and 1231 (Brabb). The top of the Lodo Formation is considered to lie within an interval equivalent to Zone P9, as differentiation between Zones P9 and P10 is often difficult, and planktic foraminifers diminish sharply in the upper 150 m of the type Lodo Formation as a result of shallowing.

Over 150 species of calcareous and agglutinated benthic foraminifers have been recorded from the Lodo Gulch section and the nearby Devils Den section some 100 km southeast of Lodo Gulch. Species of the calcareous genera *Anomalinoidea*, *Bulimina*, *Trifarina*, *Cibicidoides*, and *Lenticulina* dominate these assemblages. Agglutinated taxa (including *Bathysiphon*, *Rhabdammina*, *Haplophragmoides*,

Cyclammina, *Karreriella*, *Dorothia*, *Silicosigmoilina*, and *Tritaxilina*) occur throughout both sections and at some levels constitute almost the entire foraminiferal assemblage.

The distribution of species identified in the Brabb samples in the type Lodo Formation is shown in Table 1. Generalized foraminiferal assemblages are shown in figure 4 based on the more common or characteristic elements occurring over a given stratigraphic interval. It should be borne in mind that these assemblages are based on an examination of only 18 samples collected by Brabb and the more closely spaced samples in the Israelsky collection. Characteristic features of the benthic foraminiferal fauna of the Lodo Formation include the following:

1. There is a sequential influx of new taxa throughout the section.
2. There is a marked faunal change (reduction or disappearance of taxa from below and replacement of taxa above) in the interval above 150-100 m below the contact of the Domengine Sandstone and the Lodo Formation (above Israelsky sample 70 and Brabb sample 1231). Planktic foraminifers (particularly the angulo-conical morozovellids) exhibit a marked reduction over this interval as well and become rarer toward the upper part of the Lodo Formation.
3. An essentially threefold faunal subdivision can be made (from bottom to top): (1) The lower 15 m (approx.) contains a fauna similar to that in the Midway Group, including *Cibicidoides alleni*, *Osangularia plummerae*, *Anomalinoidea danica rubiginosa*, and species of *Silicosigmoilina*, *Spiroplectammina*, *Clavulinoides*, and *Ammodiscus*. This interval corresponds essentially to Zone P4 and Israelsky samples 3-11. This interval is below the lowest sample (1251C) collected by Brabb, which is stratigraphically located between Israelsky samples 13 and 14. (2) The interval from about 15 m to 200 m, between Israelsky samples 12 to 70, contains an assemblage characterized or dominated by various bulminids, anomalinids, osangulariids, aragonids, *Nuttallides truempyi*, *Bathysiphon*, *Rhabdammina*, *Cyclammina*, *Silicosigmoilina*, and *Clavulinoides*. This interval corresponds essentially to Zones P5-P8. (3) The upper 175 m contains an assemblage characterized by various small cibicidids, anomalinids, bolivinids, *Florilus florinense*, *Uvigerina elongata*, *Eponides primus*, *Cyclammina*, and *Trochammina*. This interval presumably corresponds to Zone P9, although planktic foraminifers are sparse or rare over this interval.





FIGURE 4.—Foraminiferal biostratigraphy and paleobathymetry of the Lodo Formation at its type locality in the Lodo Gulch area. The upper part of the section was sampled (small numbers) by Israelsky (1951, pl. 1, section A) along the first gulch south of Lodo Gulch. Samples collected by E. E. Brabb (74CB1221B and others) are projected into this section from the type section of the Lodo Formation along Lodo Gulch. (section TS). The lower part of the section was measured (small

BENTHIC FORAMINIFERAL ASSEMBLAGE		PALEOBATHYMETRY, IN METERS									PLANKTONIC FORAMINIFERAL ZONE	EUROPEAN STAGE	WEST COAST STAGES (MALLORY, 1959)	SERIES							
		NERITIC			BATHYAL																
		U	M	L	UPPER		LOWER														
AGGLUTINATED		CALCAREOUS		1	2	3	4	5	6	7	8	9									
BATHYSIPHON	Cyclammina	Anomalinoides acutus Osangularia tenuicarinata Eponides primus Boliuina crenulata Bulimina whitei Florilus florinense Uuigerina elongata																			
	Trochammina	Eponides primus Bulimina whitei Boliuina crenulata																			
	Clavulinoides californicus	Florilus florinense Hoeglundina eocenica Anomalinoides acutus Bulimina whitei Boliuina crenulata Cibicidoides felix Spiroloculina lamposa																			
		Siphonina wilcoxensis Anomalinoides acutus Osangularia tenuicarinata Spiroloculina lamposa																			
		Osangularia tenuicarinata																			
		Aragonia aragonensis																			
	Verneuilina triangulata	Loxostomoides applinae Spiroloculina lamposa																			
	Spiroplectammina richardi	Bulimina callahani Bulimina trinitatensis Anomalinoides aragonensis Cibicidoides fortunatus																			
		Aragonia aragonensis Loxostomoides applini Osangularia tenuicarinata Trifarina californica Spiroloculina lamposa																			
		Nuttallides truempyi Osangularia tenuicarinata Anomalinoides acutus Trifarina californica Hoeglundina eocenica																			
	Gaudryina coalingensis	Pleurostomella paleocenica Trifarina californica																			
	Clavulinoides californicus Gaudryina coalingensis Cyclammina simiensis Ammodiscus glabratus Silicosigmoilina californica	Anomalinoides danica rubiginosa Osangularia plummerae Eponides beisseli																			
	Silicosigmoilina californica Ammodiscus glabratus	Cibicidoides allenii																			
Spiroplectammina sp. Glomospira charoides Ammodiscus glabratus Cyclammina simiensis																					
Dorothia lodoensis Clavulinoides californicus Silicosigmoilina californica Spiroplectammina sp.																					

sample numbers) by Israelsky (1951, pl. 1, section B) along a small gulch about 300 m south of the mouth of Lodo Gulch. Samples collected by E. E. Brabb (74CB1251D and others) in this section are projected from still another section about 700 m north of Lodo Gulch. (section C). The location of all these sections is shown on figure 16 of Brabb (this volume).

TABLE 1.—Benthic foraminifers in the type Lodo Formation, Lodo Gulch, Fresno County, California

E. Brabb samples																		Benthic foraminifers	
1251C	1251D	1231G	1231F	1231E	1221J	1231D	1231C	1221H	1231B	1221G	1231A	1231	1221E	1222A	1221B	1222	1221	1214B	
																		<i>Bifarina vicksburgensis</i> (CUSHMAN) <i>Gaudryina jacksonensis</i> var. <i>coalingensis</i> MALLORY <i>Epistomina partschiana</i> (d'ORBIGNY) <i>Eponides</i> sp. <i>Lenticulina</i> cf. <i>L. terryi</i> (CORRYEL & EMBICH) <i>Lenticulina theta</i> COLE <i>Cibicidoides</i> cf. <i>C. beatus</i> (MARTIN) <i>Anomalinoides keeni</i> (MARTIN) <i>Angulogerina</i> aff. <i>A. cuneata</i> BROTZEN <i>Lenticulina</i> cf. <i>L. gyroscalprum</i> (STACHE) <i>Trifarina advena</i> var. <i>californica</i> MALLORY <i>Vaginulinopsis mexicana</i> var. <i>nudicosta</i> CUSHMAN & HANNA <i>Lenticulina pseudovortex</i> (COLE) <i>Lenticulina pseudocultrata</i> (COLE) <i>Lenticulina weaveri</i> ? (BECK) <i>Lenticulina alatolimbatus</i> (GUMBEL) <i>Lenticulina rotulata</i> (LAMARCK) <i>Lenticulina</i> spp. <i>Nodosaria deliciae</i> MARTIN <i>Dentalina</i> spp. <i>Rectoglandulina</i> sp. <i>Marginulina subbullata</i> HANTKEN <i>Gaudryina macrocamerata</i> MALLORY <i>Verneuilina triangulata</i> COOK <i>Alabamina</i> sp. <i>Anomalinoides acutus</i> (PLUMMER) <i>Cibicidoides whitei</i> (MARTIN) <i>Lenticulina vortex</i> (FICHTEL & MOLL) <i>Vulvulina curta</i> CUSHMAN & SIEGFUS <i>Rhabdammina eocenica</i> CUSHMAN & HANNA <i>Oridorsalis umbonatus</i> (REUSS) <i>Pullenia eocenica</i> CUSHMAN & SIEGFUS <i>Chilostomelloides cylindroides</i> REUSS <i>Cibicidoides fortunatus</i> (MARTIN) <i>Nodosaria arundinea</i> SCHWAGER <i>Nodosaria latejugata</i> GUMBEL <i>Nuttallides truempyi</i> (NUTTALL) <i>Vaginulinopsis</i> cf. <i>kelleyi</i> MARTIN <i>Gaudryina</i> cf. <i>G. laeyigata</i> FRANKE <i>Clavulinoides californicus</i> MALLORY <i>Spiroplectammina richardi</i> MARTIN <i>Anomalinoides welleri</i> (PLUMMER) <i>Dorothia excentrica</i> ISRAELSKI <i>Cibicidoides eponidiformis</i> (MARTIN) <i>Bulimina ovata</i> d'ORBIGNY <i>Lenticulina midwayensis</i> (in MALLORY) <i>Plectofrondicularia kerni</i> COOK <i>Anomalinoides aragonensis</i> (NUTTALL) <i>Anomalinoides judas</i> (MARTIN) <i>Bulimina curtissima</i> CUSHMAN & SIEGFUS <i>Bulimina impendens</i> PARKER & BERMUDEZ <i>Bulimina macilenta</i> CUSHMAN & PARKER <i>Bulimina trinitatis</i> CUSHMAN <i>Globobulimina</i> sp. <i>Valvulineria childsi</i> (MARTIN) <i>Dentalina subtrigata</i> (STACHE) <i>Dentalina spinosa</i> d'ORBIGNY <i>Haplophragmoides</i> cf. <i>H. nonionelloides</i> ISRAELSKY <i>Haplophragmoides</i> spp. <i>Ammodiscus</i> cf. <i>A. pennyi</i> CUSHMAN & JARVIS <i>Bulimina callahani</i> GALLOWAY & MORREY <i>Dentalina basiplanata</i> CUSHMAN <i>Buliminella grata</i> var. <i>convoluta</i> MALLORY <i>Turritilina brevispira</i> TEN DAM <i>Gyroidina soldanii</i> var. <i>octocamerata</i> CUSHMAN & HANNA <i>Stilostomella plummerae</i> (CUSHMAN) <i>Plectofrondicularia whitei</i> MARTIN <i>Marginulina curvatura</i> (CUSHMAN) <i>Bathysiphon</i> sp. B. ISRAELSKY <i>Nodosaria ewaldi</i> REUSS <i>Angulogerina wilcoxensis</i> (CUSHMAN & GARRETT) <i>Buliminella bradburyi</i> (MARTIN) <i>Bulimina</i> sp. (~ <i>instabilis</i> CUSHMAN & PARKER) <i>Haplophragmoides</i> cf. <i>H. coalingensis</i> CUSHMAN & HANNA <i>Silicosigmollina californica</i> CUSHMAN & CHURCH <i>Nodosaria</i> ? <i>gyrata</i> MALLORY	
P5																			
P8																			
P9																			
Planktic foraminiferal biozonation																			

E. Brabb samples										Benthic foraminifers									
1251C	1251D	1231G	1231F	1231E	1221J	1231D	1231C	1221H	1231B	1221G	1231A	1231	1221E	1222A	1221B	1222	1221	1214B	<i>Dentalina approximata</i> REUSS <i>Goesella</i> sp. (ISRAELSKY) <i>Bathysiphon eocenicus</i> CUSHMAN & HANNA <i>Cyclammina simiensis</i> BERRY <i>Eggerella</i> sp. (in MALLORY 1959) <i>Haplophragmoides excavatus</i> CUSHMAN & WATERS <i>Loxostomoides applinae</i> (PLUMMER) <i>Uvigerina alabamensis</i> CUSHMAN & GARRETT <i>Uvigerina elongata</i> COLE <i>Lenticulina turbinata</i> (PLUMMER) <i>Anomalinoidea crassiseptus</i> (CUSHMAN & SIEGFUS) <i>Bulimina whitei</i> MARTIN <i>Bulimina debilis</i> MARTIN <i>Cibicides spiropunctatus</i> GALLOWAY & MORREY <i>Vaginulinopsis asperuliformis</i> (NUTTALL) <i>Valvulineria allomorphinoides</i> (REUSS) <i>Pleurostomella</i> cf. <i>P. paleocenica</i> CUSHMAN <i>Siphonina wilcoxensis</i> CUSHMAN <i>Fursenkoina zetina</i> (COLE) var. <i>indirecta</i> (MALLORY) <i>Asterigerina crassaformis</i> CUSHMAN & SIEGFUS <i>Bolivina explicata</i> var. <i>californica</i> MALLORY <i>Uvigerina lodoensis</i> var. <i>miriamae</i> MALLORY <i>Osangularia tenuicarinata</i> CUSHMAN & SIEGFUS <i>Aragonia aragonensis</i> (NUTTALL) <i>Glandulina</i> sp. <i>Valvulineria</i> sp. (MALLORY 1959) <i>Spiroplectammina tejonensis</i> MALLORY <i>Bifarina eleganta</i> (PLUMMER) <i>Karrerina fallax</i> RZEHA <i>Lagena crowleri</i> MARTIN <i>Cibicoides beatus</i> (MARTIN) <i>Bolivina</i> sp. <i>Eponides lodoensis</i> MARTIN <i>Nodosaria affinis</i> REUSS <i>Marginulina</i> cf. <i>M. gardnerae</i> (PLUMMER) <i>Cassidulina globosa</i> HANTKEN <i>Cibicides "cocoaensis"</i> CUSHMAN <i>Lenticulina inornata</i> (d'ORBIGNY) <i>Dorothia excentrica</i> ISRAELSKY <i>Fronicularia naheolensis</i> CUSHMAN & TODD <i>Pleurostomella</i> cf. <i>P. acuta</i> HANTKEN <i>Hoeglundina</i> sp. <i>Vaginulinopsis mexicana</i> var. <i>kerni</i> COOK <i>Planularia</i> cf. <i>P. truncana</i> GUMBEL <i>Lenticulina rosetta</i> (GUMBEL) <i>Dentalina communis</i> (d'ORBIGNY) <i>Dentalina wilcoxensis</i> CUSHMAN <i>Marginulina glabra</i> d'ORBIGNY <i>Marginulina sischoae</i> MALLORY <i>Cibicides martinezensis</i> CUSHMAN & BARKSDALE <i>Dentalina jacksonensis</i> CUSHMAN & APPLIN <i>Amphimorphina ignota</i> CUSHMAN & SIEGFUS <i>Dentalina</i> cf. <i>D. multilineata</i> BORNEMANN <i>Plectofronicularia minuta</i> ? SULLIVAN <i>Nodogenerina lepidula</i> (SCHWAGER) <i>Nodogenerina kressenbergensis</i> (GUMBEL) <i>Marginulina adunca</i> (COSTA) <i>Gyroidina orbicularis</i> d'ORBIGNY var. <i>planata</i> CUSHMAN <i>Bifarina nuttalli</i> CUSHMAN & SIEGFUS <i>Dorothia</i> sp. <i>Martinottiella eocenica</i> CUSHMAN & BERMUDEZ <i>Cibicides felix</i> MARTIN <i>Bulimina corrugata</i> CUSHMAN & SIEGFUS <i>Lagena acuticosta</i> REUSS <i>Tritaxilina colei</i> CUSHMAN & SIEGFUS <i>Florilus florinensis</i> (COLE) <i>Pleurostomella gredalensis</i> COOK <i>Siphonina jacksonensis</i> CUSHMAN <i>Spiroloculina lamposa</i> HUSSEY <i>Cibicides</i> aff. <i>c. praecursorius</i> SCHWAGER <i>Karrerella</i> cf. <i>K. elongata</i> COLE <i>Eggerella</i> sp. <i>Lenticulina</i> cf. <i>L. antipodum</i> STACHE <i>Eponides primus</i> MARTIN <i>Cibicides kernensis</i> COOK
EXPLANATION █ Rare █ Frequent █ Common Planktic foraminiferal biozonation																			

PALEOBATHYMETRY

Strict interpretation of Paleogene paleobathymetry based on analogy with present-day faunal counterparts is unreliable in view of demonstrable differences in distributional patterns (Lohmann and Tjalsma, 1975; Tjalsma and Lohmann, 1975) and water mass structure (Douglas and Savin, 1973; Boersma and Shackleton, 1977). Nevertheless, local and regional geologic information, distribution patterns of taxa with geophysically controlled paleobathymetry (Sclater and others, 1971; Berger, 1973), and distribution of associated faunal elements with a presumed stable ecology can be combined to allow a reasonable paleodepth estimate of special faunal associations.

We interpret the paleobathymetric history of the type Lodo Formation in terms of a single sedimentary cycle in which the middle part of the formation (corresponding to Zones P5 to P8) was deposited at middle to upper bathyal depths (about 600 m), whereas the lower (Zone P4) and upper (Zone P9) parts were deposited near the neritic-bathyal boundary (above 200 m). The following line of reasoning has been used.

1. Faunas from the Midway Group have been shown (Berggren and Aubert, 1975) to represent a cosmopolitan fauna formed at neritic depths. An association of some Midway elements with somewhat deeper water elements is seen at depths interpreted as upper-middle bathyal in Tunisia (Aubert and Berggren, 1975) and Bavaria (Hillebrandt, 1962), but these elements are not observed in the lower (Zone P4) part of the type Lodo Formation.
2. In the succeeding 150 m (Israel'sky samples 13 to 63; Brabb samples 1251C to 1231C, fig. 4) a number of forms appear that are restricted to this interval but which have longer stratigraphic ranges in lower bathyal-abyssal deposits in the ocean (R. C. Tjalsma, oral commun., 1977). These include *Bulimina trinitatensis*, *Nuttallides truempyi*, *Anomalinoides aragonensis*, *Buliminella grata*, *Bulimina callahani* (= *orphanensis*), *Anomalina crassisepta*, and *Asterigerina crassaformis*. In addition several species of the agglutinated genera *Rhabdammina*, *Bathysiphon*, and *Cyclamina*, normally associated with bathyal deposits, appear within this interval as well. For example, *Nuttallides truempyi* is one of the dominant lower bathyal-abyssal taxa with an age range of Late Cretaceous (Maestrichtian) to latest Eocene (Laughton and others, 1972; Douglas, 1973; R. C. Tjalsma, oral commun., 1977). Its

upper depth limit is not known with certainty, but it occurs in the late Eocene at DSDP (Deep Sea Drilling Project) Site 116 in middle bathyal depths (above 1000 m; Laughton and others, 1972). It does not occur in neritic (Midway type) faunas, and thus we suggest an upper depth limit of 500–600 m for this taxon, between upper and middle bathyal depths. *Bulimina callahani* (= *orphanensis*) has a stratigraphic range from Zones P6a to P8 (R. C. Tjalsma, oral commun., 1977; Berggren and Aubert, 1976) in oceanic sediments yet is essentially restricted to the upper half of Zone P8 in the type Lodo Formation. At Orphan Knoll (DSDP Site 111), *B. callahani* (= *orphanensis*) occurs in the lower Eocene with faunas from depths of more than 1 km. As with *Nuttallides truempyi*, its upper depth limit is suggested to lie near the upper-middle bathyal boundary (500–600 m). Upward shallowing is suggested by the nearly simultaneous local termination or disappearance of these taxa in the middle part of the section about 190 m above the base (175 m below the top) of the Lodo Formation at a level that corresponds closely to the planktic foraminiferal Zone P8–P9 boundary (characterized by the initial appearance of *Subbotina frontosa* = *S. boweri*, *S. linaperta*, *S. inaequispira*, and *Acarinina densa*) and their replacement by shallower water forms (cibicides, bolivinids, eponidids, *Florilus*, *Valvulineria*, and hispid uvigerinids of the *elongata* group characteristic of neritic depths: 50–200 m; Boersma, 1974).

3. A preliminary examination of the ostracode fauna in the Israel'sky collections by R. H. Benson (Smithsonian Institution, written commun., 1977) indicates the presence of *Trachyleberidea* and *Cytherella*, both typical of bathyal deposits, *Argilloecia*, and some allochthonous specimen of *Loxoconcha*. *Krithe* (with markedly open vestibule) suggests that oxygen ranged between 4.5 and 5.5 ml/l. Very small eye tubercles present on an *Actinocythereis*-like form suggest (to Benson) water depths not in excess of 600–800 m, in good accord with estimates made above.

The presence of cyclaminids and *Bathysiphon* to the top of the type Lodo Formation at depths interpreted from other faunal evidence to have been near the shelf-slope break (about 200 m) suggest that the upper depth limits of these agglutinated benthic foraminiferal faunas were somewhat shallower during the Paleogene than during the Neogene and present day, perhaps because of the lack of a pronounced thermocline during this time.

DEVILS DEN (AQUEDUCT SECTION AREA)

The aqueduct section in the Devils Den area is located in Kern County some 100 km southeast of Lodo Gulch (fig. 3). Several hundred meters of Cretaceous, Paleocene, and Eocene strata are exposed there (Brabb and others, 1977, fig. 37), of which only a part (fig. 5) has been selected for study. Our preliminary examination of the planktic foraminiferal fauna suggests that the sequence studied spans Zones P5 to P9 and may include P10. Zones P5 to P6 are represented only in a fault-bounded part of the Lodo Formation.

Warren (1977) reported the presence of the calcareous nannoplankton *Tribrachiatus orthostylus* (NP12) to *Nannotetrina quadrata* (NP15) Zones in the aqueduct section. The presence of the *Discoaster lodoensis* (NP13) and *D. sublodoensis* (NP14) Zones in the lower part of the Gredal Shale Member of the Kreyenhagen Formation supports our assignment of this same interval to planktic foraminiferal Zones P8 and P9 (see also Wise and Constans, 1976). The upper 40-50 m of the Gredal Shale Member was placed by Warren in the *Nannotetrina quadrata* (NP15) Zone of middle Eocene age.

In summary, the sequence studied in the aqueduct section of Devils Den is correlative with most of the type Lodo Formation.

Planktic foraminifers (particularly acarininids and anguliconical morozovellids) are abundant and well preserved throughout the aqueduct section, in contrast to those in the Lodo Gulch area.

FORAMINIFERAL BIOSTRATIGRAPHY

The benthic foraminiferal taxa recorded from the Devils Den (aqueduct section) area (table 2), form the basis for the assemblage subdivision shown in figure 5. The basic features of the assemblages that should be noted are as follows:

1. Faunal elements similar to those from the Midway Group are absent at the base of the sequence studied, that is, in the lowest beds of the Lodo Formation, whereas the *Florilus-Eponides-Uvigerina* assemblage is absent at the top.
2. The assemblages are nearly identical with those present in the middle part of the type Lodo Formation at Lodo Gulch (*Nuttallides truempyi*, buliminids, osangulariids, anomalinids, cyclaminids, *Tritaxilina*, *Rhabdammina*, *Bathysiphon*).
3. Several of the taxa that disappear near the boundary between Zones P8 and P9 in the Lodo Gulch area appear to range well into Zone P9 in the

Devils Den (aqueduct section) area (*N. truempyi*, *Anomalinoides crassiseptus*, *Asterigerina crassaformis*).

4. The upper part of the aqueduct section contains a fauna composed primarily of elements that range through a major part of the section: *Rhabdammina eocenica*, *Bathysiphon eocenicus*, *Cyclamina simiensis*, *Haplophragmoides* sp. cf. *H. eggeri*, *H. excavatus*, *Tritaxilina colei*, *T. principiensis*, *Karrerella mediaguaensis*, *Plectina cubensis*, *Anomalinoides crassiseptus*, *Cibicidoides fortunatus*, *Lenticulina* spp.

PALEOBATHYMETRY

The similarity of faunas in the aqueduct section with those in the Lodo Gulch area suggests similar water depths. The fact that some of the bathyal taxa (*Nuttallides truempyi*) range into Zone NP14 (equivalent to P9) and some (*Anomalinoides crassiseptus*, *Asterigerina crassaformis*) as high as NP15 (equivalent to P10) suggests that water depths were greater in the Devils Den area than in the Lodo Gulch area and that the shallowing near the boundary between Zones P8 and P9 in the Lodo Gulch area occurs within Zone P9 in the Devils Den area. The shallowing eliminated some of the taxa. Other taxa, however, persisted because the water depth was still well within their upper depth limit. It should be noted that *Bulimina callahani* (= *Bulimina orphanensis*) disappears near the boundary between Zones P8 and P9 in the Devils Den area as in the Lodo Gulch area, an apparent example of an evolutionary extinction as opposed to bathymetrically controlled local disappearance. A middle bathyal depth (below 600 m) is suggested for the Lodo Formation and lower part of the Gredal Shale Member in the Devils Den section, an upper bathyal depth (above 600 m) for the upper 40-50 m of the Gredal Shale Member, above the disappearance of *N. truempyi*.

SANTA CRUZ MOUNTAINS

Fourteen samples have been examined from the terrigenous arkosic sandstone and siltstone of the Locatelli Formation and the Butano Sandstone in the Smith Grade-Empire Grade and Lompico areas, respectively, of the Santa Cruz Mountains.

SMITH GRADE-EMPIRE GRADE AREA

The seven samples examined from the Locatelli Formation (Paleocene) span a stratigraphic interval of approximately 250 m (fig. 6).

BENTHIC FORAMINIFERAL ASSEMBLAGES		PALEOBATHYMETRY									PLANKTONIC FORAMINIFERAL ZONE	BENTHONIC FORAMINIFERAL STAGES OF MALLORY (1959) PROVISIONAL STAGE DETERMINATION BY A.D. WARREN (THIS VOLUME)	EUROPEAN STAGE	SERIES OR SUBSERIES
		NERITIC			BATHYAL									
		U	L	M	UPPER			MIDDLE						
AGGLUTINATED	CALCAREOUS	1	2	3	4	5	6	7	8	9				
													No age data	
EOCENICA	<i>Spiroplectammina directa</i> <i>Cyclammina simiensis</i> <i>Cyclammina clarki</i> <i>Tritaxilina colei</i> <i>Plectina cubensis</i> <i>Trochammina</i> spp.										P10(?)	Narizian	Ypresian or Lutetian	Lower or middle Eocene
	<i>Cyclammina simiensis</i> <i>Cyclammina incisa</i> <i>Spiroplectammina richardi</i>										P9	Narizian or Ulatisian		Lower and(or) middle Eocene
RHABDAMMINA												Ulatisian		Lower Eocene
	<i>Clavulinoides californicus</i> <i>Spiroplectammina richardi</i> <i>Tritaxilina colei</i> <i>Tritaxilina principiensis</i>										P8	Penutian or Ulatisian		
	<i>Clavulinoides californicus</i> <i>Tritaxilina colei</i> <i>Karriella media aquaensis</i> <i>Vulvulina curta</i>										P5 and(or) P6	Penutian		
	<i>Anomalinoides aragonensis</i> <i>Osangularia tenuicarinata</i> <i>Trifarina californica</i>											Bulitian and Ynezian	Thanetian	Paleocene
BY FAULTING														
EOCENICA	<i>Cyclammina simiensis</i> <i>Cyclammina incisa</i> <i>Verneuilina triangulata</i> <i>Spiroplectammina richardi</i> <i>Tritaxilina colei</i> <i>Gaudryina laevigata</i> <i>Clavulinoides californicus</i> <i>Karriella media aquaensis</i> <i>Vulvulina curta</i>										P9 and(or) P8	Penutian Ulatisian	Ypresian	Lower Eocene
RHABDAMMINA	<i>Cibicidoides fortunatus</i> <i>Cibicidoides martinezensis</i> <i>Cibicidoides whitei</i> <i>Osangularia mexicana</i> <i>Buliminella grata</i> var. <i>convoluta</i> <i>Nuttallides truempyi</i> <i>Oridorsalis umbonatus</i> <i>Bulimina trinitatensis</i> <i>Aragonia aragonensis</i> <i>Valvulinella childsi</i>													

FIGURE 5.— Continued.

STUDIES IN TERTIARY STRATIGRAPHY, CALIFORNIA COAST RANGES

TABLE 2.—Benthic foraminifers in the Devils Den (aqueduct) area, Kern County, California—Continued

E. Brabb samples		Benthic foraminifers	
1272			
1273			
1273A			
1273B			
1281			
1281C			
1281D			
1633			
1281E			
1281H			
1281J			
1281K			
1281L			
1281M			
1291			
1231A			
		<i>Osangularia mexicana</i> (COLE) <i>Bulimina truncanella</i> FINLAY ? <i>Ammobaculites</i> sp. ? <i>Epistomina partschiana</i> (d'ORBIGNY) <i>Gaudryina coalingsensis</i> CUSHMAN & HANNA <i>Cyclammina incisa</i> STACHE <i>Dorothia cubensis</i> CUSHMAN & BERMUDEZ <i>Glomospira charoides</i> PARKER & JONES <i>Karrerella monumentensis</i> MALLORY <i>Dorothia germanica</i> CUSHMAN <i>Plectina cubensis</i> CUSHMAN & BERMUDEZ <i>Conotrochammina</i> spp. <i>Ramulina</i> cf. <i>R. navaroana</i> CUSHMAN <i>Nodogenerina kressenbergensis</i> GUMBEL <i>Nodogenerina lepidula</i> (SCHWAGER) <i>Lenticulina theta</i> COLE <i>Planularia truncana</i> GUMBEL <i>Baggatella californica</i> MALLORY <i>Bulimina semicostata</i> var. <i>lacrima</i> MALLORY <i>Bolivina kleinpell</i> BECK <i>Plectofrondicularia minuta</i> SULLIVAN <i>Bulimina impendens</i> PARKER <i>Bifarina vicksburgensis</i> CUSHMAN <i>Angulogerina</i> cf. <i>A. wilcoxensis</i> (CUSHMAN & PONTON) <i>Eponides minima</i> CUSHMAN <i>Gyroidina orbicularis</i> var. <i>obliquata</i> CUSHMAN & MASTER <i>Gyroidina soldanii</i> var. <i>octocamerata</i> CUSHMAN & HANNA <i>Trifarina advena</i> var. <i>californica</i> MALLORY <i>Siphonina wilcoxensis</i> CUSHMAN <i>Pleurostomella alazanensis</i> var. <i>cubensis</i> CUSHMAN & BERMUDEZ <i>Cibicides kernensis</i> MALLORY <i>Cibicoides</i> aff. <i>C. cushmani</i> (NUTTALL) <i>Globulina lacrima</i> REUSS <i>Bulimina</i> sp. <i>Bulimina</i> cf. <i>B. trihedra</i> CUSHMAN <i>Uuigerina elongata</i> COLE <i>Cibicoides allenii</i> (PLUMMER) <i>Gonatosphaera</i> sp. <i>Siphonina</i> sp. <i>Nonionella</i> cf. <i>N. ansata</i> CUSHMAN <i>Lenticulina</i> cf. <i>L. convergens</i> (BORNEMANN) <i>Anomalinoidea keeni</i> (MARTIN) ? " <i>Zeuuigerina</i> " sp. <i>Pleurostomella paleocenica</i> CUSHMAN <i>Nodosaria</i> cf. <i>N. velascoensis</i> CUSHMAN <i>Osangularia midwayana</i> (CUSHMAN & TODD) <i>Siphonina</i> cf. <i>S. jacksonensis</i> CUSHMAN <i>Lenticulina welchi</i> (CHURCH) <i>Cibicidina</i> cf. <i>C. mauricensis</i> HOWE <i>Cibicides</i> sp. <i>B.</i> (in GRAHAM & GLASSEN) <i>Eponides</i> sp. nov. ? <i>Chilostomelloides</i> sp. <i>Angulogerina abbreviata</i> var. <i>tubulifera</i> BATJES <i>Dorothia principiensis</i> CUSHMAN & BERMUDEZ <i>Pleurostomella</i> cf. <i>P. nuttalli</i> (CUSHMAN & SIEGFUS) <i>Anomalina garzaensis</i> CUSHMAN & SIEGFUS <i>Vaginulinopsis mexicana</i> var. <i>nudicostatus</i> CUSHMAN & HANNA <i>Vaginulinopsis</i> sp. <i>Bulimina consanguina</i> CUSHMAN & PARKER <i>Lenticulina</i> cf. <i>L. turbinata</i> (PLUMMER) <i>Siphonodosaria gracilis</i> (PALMER & BERMUDEZ) <i>Anomalinoidea crassiseptus</i> (CUSHMAN & SIEGFUS) <i>Silicosigmoina californica</i> CUSHMAN & CHURCH <i>Cribrostomoides</i> sp. <i>Eponides dorfi</i> TOULMIN <i>Nodogenerina adolphina</i> (d'ORBIGNY) <i>Globulina gibba</i> var. <i>globosa</i> (HANTKEN) <i>Haplophragmoides excavatus</i> CUSHMAN & WATERS <i>Pullenia quinqueloba</i> var. <i>angusta</i> (CUSHMAN & TODD) <i>Vaginulinopsis mexicana</i> var. <i>kerni</i> (COOK) <i>Ammobaculites</i> cf. <i>cubensis</i> CUSHMAN <i>Plectofrondicularia garzaensis</i> CUSHMAN & SIEGFUS " <i>Valvulinera</i> " sp. (in MALLORY) <i>Fursenkoina zetina</i> var. <i>indirecta</i> MALLORY <i>Bolivina pisciformis</i> GALLOWAY & MORREY <i>Cibicides grimsdalei</i> NUTTALL <i>Cibicides pachyderma</i> (RZEHA) <i>Bulimina excavata</i> CUSHMAN & PARKER <i>Dentalina basiplanata</i> CUSHMAN <i>Asterigerina crassaformis</i> CUSHMAN & SIEGFUS <i>Spiroplectammina tejonensis</i> MALLORY <i>Lenticulina antipoda</i> (STACHE) <i>Cyclammina</i> cf. <i>C. clarki</i> (HANNA)	
		Planktic foraminiferal biozonation	
		P8 P9 P10	

EXPLANATION
| Rare
| Frequent
| Common




FIGURE 6.—Foraminiferal biostratigraphy of Locatelli Formation, Smith Grade-Empire Grade area.

Foraminiferal faunas consist of predominantly agglutinated forms with minor amounts of calcareous benthic and planktic elements. Planktic foraminifers in samples from the Locatelli Formation include *Subbotina triloculinoides*, *S. triangularis*, *S. velascoensis*, *Acarinina mckannai*, *A. acarinata* and *Morozovella aequa* and indicate a later Paleocene age (*Planorotalites pseudomenardii* (P4)—*Morozovella velascoensis* (P5) Zones). Agglutinated benthic foraminifers of the Locatelli Formation are markedly similar to those that occur in the type Lodo Formation, some 250 km (150 mi) southeast of the Santa Cruz Mountains. Among the more characteristic are *Bathysiphon eocenicus*, *Cyclammina simiensis*, *Glomospira charoides*, *Haplophragmoides excavatus*, *Hyperammina elongata*, *Karreriella media-aguaensis*, *Rhabdammina eocenica*, *Rzehakina epigonalata*, *Spiroplectammina richardi*, and *Silicosigmoilina californica*.

Calcareous benthic taxa are sparsely represented in samples from the Locatelli Formation. Among the more diagnostic forms present are: *Anomalinoidea acutus*, *A. rubiginosus*, *Gavelinella beccariiiformis*, *Cibicidoides martinezensis*, *C. whitei*, *Osangularia* sp., and *Vaginulinopsis saundersi*.

The agglutinated benthic foraminiferal fauna of the Locatelli Formation, accompanied by infrequent calcareous benthic and planktic elements, has all the characteristics of a classic flysch assemblage. Most of the calcareous benthic taxa present are indicative of bathyal depths, although there is also a fairly diverse component of neritic elements (predominantly lagenids) similar to those in the Midway Group (see faunal list of Clark, 1966, table 2, also Clark in Brabb and others, 1977, table 5; we have also examined Clark's prepared slides of samples JC60-126, JC60-120, and JC60-93 in connection with this study). Precise estimates of deposition are not possible on the basis of this preliminary faunal examination but a general depth of 1-2 km is suggested.

LOMPICO AREA

The seven samples examined from the Butano Sandstone span a stratigraphic interval of approximately 1,000 m (fig. 7).

Calcareous faunas are exceedingly sparse in the samples of the Butano Sandstone we have examined, although Clark (1966, table 3, also Clark in Brabb and others, 1977, table 4) records a relatively diverse, albeit sparse, calcareous fauna (including lagenids, buliminids, pleurostomellids and anomalinids) from the middle siltstone member of the Butano Sandstone near Lompico.

Planktic foraminiferal faunas of the Butano Sandstone contain *Subbotina* sp. cf. *S. linaperta*, *S. turgida*, *Acarinina acarinata*, *A. coalingsensis*, *A. wilcoxensis*, and *Morozovella subbotinae* and indicate an early Eocene age (*Morozovella aragonensis* (P8 Zone) for at least part of the middle siltstone member.

Agglutinated assemblages from the Butano Sandstone are poorly preserved and not directly comparable with those observed at equivalent levels in the Lodo Formation or Gredal Shale Member of the Kreyenhagen Formation. In general the samples from the Butano lack the diversity seen in samples from the Lodo (absence of *Cyclammina*, *Silicosigmoilina*, *Rzehakina*, *Rhabdammina*, *Verneuilina*, *Gaudryina*), but this may be due to the small number of samples studied. Among the more diagnostic forms present in the Butano Sandstone are *Bathysiphon eocenicus*, *Karreriella mediaguaensis*, and *Haplophragmoides obliquicameratus*. Calcareous benthic foraminifers, although sparse, include several forms also present in the type Lodo Formation. Among these are *Buliminella bradburyi*, *Bolivina crenulata*, *Anomalinoidea welleri*, *Pleurostomella paleocenica*, *Hoeglundina eocenica*, and *Nuttallides truempyi*.

The agglutinated benthic fauna of the Butano suggests a depositional environment similar to that of the Locatelli Formation with water between 1 and 2 km.

SUMMARY

We have made a preliminary examination of samples from the type Lodo Formation in the Lodo Gulch area; the Lodo Formation and Gredal Shale Member of the Kreyenhagen Formation in the Devils Den area, and the Locatelli Formation and Butano Sandstone of the Santa Cruz Mountains, California. The following are our tentative conclusions.

1. The type Lodo Formation is of late Paleocene and early Eocene (Zones P4-P9) age and contains a benthic foraminiferal fauna that indicates deposition during a single sedimentary cycle ranging from lower neritic (somewhat shallower than 200 m) near the base to middle bathyal (about 600 m) in the lower part to outer neritic (200 m) in the upper part.
2. A relatively abrupt shallowing occurs near the boundary between Zones P8 and P9 in the Lodo Gulch area and is denoted by the disappearance of bathyal taxa (including *Nuttallides truempyi*) that have more extensive stratigraphic ranges elsewhere in deep-sea sections.

3. The common occurrence of cyclamminids and *Bathysiphon* in the type Lodo Formation at depths interpreted from other faunal evidence to have been near the shelf-slope break (about 200 m depth) suggests that the upper depth limit of the agglutinated benthic foraminiferal fauna was somewhat shallower during the Paleogene than during the Neogene and present day. This discrepancy may, in turn, be linked with the lack of a pronounced thermocline at the time.
4. The similarity of faunas from the Devils Den section with those in the Lodo Gulch area is marked; however, the presence of a rich planktic foraminiferal assemblage (particularly the angulo-conical morozovellids and acarininids) throughout the Devils Den section and the absence of shallow-water calcareous benthic assemblages (as seen in the type Lodo Formation) suggests deposition at middle bathyal depths (below 600 m) shallowing to upper bathyal (above 600 m) within Zone P9.
5. The relatively sudden shallowing in the late early Eocene (Zone P9) in these two sections (fig. 8) is remarkably close to a major eustatic sea-level change identified by Vail, Mitchum, and Thompson (1977) by seismic stratigraphic methods and merits further investigation.
6. The Locatelli Formation and Butano Sandstone of the Santa Cruz Mountains contain an agglutinated benthic foraminiferal fauna (with minor amounts of calcareous benthic and planktic elements) characteristic of flysch deposits elsewhere. Deposition in the distal part of turbidite fans at depths between 1 and 2 km is suggested.
7. In addition to containing elements that appear to be of Pacific affinities, the faunas of the Lodo Formation and the Gredal Shale Member of the Kreyenhagen Formation contain numerous cosmopolitan elements known from western Europe (Aquitaine Basin), West Africa (Angola), the North Atlantic (Orphan Knoll), and elsewhere. The agglutinated fauna appears to be less cosmopolitan, that is, not as closely related to the Paleogene flysch faunas of the Carpathians and North Atlantic.




FIGURE 8.—Late Paleocene and early Eocene bathymetric history of Lodo Gulch and Devils Den (aqueduct) areas. N.t., *Nuttallides truempyi*; As.c., *Asterigerina crassaformis*; A.c., *Anomalinoides crassiseptus*; A.a., *Anomalinoides aragonensis*; B.g., *Buliminella grata*; B.t., *Bulimina trinitatensis*. The large black "T" refers to the last occurrence of the selected species; the vertical squares indicate a rare or sporadic occurrence of species.

LOWER TERTIARY NANNOPLANKTON BIOSTRATIGRAPHY IN THE CENTRAL COASTAL RANGES, CALIFORNIA

By A. D. WARREN¹

ABSTRACT

Nannoplankton from the Devils Den, Lodo Gulch, and Media Agua Creek areas were correlated with Paleogene California benthic foraminiferal stages along the west side of the San Joaquin Valley. Based on nannoplankton chronology, the Ynezian Stage is of Paleocene and earliest Eocene age, the Bulitian, Penutian, and most of the Ulatisian Stages are of early Eocene age, and the upper Ulatisian and lower Narizian Stages are of early middle Eocene age. Estimated absolute ages for the tops of the Ynezian and Ulatisian Stages are 52 and 48 million years respectively.

Data from this and previous studies on the lower and middle Tertiary of California indicate the relations of the provincial benthic foraminiferal stages to nannoplankton and planktic foraminiferal age and zonal assignments. The Paleogene of California includes, from oldest to youngest, the Cheneyyan, Ynezian, Bulitian, Penutian, Ulatisian, Narizian, Refugian, and Zemorrian Stages.

INTRODUCTION

Calcareous nannoplankton are one of the most important fossil groups for correlation of marine deposits, particularly in the middle- and low-altitude areas of the world. The main purpose of this study is to examine the vertical distribution of calcareous nannofossils within the Paleogene California benthic foraminiferal stages as represented in three areas, Devils Den aqueduct, Lodo Gulch, and Media Agua Creek (fig. 9), along the west side of the San Joaquin Valley. Each of these areas contains exposures of marine sedimentary rocks which may be assigned to the Ynezian, Bulitian, Penutian, Ulatisian, and Narizian provincial stages of Mallory (1959) based on vertical distribution of benthic foraminifers. The secondary purpose of this paper is to reexamine the Paleogene of California by extracting from this and previously published studies a synthesis of the relations of the lower and middle Tertiary provincial stages to nannoplankton zonations and, in turn, to planktic foraminiferal zonations, which are in wide use today by biostratigraphers everywhere.

The most commonly used nannoplankton and planktic foraminiferal zonations are those of Bukry (1973,

1975), Martini (1971), Blow (1969), Berggren (1972), and Hardenbol and Berggren (1978). The relationship between these zonations is shown of figure 10. The series designations of Bukry (1973, 1975) will be followed in this report.

Acknowledgments.—For providing samples and a measured section from the Devils Den (aqueduct) area, I thank D. W. Weaver. I am also grateful to J. H. Newell for much additional supplemental nannofossil data and discussion on the Lodo Gulch and Devils Den areas.

DEVILS DEN (AQUEDUCT) AREA

The aqueduct section in the Devils Den area, in sec. 34, T. 25 S., R. 18 E., Kern County, California, exposes approximately 300 m of fossiliferous marine sedimentary rocks (fig. 11). Rock units exposed there are, from oldest to youngest, the upper part of the Panoche Formation, Lodo Formation, Avenal Sandstone, and Gredal Shale and Point of Rocks Sandstone Members of the Kreyenhagen Formation. A fault approximately 210 m below the top of the exposure repeats the Lodo, the Avenal, and part of the Gredal. The aqueduct section was measured and collected in 1970 by D. W. Weaver, and his samples and stratigraphic column form the basis for data presented on this exposure. Geologic and locality maps for this area have been provided by Brabb, Clark, and Throckmorton (1977, figs. 36 and 38).

Age assignments using the provincial benthic foraminiferal stages of Mallory (1959) are as follows (fig. 11): Samples PR-1 to PR-26 are assigned to the lower Narizian Stage based on the presence of the following selected foraminiferal taxa:

Alabamina wilcoxensis californica Mallory
Anomalina crassisepta Cushman and Siegfus
A. dorri aragonensis Nuttall
Asterigerina crassaformis Cushman and Siegfus
Bulimina corrugata Cushman and Siegfus
Buliminella grata convoluta Mallory
Caucasina schencki (Beck)
Cibicides laimingi Mallory
C. spiropunctatus Galloway and Morrey
Vaginulinopsis asperuliformis Nuttall

¹A. D. Warren, Consulting Micropaleontology, Inc., 7202 Clairemont Mesa Blvd., San Diego, CA 95211.




FIGURE 9.—Index map of the counties of California showing nannofossil localities.

STUDIES IN TERTIARY STRATIGRAPHY, CALIFORNIA COAST RANGES

PALEOGENE LOW-LATITUDE NANNOPLANKTON ZONATION OF BUKRY (1973, 1975)			PALEOGENE CALCAREOUS NANNOPLANKTON ZONATION OF MARTINI (1971) Boundaries between zones and series are dashed where relationship to Bukry's zonation approximate		
SERIES OR SUBSERIES	NANNOPLANKTON ZONES		SERIES OR SUBSERIES	NANNOPLANKTON ZONES	
Miocene	Triquetrorhabdulus carinatus	Discoaster druggii	Miocene	NN2 Discoaster druggi	
		Discoaster deflandrei		NN1 Triquetrorhabdulus carinatus	
		Cydicargolithus abisectus			
Oligocene	Sphenolithus ciperoensis		Upper Oligocene	NP25 Sphenolithus ciperoensis	
	Sphenolithus distentus			NP24 Sphenolithus distentus	
	Sphenolithus predistentus		Middle Oligocene	NP23 Sphenolithus predistentus	
	Helicopontosphaera reticulata	Reticulofenestra hillae		Lower Oligocene	NP22 Helicopontosphaera reticulata
		Coccolithus formosus	NP21 Ericsonia ? subdisticha		
		Coccolithus subdistichus			
	Upper Eocene	Discoaster barbadiensis	Isthmolithus recurvus	Upper Eocene	NP20 Sphenolithus pseudoradians
Chiasmolithus oamaruensis			NP19 Isthmolithus recurvus		
Middle Eocene	Reticulofenestra umbilica	Discoaster saipanensis	NP18 Chiasmolithus oamaruensis		
		Discoaster bifax	NP17 Discoaster saipanensis		
	Nannotetrina quadrata	Coccolithus staurion	Middle Eocene	NP16 Discoaster tani nodifer	
		Chiasmolithus gigas		NP15 Chiphragmalithus alatus	
		Discoaster strictus		NP14 Discoaster sublodoensis	
	Discoaster sublodoensis	Rhabdosphaera inflata	Middle Eocene	NP13 Discoaster lodoensis	
		Discoasteroides kuepperi		NP12 Marthasterites tribrachiatum	
Lower Eocene	Discoaster lodoensis		Lower Eocene	NP11 Discoaster binodosus	
	Tribrachiatus orthostylus			NP10 Marthasterites contortus	
	Discoaster diastypus	Discoaster binodosus		Upper Paleocene	NP9 Discoaster multiradiatus
		Tribrachiatus contortus			NP8 Heliolithus riedeli
Paleocene	Discoaster multiradiatus		Upper Paleocene	NP7 Discoaster gemmeus	
	Discoaster nobilis			NP6 Heliolithus kleinpellii	
	Discoaster mohleri		Middle Paleocene	NP5 Fasciculithus tympaniformis	
	Heliolithus kleinpellii			NP4 Ellipsolithus macellus	
	Fasciculithus tympaniformis		Lower Paleocene	NP3 Chiasmolithus danicus	
	Cruciplacolithus tenuis			NP2 Cruciplacolithus tenuis	
		NP1 Markalius inversus			
Upper Cretaceous			Upper Cretaceous	?	

FIGURE 10.—Diagram showing relationship between the nannoplankton zones of Bukry (1973, 1975) and Martini

PALEOGENE PLANKTONIC FORAMINIFERAL ZONATION (1978) MODIFIED AFTER BLOW (1969), BERGGREN (1972), AND HARDENBOL AND BERGGREN (1982)		
SERIES OR SUBSERIES	EUROPEAN STAGE BOUNDARY AGE IN M.Y.B.P.	PLANKTONIC FORAMINIFERAL ZONE
Miocene	Aquitainian	N5 <i>Globoquadrina dehiscens praedehiscens</i> - <i>Globoquadrina dehiscens</i>
		N4 <i>Globigerinoides quadrilobatus primordius</i> / <i>Globorotalia kugleri</i>
Upper Oligocene	Chattian	P22 <i>Globigerina angulissuturalis</i>
		P21 <i>Globigerina angulissuturalis</i> / <i>Globorotalia opima opima</i>
		P20 <i>Globigerina ampliapertura</i>
Lower Oligocene	Rupelian	P19 <i>Globigerina sellii</i> / <i>Pseudohastigerina barbadoensis</i>
		P18 <i>Globigerina tapuriensis</i>
		P17 <i>Globigerina gortanii gortanii</i> / <i>Globorotalia centralis</i>
Upper Eocene	Priabonian	P16 <i>Cribrohantkenina inflata</i>
		P15 <i>Globigerapsis mexicana</i>
Middle Eocene	Bartonian	P14 <i>Truncorotaloides rohri</i> - <i>Globigerinita howei</i>
		P13 <i>Orbulinoides beckmanni</i>
	Lutetian	P12 <i>Globorotalia lehneri</i>
		P11 <i>Globigerapsis kugleri</i>
		P10 <i>Hantkenina aragonensis</i>
Lower Eocene	Ypresian	P9 <i>Acarinina densa</i>
		P8 <i>Globorotalia aragonensis</i>
		P7 <i>Globorotalia formosa</i>
		P6 <i>Globorotalia subbotinae</i> / <i>Acarinina wilcoxensis</i>
Upper Paleocene	Thanetian	P5 <i>Globorotalia velascoensis</i>
		P4 <i>Globorotalia pseudomenardii</i>
		P3 <i>Globorotalia pusilla</i> - <i>Globorotalia angulata</i>
Lower Paleocene	Danian	P2 <i>Globorotalia uncinata</i> - <i>Globigerina spiralis</i>
		P1 <i>Globoconusa daubjergensis</i>
Upper Cretaceous	Maestrichtian	?

(1971) and the planktic foraminiferal zones of Blow (1969), Berggren (1972) and Hardenbol and Berggren (1978).






FIGURE 11A.—Biostratigraphy of the aqueduct section of the Devils Den Area. A is the upper part of the section exposed near the penstocks.



EXPLANATION

FREQUENCY SYMBOLS



?

Rare, possibly reworked or questionable occurrences

Rare

Rare to few

Common to abundant

METERS

30

20

10

0

FIGURE 11B.—Lower part of the aqueduct section in the Devils Den area. Section measured and collected in 1970 by D. W. Weaver (PR sample numbers). Other samples were collected by E. E. Brabb. See Berggren and Aubert, this volume, for discussion. Lithostratigraphy from Dibblee (1973). Series assignments for formations are based on repeated faulted nannoplankton zonal assignments of Bukry (1973a, b). (See also Berggren and Aubert, this volume.)

Samples PR-27 to PR-31 cannot be assigned with certainty to either the Ulatisian or Narizian Stage although most workers would probably classify these samples as Ulatisian based on rare to few occurrences of the following selected taxa:

Bulimina debilis Martin

Gaudryina jacksonensis coalingensis Cushman and Hanna

Nodosaria latejugata Gumbel

Vaginulinopsis mexicana (Cushman) vars.

Samples PR-32 to PR-49 are assigned to the Ulatisian Stage on the basis of the following selected taxa:

Bifarina nuttalli Cushman and Siegfus

Globorotalia aragonensis Nuttall

Nodosaria latejugata Gumbel

Lenticulina ulatisensis (Boyd)

Trifarina advena californica Mallory

Uvigerina lodoensis Martin and vars.

Vaginulinopsis mexicana (Cushman) vars.

Valvulineria indiscriminata Mallory

McKeel and Morin (1979) have investigated the taxonomic status of *Valvulineria indiscriminata* Mallory and have determined that it is a planktic species referable to the genus *Planorotalites*. In addition, they consider *Globorotalia pseudochapmani* Gohrbandt to be a junior synonym of *Planorotalites indiscriminata* (Mallory), having had the opportunity of comparing hypotypes of the latter species to paratypes or topotypes of the former provided by Gohrbandt and finding them to be conspecific. Mallory (1959) indicates that this species is restricted to his Ulatisian Stage.

Samples PR-50 and PR-51 are either Penutian or Ulatisian as the taxa present are not clearly diagnostic of either stage.

The Avenal Sandstone is assigned to the Penutian Stage on the basis of *Pseudosphragmina psila* (Woodring), and *Rotularia (Spiroglyphus) tinajasensis* (Hanna and Hertlein), a worm tube, collected by the author from the Avenal Sandstone.

Samples PR-52 and PR-53 are assigned to the Penutian Stage based on the occurrence of the following selected taxa:

Angulogerina wilcoxensis (Cushman and Ponton)

Eponides lodensis Martin

Pseudosphragmina psila (Woodring)

Lenticulina laimingi (Israelsky)

Siphonina wilcoxensis Cushman

Verneuilina triangulata Cook

Lenticulina laimingi (Israelsky) was first recog-

nized as a distinctive species by Cook (1950) but was not formally described at that time. The species appears to range from within the Bulitian Stage to the top of the lower Penutian Stage and is present in the Lodo Formation at its type locality in the Lodo Gulch area and in the Lodo Formation in the Devils Den and Media Agua Creek areas.

Samples PR-54 to PR-64 are assigned to the Bulitian Stage on the basis of the following selected taxa:

Bulimina excavata Cushman and Parker (rare)

Cibicides kernensis Cook

Globorotalia californica Smith (rare)

Lenticulina laimingi (Israelsky)

Samples PR-65 to PR-70 are assigned to the upper Ynezian Stage on the basis of the following selected taxa:

Anomalina dorri Cole

Bulimina excavata Cushman and Parker (common)

Chiloguembelina midwayensis subcylindrica Beckman

Globorotalia californica Smith (common)

G. marginodentata Subbotina

Samples PR-71 to PR-98 are in that part of the section repeated by faulting and represent part of the Ulatisian, Penutian, Bulitian, and Ynezian Stages. These stage assignments are based on the occurrence of the same foraminifers below the fault as above the fault, thus providing paleontological evidence of the repeated section.

Samples PR-99 to PR-102 are from the upper part of the Panoche Formation and contain only crushed arenaceous foraminifers except for sample PR-101, in which Cretaceous radiolarians were recognized by M. B. Mickey (oral commun., 1975). Radiolarian species present are *Amphipyndax plousios*, *Dictyomitra* cf. *multicostata* and *Stichomitra asymbatos*, all of which have been reported from the Late Cretaceous of California by Foreman (1968).

Some of the nannoplankton from the Devils Den section are shown on Figure 10. Additional nannoplankton from samples collected by E. E. Brabb are shown on table 3. The stratigraphic position of the Brabb samples is shown in the report by Aubert and Berggren (this volume, fig. 4).

The ages of nannoplankton from the aqueduct section in the Devils Den area are also shown on figure 11 and table 3. These age assignments follow the zonal criteria and series correlations proposed by Bukry (1973a and b, 1975). Nannoplankton zones recognized are the *Discoaster diastypus*, *Tribrachia-*

tus orthostylus, *Discoaster lodoensis*, *Discoaster sublodoensis*, and *Nannotetrina quadrata* Zones. Placement of boundaries between the *Tribrachiatus orthostylus* and *Discoaster lodoensis* Zones and the *Discoaster sublodoensis* and *Nannotetrina quadrata* Zones are not precise. The boundary between the *Tribrachiatus orthostylus* and *Discoaster lodoensis* Zones is arbitrarily and tentatively drawn here to approximate the highest occurrence of persistent and common to abundant *Tribrachiatus orthostylus* and the lowest occurrence of *Coccolithus crassus* s. ampl. The boundary between the *Discoaster sublodoensis* and *Nannotetrina quadrata* Zones must be below sample PR-21 and above sample PR-30, but the absence of both *Rhabdosphaera inflata* and *Nannotetrina quadrata* in samples from that interval precludes a more precise placement based on the criteria proposed by Bukry (1973). The highest occurrence of persistent and common to abundant *Discoaster sublodoensis* as noted in sample PR-27 might serve locally as a useful reference point approximating the top of the *Discoaster sublodoensis* Zone.

Having now resolved the placement of both provincial stage and nannoplankton zonal boundaries within the aqueduct section in the Devils Den area, we may draw the following conclusions regarding that section:

1. The boundary between the Ynezian and Bulitian States lies within the lower part of the *Tribrachiatus orthostylus* Zone and very near the top of the *Discoaster diastypus* Zone, and the boundary between the Ulatisian and Narizian Stages is closely associated with the top of the *Discoaster sublodoensis* Zone.
2. Age assignments based on nannoplankton as proposed by Bukry (1973, 1975) require that the upper Ynezian, Bulitian, Penutian, and lower and middle Ulatisian Stages be of early Eocene age and that the uppermost Ulatisian and lower Narizian be of early middle Eocene age.
3. A hiatus or unconformity at the contact between the Upper Cretaceous Panoche Formation and the Lodo Formation is suggested by the absence of any Paleocene nannoplankton zones.
4. At Devils Den, the Panoche Formation is of Late Cretaceous age, and, on the basis of nannoplankton, the Lodo Formation, Avenal Sandstone, and lower part of the Gredal Shale Member of the Kreyenhagen Formation are of early Eocene age, and the upper part of the Gredal Shale Member and lower part of the Point of Rocks Member of

the Kreyenhagen are of middle Eocene age.

LODO GULCH AREA

The Lodo Formation at its type locality at Lodo Gulch, secs. 28 and 29, T. 15 S., R. 12 E., Fresno County, California, is a continuous section of marine sedimentary rocks approximately 360 m thick. Martin (1943) and Israelsky (1951, 1955) provided foraminiferal data on the type Lodo, which Mallory (1959, p. 29, 33, 37, 51, 53 and fig. 7) used to generalize the approximate position of his lower Tertiary benthic foraminiferal stages within that section. In a classic report, Bramlette and Sullivan (1961) described in detail the nannoplankton flora of the type Lodo Formation including data on the overlying Domenigine and Kreyenhagen Formations from other localities. Utilizing distributional data of Bramlette and Sullivan (1961, table 1) and Hay and others (1967, fig. 5), Poore (1976) reinterpreted zonal assignments within the type Lodo Formation to prove the presence of six nannoplankton zones within that unit, the *Discoaster mohleri*, *D. nobilis*, *D. multiradiatus*, *D. diastypus*, *Tribrachiatus orthostylus*, and *Discoaster lodoensis* Zones of Bukry (1973); the lower three zones are placed in the Paleocene by the majority of nannoplankton workers, and the higher three are considered to be early Eocene. Figure 12 illustrates the relations in the Lodo Gulch area of the formations, provincial benthic foraminiferal stages, vertical distribution of key nannofossil taxa, and nannoplankton zones to one another as derived from published data of the authors cited above. Additional nannoplankton from this section are shown on table 3.

Keeping in mind that the placement of benthic foraminiferal stage boundaries in figure 12 is only approximate, the following conclusions may be drawn as regards the exposures in the Lodo Gulch area:

1. The Ynezian-Bulitian boundary as approximately delimited by Mallory (1959, p. 29 and 33) appears to be within the *Discoaster diastypus* Zone.
2. Following Bukry's (1973a and b, 1975) correlation of nannoplankton zones with European Tertiary series, and the correlations in this report between the nannoplankton zones and the benthic foraminiferal stages, most of the Ynezian Stage is of Paleocene age, and the uppermost Ynezian, Bulitian, Penutian, and lower and middle Ulatisian Stages are of early Eocene age.
3. A hiatus may be present between the Moreno Shale and the overlying type Lodo Formation as

collected by E. E. Brabb in the Devils Den and Lodo Gulch areas
report by Berggren and Aubert (this volume, fig.4)]

				••			••	•	•		••	•		•	Nannotetrina quadrata	Discoaster strictus
•	•			••		•		••	•		••	•		•	Discoaster subloboensis	Discoasteroides kuepperi
	••	•		••	••		•			•	•	•	••	•	Discoaster lodoensis	
	•	••		••	•						•	•	••	•		
	••	•		••	•						•	•	••	•		
	••	•		••	•						•	•	••	•		
	••	•		••	•						•	•	••	•	Tribrachiatus orthostylus/ Discoaster lodoensis undifferentiated	
	••	•		••	•						•	•	••	•		
	••	•		••	•						•	•	••	•		
	••	•		••	•						•	•	••	•	Tribrachiatus orthostylus	
		•	••	••		•					•	•	••	•	Discoaster multiradiatus	
D. subloboensis															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
D. tari																
D. tari ornatus															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
D. uermelensis																
Discoasteroides kuepperi															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
D. megastypus																
Discolithina plana															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
Ellipsolithus macellus																
Fasciculithus involutus															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
F. mitreus																
F. schaubi															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
F. tymaniformis																
Helicopontosphaera lophota															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
H. seminulum																
Isthmolithus recurvus															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
Lophodolichus mochiophonus																
L. nactus															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
L. reniformis																
Markalius astroponus															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
Microantholithus entaster																
M. vesper															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
Nannotetrina fulgens																
N. quadrata															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
Neococcolithes dubius																
N. protenus															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
Reticulofenestra dictyoda																
R. hillae															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
R. pseudoumbilica																
R. samodurovi															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
R. umbilica																
Rhabdosphaera crebra															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
R. inflata																
Sphenolithus anarhogus															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
S. furciculoides																
S. moriformis															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
S. pseudoradians																
S. radians															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
S. spiniger																
Touletia craticulus															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
T. eminens																
Transversopontis pulcher															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
Tribrachiatus orthostylus																
Zygodiscus adamas															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
Z. plectopons																
Zugrhablithus bijugatus															NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)	
	•				•				•						D. barbadiensis to H. reticulata	I. recurvus to C. subdistichus
	••				•			••	•			••	•		Nannotetrina quadrata	Chiasmolithus gigas
	•	•			•			••	•			••	•			Discoaster strictus
•					•			••	•			•	•		Probable Nannotetrina quadrata	
••	•				••	•		•	•			••	•		Discoaster subloboensis	Rhabdosphaera inflata
••	••				••	•		•	•			••	•			Discoasteroides kuepperi
	•				••			•				••	•		Discoaster lodoensis	
	•	•			••							•	•	•	Tribrachiatus orthostylus	
	•	•			••			•				•	•	•	Discoaster diastypus	Discoaster binodosus
BY FAULTING																
•	•			••	•		•	••			•	•		•	Discoaster subloboensis	D. kuepperi
	••			••	•			•				•		•	Discoaster lodoensis	
				••	••			•				••	•	•		
	•				••			•				••	•	•	Tribrachiatus orthostylus	




FIGURE 12.—Selected nannofossil distributions, nannoplankton zonation, and benthic foraminiferal stages in the Lodo Gulch area. The lithostratigraphy is from Israelsky (1951, fig. 2). Sample numbers and nannoplankton frequency and range are from Bramlette and Sullivan (1961). Nannoplankton zonal assignments are by Poore (1976). Benthic foraminiferal stage assignments are by Mallory (1959). Dashed lines indicate stage and zone boundaries are approximately located.

suggested by the absence of any Paleocene nannofossil zones older than the *Discoaster mohleri* Zone within the type Lodo.

- On the basis of nannoplankton in the Lodo Gulch area, the Lodo Formation is of late Paleocene and early Eocene age.

MEDIA AGUA CREEK AREA

At Media Agua Creek, in sec. 22, T. 28 S., R. 19 E., Kern County, California, a continuous section of lower Tertiary marine sedimentary rocks over 700 m thick is exposed. Rock units present, from oldest to youngest, are the Lodo(?) Formation, Avenal Sand-

stone, Gredal Shale and Point of Rocks Sandstone Members of the Kreyenhagen Formation following stratigraphic nomenclature modified from Dibblee (1973a, p. 8-13). Mallory (1959, 1970), after detailed foraminiferal analysis, selected the Media Agua Creek exposures as the type sections for the benthic foraminiferal zones of his Ynezian and Bulitian Stages and the type section of his Penutian Stage; he also delimited the position of the Ulatisian and lower Narizian Stages within that section. Sullivan (1964, 1965) provided detailed nannoplankton data from Mallory's samples; Poore (1976) used these data to establish the position of the nannoplankton zones of Bukry (1973) in the Media Agua Creek section. Consequently, the Media Agua Creek section is perhaps the single most important reference section for the lower Tertiary in California by virtue of its wealth of disciplinary controls.

Figure 13 illustrated the relations at Media Agua Creek of the provincial stages, formations, vertical distribution of key nannofossil taxa, and nannoplankton zones to one another as derived from published data of the authors cited in the preceding paragraph. However, I have placed the boundary between the *Discoaster lodoensis* and *Discoaster sublodoensis* Zones at the approximate middle of the Gredal Shale Member rather than at the base of the Gredal as shown by Poore (1976, pl. 1); this interpretation is based on the first stratigraphic occurrence of *Discoaster sublodoensis* in Sample A-7052 as reported by Sullivan (1965).

From previously published data from the various authors cited above, the following conclusions may be drawn as regards the Media Agua Creek section:

1. The Ynezian-Bulitian boundary at Media Agua Creek is coincident with the top of the *Discoaster diastypus* Zone.
2. Age assignments based on nannoplankton as proposed by Bukry (1973, 1975) require that the lower Ynezian Stage be Paleocene and that the upper Ynezian, Bulitian, Penutian, and the lower Ulatisian be of early Eocene age. It is not possible from the available data to determine if the middle and upper Ulatisian at Media Agua Creek is in part of early middle Eocene age as noted in other exposures along the west side of the San Joaquin Valley, but the lower Narizian here is clearly of early middle Eocene age.
3. The absence of the *Discoaster multiradiatus* Zone suggests a hiatus within that part of the Lodo(?) Formation assigned by Mallory to the Ynezian Stage. A second hiatus or unconformity may be present at or near the contact between the Panoche and Lodo(?) Formations as suggested by

the absence of any recognizable Paleocene nannoplankton zones older than the *Discoaster nobilis* Zone.

4. At Media Agua Creek, on the basis of nannoplankton, the lowermost 10 m of the Lodo(?) Formation is Paleocene, and the remainder of the Lodo(?), the Avenal Sandstone, and the lower part of the Gredal Shale Member of the Kreyenhagen Formation are of early Eocene age. The upper part of the Gredal Shale Member and the lower part of the Point of Rocks Sandstone Member of the Kreyenhagen Formation (Mallory, 1970, fig. 3) are late early Eocene and (or) early middle Eocene age, and the middle and upper parts of the Point of Rocks Sandstone Member are of early middle Eocene age.

COMPARISON OF THE THREE AREAS

New and previously published data from the Devils Den, Lodo Gulch, and Media Agua Creek areas make possible a comparison of lower Tertiary benthic foraminiferal stages of California and nannoplankton zones as represented along the west side of the San Joaquin Valley. Figure 14 illustrates those relations and permits the following observations to be made:

1. The Ynezian Stage in this area has been shown to include the *Discoaster mohleri*, *Discoaster nobilis*, *Discoaster multiradiatus*, *Discoaster diastypus* and lowermost part of the *Tribrachiatus orthostylus* Zones and thus is of late Paleocene to early Eocene age based on the nannoplankton chronology of Bukry (1973a and b, 1975). The Bulitian, Penutian, and Ulatisian Stages include the *Discoaster diastypus*, *Tribrachiatus orthostylus*, *Discoaster lodoensis*, and the *Discoaster sublodoensis* Zones and are thus of early Eocene and early middle Eocene age. The top of the Ulatisian Stage appears to coincide with the top of the *Discoaster sublodoensis* Zone, for which Bukry estimated an absolute age of 48 million years. The lower Narizian Stage is represented by the *Nannotetrina quadrata* Zone of early middle Eocene age, for which Bukry estimated an absolute age at the zonal top of 45 million years.
2. The time-transgressive nature of rock units is rather strikingly demonstrated along the west side of the San Joaquin Valley by the time-stratigraphic relations between the Gredal Shale and the Point of Rocks Sandstone Members of the Kreyenhagen Formation at the Media Agua Creek and at the Devils Den aqueduct sections. Mallory (1970, fig. 3) divided his Point of Rocks Formation at Media Agua Creek into lower, middle, lower-

SERIES	PROVINCIAL CALIFORNIA STAGE	ROCK UNIT	SAMPLE NUMBER	METERS FROM BASE OF POINT OF ROCKS SANDSTONE	SELECTED NANNOFOSSILS										FREQUENCY AND RANGE SYMBOLS		NANNOPLANKTON ZONES AND SUBZONES OF BUKRY (1973)
					<i>Chiasmolithus gigas</i>	<i>Coccolithus crassus</i>	<i>Discoaster diastypus</i>	<i>Discoaster lodoensis</i>	<i>Discoaster multiradiatus</i>	<i>Discoaster nobilis</i>	<i>Discoaster subloboensis</i>	<i>Fasciculithus involutus</i>	<i>Helolithus riedeli</i>	<i>Nannotetrina quadrata</i>	<i>Tribrachiatus orthostylus</i>	? Rare, possibly reworked or questionable occurrences Rare Rare to few Common to abundant	
Eocene	Early Narizian	Point of Rocks Sandstone Member	A-7025	371												Nannotetrina quadrata	Chiasmolithus gigas
			A-7023	347													Discoaster strictus
			A-7020	329													
			A-7009	196													
	Ulatisian	Kreyenhagen Formation (part)	A-7038	159											?		Discoaster subloboensis
			A-7037	157											?		
			A-7046	-0.3											?		
			A-7052	-10													Discoaster lodoensis
	Penutian	Gredal Shale Member	A-7055	-15													
			A-7061	-22													
			A-7088	-25													
	Bulitian	Avenal Sandstone	A-7091	-27													Tribrachiatus orthostylus
			A-7062	-30.8					?								
			A-7063	-31.4													
			A-7064	-36													
Paleocene	Ynezian	Lodo(?) Formation	A-7070	-48					?								Discoaster diastypus
			A-7071	-51									?				
			A-7075	-70									?				
			A-7076	-72													
			A-7078	-78													Hiatus ?
			A-7079	-79.5													
Upper Cretaceous	Ciervian or older	Panoche Formation	A-7083	-85													Hiatus ?

FIGURE 13.—Formations, selected nannofossil distributions, nanoplankton zonation, and benthic foraminiferal stages in the Media Agua Creek area. Lithostratigraphy is from Dibblee (1973a, p. 8-13). Sample numbers and nanoplankton frequency and range are from Sullivan (1964, 1965). Nanoplankton zonal assignments are by Poore (1976). Dashed lines indicate boundaries of the zone are approximately located. Benthic foraminiferal stage assignments are by Mallory (1959).

SERIES	CALIFORNIA PROVINCIAL STAGE	MEDIA AGUA CREEK		DEVILS DEN (AQUEDUCT SECTION)		LODO GULCH AREA	
Eocene	Narizian (part)	Not studied		Nannotetrina quadrata	Coccolithus staurion	Not studied	
		Nannotetrina quadrata	Chiasmolithus gigas		Chiasmolithus gigas		
			Discoaster strictus		Discoaster strictus		
	Ulatisian	Discoaster sublodoensis		Discoaster sublodoensis	Rhabdosphaera inflata	Discoaster sublodoensis	Rhabdosphaera inflata
					Discoasteroides kuepperi	Discoasteroides kuepperi	
						Hiatus	
		Discoaster lodoensis		Discoaster lodoensis		Discoaster lodoensis	
	Penutian						
	Bulitian	Tribrachiatus orthostylus		Tribrachiatus orthostylus		Tribrachiatus orthostylus	
Ynezian	Discoaster diastypus		Discoaster diastypus		Discoaster diastypus		
	Hiatus		Hiatus		Discoaster multiradiatus		
					Hiatus		
					Discoaster nobilis		
	?				Discoaster mohleri		
Cheneyan	Hiatus						Hiatus

FIGURE 14.—Correlations of provincial benthic foraminiferal stages of California to nannoplankton zones and subzones, Devils Den, Lodo Gulch, and Media Agua Creek areas.

upper and upper-upper members. Dibblee (1973a, p. 10 and 12) assigned clay shale underlying his Point of Rocks Sandstone at Media Agua Creek to the Gredal Shale Member of the Kreyenhagen. Nannoplankton and foraminiferal data from this and previous studies indicate that the Gredal and lower and middle parts of the Point of Rocks at Media Agua Creek are equivalent in a time sense to the entire Gredal exposed in the aqueduct section in the Devils Den area and that the unit referred to as the lower-upper member of the Point of Rocks at Media Agua Creek is equivalent to that part of the Point of Rocks exposed at the Devils Den aqueduct section. Thus the base of the Point of Rocks at Media Agua Creek is much older than the base of the Point of Rocks at the Devils Den aqueduct; the difference in age may be estimated at 2.5 to 3 million years by referring to the estimated absolute ages of the nannoplankton zones and subzones of Bukry (1975, table 2). At Point of Rocks, a locality within the Devils Den area only 2.5 km southeast of the aqueduct, the basal 23 m of the Point of Rocks Sandstone Member was assigned by Mallory (1959, p. 55 and 57) to the Ulatisian, whereas at the aqueduct exposure the top of the Ulatisian is within the Gredal Shale Member at least 50 m below the base of the Point of Rocks Sandstone Member. Another case of a formation boundary transgressing time is exemplified by the difference in age of the top of the type Lodo Formation in the Lodo Gulch area compared to the top of the Lodo Formation in the Devils Den area and in the Lodo(?) Formation at Media Agua Creek. Nannoplankton and foraminiferal data indicate that the type Lodo Formation is equivalent in a time sense to the Lodo and Lodo(?) Formations, Avenal Sandstone and the lower part of the Gredal Shale Member of the Kreyenhagen Formation as exposed at Devils Den and Media Agua Creek. Thus the top of the Lodo Formation at its type locality in the Lodo Gulch area is demonstrably younger than the top of the Lodo Formation at Devils Den and the Lodo(?) Formation at Media Agua Creek as defined by Dibblee (1973a).

3. Nannoplankton data from this study and from Bramlette and Sullivan (1961) and Sullivan (1965) further suggest that the Domengine Sandstone in the Lodo Gulch area, the Domengine Formation and Canoas Siltstone Member of the Kreyenhagen Shale at Oil City (fig. 9), and the Canoas at Garza Creek are time equivalents to at least part of the upper 82 m of the Gredal Shale Member in the Devils Den aqueduct section and to at least parts

of the lower and middle Point of Rocks Sandstone Member in the Media Agua Creek section.

THE PALEOGENE OF CALIFORNIA: SUMMARY

It is now feasible to summarize the relations of the lower and middle Tertiary provincial benthic foraminiferal stages of California to nannoplankton and planktic foraminiferal zonations that are being widely used today for correlation of marine sedimentary deposits throughout the world. Figure 15 illustrates those relations and is a synthesis of nannoplankton, foraminiferal, and stratigraphic data from this report and from the following authors: Berggren (1972), Blow (1969), Brabb, Bukry, and Pierce (1971), Bramlette and Sullivan (1961), Bukry (1973, 1975), Bukry, Brabb, and Vedder (1977), Cifelli (1951), Cook (1950), Goudkoff (1945), Hardenbol and Berggren (in press), Hay and others (1967), Kleinpell (1938), Loeblich (1958), Mallory (1959, 1970), Martin (1964), Martini (1971), Poore (1976), Schenck and Kleinpell (1936), Schmidt (1970, 1975), Smith (1974), Sullivan (1964, 1965), Tipton (1980), Warren (1980), and Warren and Newell (1976). Most of the nannoplankton data used for determining relations between nannoplankton zones and provincial benthic foraminiferal stages as shown in figure 15 were obtained from the type sections and type areas for these stages or from the type localities of the foraminiferal zones used to define such stages with three exceptions; no nannoplankton data have as yet been obtained from the type sections or areas of the Cheneyan, upper Narizian, or Zemorrian Stages. The *Cruciplacolithus tenuis* and *Sphenolithus distentus* Zones are shown on figure 15 in their normal positions within the zonal sequence of Bukry (1973), although I believe neither zone has been recognized to date in California. Figure 15 shows that on the basis of nannoplankton data the Cheneyan and most of the Ynezian are of Paleocene age, the uppermost Ynezian, Bulitian, Penutian, and most of the Ulatisian are of early Eocene age, the upper Ulatisian and most of the Narizian are of middle Eocene age. Most of these age assignments are at variance with ages (based essentially on benthic foraminifers and megafossils) assigned by Kleinpell (1938), Mallory (1959), and Goudkoff (1955).

To be sure, additional data would be welcome before every detail of figure 15 is accepted as fact, particularly precise placement of most of the planktic foraminiferal zones P1 through P14. However, assuming that future data would cause no major readjustment in age at the base of the Cheneyan or the top of the Zemorrian Stages (and there is good reason for

ORIGINAL PROVINCIAL AGE/STAGE ASSIGNMENTS OF KLEINPELL (1938), SCHENCK AND KLEINPELL (1936), MALLORY (1959) AND GOUDKOFF (1945)		AGE ASSIGNMENTS USED IN THIS REPORT BASED ON PALEOGENE LOW-LATITUDE NANNOPLANKTON ZONATION OF BUKRY (1973 AND 1975)			PALEOGENE NANNOPLANKTON ZONATION OF MARTINI (1971)	PALEOGENE PLANKTONIC FORAMINIFERAL ZONATION MODIFIED AFTER BLOW (1969), BERGGREN (1972), AND HARDENBOL AND BERGGREN (1982)																						
SERIES OR SUBSERIES	STAGE	SERIES OR SUBSERIES	ZONE/SUBZONE		ZONE	SERIES OR SUBSERIES	EUROPEAN STAGE BOUNDARY AGE IN M.Y.B.P.	ZONE																				
Upper Eocene	Narizian	Middle Eocene	Reticulofenestra umbilica	Discoaster saipanensis	NP17	Middle Eocene	40	Bartonian	P14																			
				Discoaster bifax	NP16				P13																			
			Nannotetrina quadrata	Coccolithus stauroion	NP15		44	Lutetian	P12																			
				Chiasmolithus gigas					P11																			
				Discoaster strictus					P10																			
Middle Eocene	Ulatisian	Lower Eocene	Discoaster subloboensis	Rhabdosphaera inflata Discoasteroides kuepperi	NP14	Lower Eocene	49	Ypresian	P9																			
Lower Eocene	Penutian		Discoaster lodoensis		NP13				P8																			
			Tribachiatus orthostylus		NP12				P7																			
Paleocene	Bulitian		Discoaster diastypus	Discoaster binodosus					NP11	Upper Paleocene	53.5	Thanetian	P6															
				Tribachiatus contortus					NP10				P5															
	Ynezian		Discoaster multiradiatus		NP9				NP8				NP7	NP6	NP5	NP4	P4											
			Discoaster nobilis		Cruciplacolithus tenuis													NP3	Lower Paleocene	60	Danian	P2						
			Discoaster mohleri																				NP2	NP1	Upper Cretaceous	65	Maestrichtian	-
			Heliolithus kleinpellii																									
			Fasciculithus tympaniformis																									
	Cheneyan																											
	Upper Cretaceous		Ciervian	Upper Cretaceous																								

FIGURE 15.—Correlations of Paleogene provincial benthic foraminiferal stages of California to calcareous microplankton zonations.

making that assumption), then the Paleogene of California should be recognized as including all of those provincial stages from the base of the Cheney-

an to the top of the Zemorrian which have been shown by nannoplankton chronology to range in age from Paleocene to Oligocene.

CALCAREOUS NANNOFOSSIL BIOSTRATIGRAPHY OF THE PALEOGENE OF THE SANTA CRUZ MOUNTAINS, CALIFORNIA

By BILAL U. HAQ¹

Eighteen samples of calcareous nannofossils (centrifugally concentrated) were examined in detail from the sections of the Locatelli Formation in the Smith Grade area, the Butano Sandstone in the Lompico area, the Butano Sandstone and Twobar Shale Member of the San Lorenzo Formation in the Bear Creek area, and the Twobar Shale Member along the San Lorenzo River. The preservation of the nannofossils is generally very poor, specimens are rare and zone diagnostic species are lacking. Age assignments are, consequently, difficult and time-consuming and have to be based on the overall composition of the

assemblages. Zonal assignments are thus approximate, based mainly on the current ranges of taxa (after Bukry, 1971 and Martini, 1971).

In the Smith Grade area, collection 76CB1511A from the Locatelli Formation has only a few fragments of late Paleocene discoasters, but collection JC60-120d (table 4) from 4.5 km to the east has an assemblage that can be correlated with the *Fasciculithus tympaniformis* (NP5) Zone of late Paleocene age. In the Lompico area, collections JC60-37d, 76CB1521, JC60-51 and JC62-24b indicate that the lower part of the Butano Sandstone is of early Eocene (NP12 and NP13 Zones) and middle Eocene (NP15 and NP16 Zones) age. Collection 76CB1573 and

¹Woods Hole Oceanographic Institution, Woods Hole, Mass. 02543

TABLE 4.—*Nannoplankton from the Paleogene of the Santa Cruz Mountains*
[JC samples collected by J. C. Clark, CB samples collected by E. E. Brabb. n.d., not determinable]

Nannoplankton	¹ JC60-120d	² JC60-37d	² 76CB-1521	² JC60-51	² JC62-24b	² 76CB-1574	² 76CB-1573	² 68CB-202	² 68CB-204	² 68CB-205	² 68CB-206	² 76CB-1571	² 76CB-1451A
<i>Chiasmolithus californicus</i>								X					
<i>C. consuetus</i>	X												
<i>C. expansus</i>						X							
<i>C. gigas</i>													X
<i>Coccolithus bisectus</i>										X	X	X	
<i>C. eopelagicus</i>		X			X	X	X	X	X	X	X	X	X
<i>C. inflatus</i>	X	X			X	X	X	X	X	X	X	X	X
<i>C. inversus</i>		X			X		X	X	X	X	X	X	X
<i>Cyclolithella inflexa</i>			X	X	X								X
<i>Dictyococites hesslandii</i>							X						X
<i>Discoaster barbadensis</i>					X		X	X		X		X	X
<i>D. binodosus</i>							X						?
<i>D. delicatus</i>							X						X
<i>D. distinctus</i>													
<i>D. lodoensis</i>		X	X										
<i>D. mirus</i>			X					X	X				
<i>D. saipanensis</i>							X	X					
<i>Ericsonia subpertusa</i>	X	X											
<i>Fasciculithus tympaniformis</i>	X												
<i>Helicosphaera seminulum</i>					X								
<i>Heliorthus concinnus</i>						X							
<i>Impatiaster obscurus</i> ?								X					
<i>Lithostromation perdurum</i>													X
<i>Nannotetrina</i> aff. <i>N. cristata</i>			X										X
<i>Pemma angulatum</i>													
<i>Pontosphaera multipora</i>					X								
<i>P. plana</i>					X	X							
<i>Reticulofenestra dictyoda</i>		?					X						X
<i>R. umbilica</i>				X			X	X					
<i>Rhabdosphaera</i> sp.					X								
<i>Sphenolithus moriformis</i>								X		X			
<i>S. orphanknolli</i>								X					
<i>S. predistentus</i>										X			
<i>S. aff. S. radians</i>								X					
<i>Thoracosphaera saxea</i>								X					
<i>T. spp.</i>											X		
<i>Zygodiscus sigmoides</i>	X												
<i>Zygodiscus</i> sp.			X					X					X
Zone assignment	NP5	NP12-13	NP14-15	NP14 ² -15 ²	NP15-16	n.d.	NP16-17	NP16-17	n.d.	NP19-20	n.d.	n.d.	NP15

¹Locatelli Formation.

²Butano Sandstone.

³Twobar Shale Member of San Lorenzo Formation.

68CB202 from the Bear Creek area shows that the upper part of the Butano Sandstone is of middle Eocene (NP16 and NP17 Zones) age.

Collection 76CB1451A from the Twobar Shale Member of the San Lorenzo Formation along the San Lorenzo River indicates that the lower part of this member is of middle Eocene (NP15 Zone) age. This

collection also contains *Neochiastozygus dubius* fragments that may be reworked from older strata.

The middle part of the Twobar Shale Member, represented by collection 68CB205 from the Bear Creek area, is of late Eocene (NP19 and NP20 Zones) age. Thus the boundary between the middle and late Eocene is probably within the Twobar Shale Member.

DEFINITION OF THE LODO FORMATION

By EARL E. BRABB

ABSTRACT

The section along Lodo Gulch, described in general terms by White (1940) and Martin (1943, section I-S), is the type section of the Lodo Formation. Sections along the first gulch south of Lodo Gulch (section A of Israelsky, 1951 and the so-called Lodo Gulch section of Schoellhamer and Kinney, 1953), along the second gulch south of Lodo Gulch (section I-X of Martin, 1943 and section B of Israelsky, 1951), and about 700 m north of Lodo Gulch (section C of Brabb, Clark, and Throckmorton, 1977) are reference sections.

DISCUSSION

The name Lodo, Spanish for mud, was proposed in an abstract by R. T. White (1938) for a small gulch in the Tumey Hills, about 75 km west of Fresno, Calif. (figs. 1 and 16, this report); the exact location of Lodo Gulch is shown on maps by Martin (1943, fig. 1) and Israelsky (1951, fig. 2). In this same abstract, White also proposed the name Lodo Formation for 350 m of predominantly claystone that rests, according to White, disconformably on the Moreno Shale and is overlain conformably by the Domengine Sandstone.

Two years later, White (1940, fig. 1), in an expanded report, showed a generalized stratigraphic column for the Lodo Formation in Lodo Gulch. He used the term "type locality" for this section in both his 1938 abstract and his report, and it is clear from his descriptions that he intended this section to be the type section for the formation. This definition is accepted here as the type section of the Lodo (section TS, fig. 16, this report). Martin (1943), in a report that described foraminifers from samples collected by White, provided a figure entitled, "Sketch map of the Lodo Formation in the type area." Taken literally, the map indicates that the type area includes all of the Lodo Formation in secs. 20, 28, and 29, T. 15 S., R. 12 E. This definition of the type area is accepted here (see fig. 16, this report).

Some confusion has been introduced by authors who measured sections in the type area of the Lodo

Formation but not in the type section along Lodo Gulch. According to the International Stratigraphic Guide (Hedberg, 1976, p. 26) and the Code of Stratigraphic Nomenclature (American Commission on Stratigraphic Nomenclature, 1961, p. 654), sections subsidiary to the type section (holostratotype) are reference sections. Martin (1943, fig. 1), for example, showed two sections (I-S and I-X) as a composite type section of the Lodo. This definition is not accepted here. Martin's section I-S is in Lodo Gulch and is the type section (TS, fig. 16, this report), but section I-X (section B, fig. 16, this report) is in the second gulch south of Lodo Gulch and is therefore here designated a reference section.

The label "Lodo Gulch section" on the map by Schoellhamer and Kinney (1953) is misleading. This section (section A, fig. 16, this report) is not in Lodo Gulch, but is in the first gulch south of Lodo Gulch. The label on Schoellhamer and Kinney's map led me to conclude wrongly that Israelsky's (1951) section A (section A, fig. 16, this report) is along Lodo Gulch. Thus, section A on the stratigraphic column of Brabb, Clark, and Throckmorton (1977, fig. 42) is a composite of samples collected by me along Lodo Gulch, the type section, and samples described by Israelsky from section A, the first gulch south of Lodo Gulch. Section A of Israelsky is in the type area but is not the type section. The "Lodo Gulch section" of Schoellhamer and Kinney (1953) and section A of Israelsky (1951) are the same as section A of this report (fig. 16), which is here designated the principal reference section.

Section B of Israelsky (1951), which gives supplementary information on the lower part of the Lodo, is the same as White's (1943) section I-X and section B of this report; it is a reference section. Section C of Brabb, Clark, and Throckmorton (1977, p. 105-109) also gives supplementary information on the lower part of the formation; it is the same as section C of

68CB202 from the Bear Creek area shows that the upper part of the Butano Sandstone is of middle Eocene (NP16 and NP17 Zones) age.

Collection 76CB1451A from the Twobar Shale Member of the San Lorenzo Formation along the San Lorenzo River indicates that the lower part of this member is of middle Eocene (NP15 Zone) age. This

collection also contains *Neochiastozygus dubius* fragments that may be reworked from older strata.

The middle part of the Twobar Shale Member, represented by collection 68CB205 from the Bear Creek area, is of late Eocene (NP19 and NP20 Zones) age. Thus the boundary between the middle and late Eocene is probably within the Twobar Shale Member.

DEFINITION OF THE LODO FORMATION

By EARL E. BRABB

ABSTRACT

The section along Lodo Gulch, described in general terms by White (1940) and Martin (1943, section I-S), is the type section of the Lodo Formation. Sections along the first gulch south of Lodo Gulch (section A of Israelsky, 1951 and the so-called Lodo Gulch section of Schoellhamer and Kinney, 1953), along the second gulch south of Lodo Gulch (section I-X of Martin, 1943 and section B of Israelsky, 1951), and about 700 m north of Lodo Gulch (section C of Brabb, Clark, and Throckmorton, 1977) are reference sections.

DISCUSSION

The name Lodo, Spanish for mud, was proposed in an abstract by R. T. White (1938) for a small gulch in the Tumey Hills, about 75 km west of Fresno, Calif. (figs. 1 and 16, this report); the exact location of Lodo Gulch is shown on maps by Martin (1943, fig. 1) and Israelsky (1951, fig. 2). In this same abstract, White also proposed the name Lodo Formation for 350 m of predominantly claystone that rests, according to White, disconformably on the Moreno Shale and is overlain conformably by the Domengine Sandstone.

Two years later, White (1940, fig. 1), in an expanded report, showed a generalized stratigraphic column for the Lodo Formation in Lodo Gulch. He used the term "type locality" for this section in both his 1938 abstract and his report, and it is clear from his descriptions that he intended this section to be the type section for the formation. This definition is accepted here as the type section of the Lodo (section TS, fig. 16, this report). Martin (1943), in a report that described foraminifers from samples collected by White, provided a figure entitled, "Sketch map of the Lodo Formation in the type area." Taken literally, the map indicates that the type area includes all of the Lodo Formation in secs. 20, 28, and 29, T. 15 S., R. 12 E. This definition of the type area is accepted here (see fig. 16, this report).

Some confusion has been introduced by authors who measured sections in the type area of the Lodo

Formation but not in the type section along Lodo Gulch. According to the International Stratigraphic Guide (Hedberg, 1976, p. 26) and the Code of Stratigraphic Nomenclature (American Commission on Stratigraphic Nomenclature, 1961, p. 654), sections subsidiary to the type section (holostratotype) are reference sections. Martin (1943, fig. 1), for example, showed two sections (I-S and I-X) as a composite type section of the Lodo. This definition is not accepted here. Martin's section I-S is in Lodo Gulch and is the type section (TS, fig. 16, this report), but section I-X (section B, fig. 16, this report) is in the second gulch south of Lodo Gulch and is therefore here designated a reference section.

The label "Lodo Gulch section" on the map by Schoellhamer and Kinney (1953) is misleading. This section (section A, fig. 16, this report) is not in Lodo Gulch, but is in the first gulch south of Lodo Gulch. The label on Schoellhamer and Kinney's map led me to conclude wrongly that Israelsky's (1951) section A (section A, fig. 16, this report) is along Lodo Gulch. Thus, section A on the stratigraphic column of Brabb, Clark, and Throckmorton (1977, fig. 42) is a composite of samples collected by me along Lodo Gulch, the type section, and samples described by Israelsky from section A, the first gulch south of Lodo Gulch. Section A of Israelsky is in the type area but is not the type section. The "Lodo Gulch section" of Schoellhamer and Kinney (1953) and section A of Israelsky (1951) are the same as section A of this report (fig. 16), which is here designated the principal reference section.

Section B of Israelsky (1951), which gives supplementary information on the lower part of the Lodo, is the same as White's (1943) section I-X and section B of this report; it is a reference section. Section C of Brabb, Clark, and Throckmorton (1977, p. 105-109) also gives supplementary information on the lower part of the formation; it is the same as section C of

LARGE FORAMINIFERS OF EOCENE AGE FROM THE COAST RANGES OF CALIFORNIA

By ALPHONSE BLONDEAU and EARL E. BRABB

ABSTRACT

Large foraminifers from an unnamed limestone of Eocene age and from reworked boulders within the unconformably overlying Temblor Formation of Miocene age in the vicinity of San Jose, from the Domingue Formation of early and middle Eocene age near Mount Diablo, and from the Avenal Sandstone of early and middle Eocene age in the Devils Den area, all in California include *Nummulites willcoxi* (Heilprin), *N. striatoreticulatus* (L. Rutten), *Euconuloides lopeztrigoi* (Palmer), *Amphistegina parvula* (Cushman), *Discocyclina* (*Discocyclina*) *marginata* (Cushman), *Pseudophragmina* (*Proporocyclina*) *clarki* (Cushman), *P. (P.) teres* (Cole and Gravell), *Asterocyclina aster* (Woodring), and *Eofabiana grahami* (Kupper). These species, which collectively have been reported previously from other areas in California, from Oregon and Washington, and from the Gulf Coast and Caribbean region, indicate a middle Eocene age.

INTRODUCTION

Many reports have been published on large foraminifers of Eocene age from the Pacific Coast region, but almost all of these date from the 1920's and 1930's. (See for example the reports by Palmer, 1929, Nelson and Schenck, 1930, Woodring, 1930, Keenan, 1932, and Schenck, 1933). The timing of these reports is significant because the California Eocene zonations based on benthic foraminifers were not published until the 1940's (Laiming, 1943) and 1950's (Mallory, 1959), and the application of zonations based on planktic foraminifers have only been made in the last few years. Thus, the stratigraphic relation between the large foraminifers from the Pacific Coast and those used for interregional and intercontinental correlation is not well established. Moreover, knowledge about the evolution and distribution of large foraminifers has increased so much in the past 30 years that the time is opportune for a new look at the Pacific Coast assemblages.

The fossils described in this study were all collected east of the San Andreas fault from the Mount Diablo to the Devils Den areas (fig. 1). In the conceptual scheme of Nilsen and Clarke (1975), the large foraminifers were deposited on the ocean floor in a linear basin along the margins of the continent. The sandstone and shale that make up most of the Eocene sequence were derived from the Sierra Nevada region and narrow islands within the linear basin. The sands were probably displaced downslope from upper

neritic depths to a bathyal environment, carrying the larger foraminifers with them.

The specimens were nearly all cemented together by calcite, so that it was necessary to study them in thin sections that could not always be oriented in the most desirable direction. The study was further complicated by the great variety and plasticity of forms that create nomenclatural, classification, and correlation problems to an extent generally unknown in Europe.

Acknowledgments.—We are grateful to Erik Osburn, who provided several collections of large foraminifers from the Sveadal area. A. D. Warren kindly pointed out the occurrence of large foraminifers in the aqueduct section near Devils Den and provided several unattached specimens for study. J. Butterlin provided much useful information on the stratigraphic occurrence of similar large foraminifers in the Caribbean region. K. N. Sachs and René Herb reviewed the manuscript and provided many helpful suggestions.

STRATIGRAPHY

DEVILS DEN AREA

The Devils Den area has a nearly complete section of Paleogene rocks that is easily accessible, moderately well exposed, and not too complicated structurally. Unfortunately, calcareous microfossils have been leached out of most exposures. In the late 1960's, however, the Barrenda Mesa Water District made a cut for a canal that exposed rocks containing well-preserved calcareous microfossils. The nannoplankton are described by Warren (this volume) and the smaller foraminifers by Berggren and Aubert (this volume). We report on a single bed containing large foraminifers. The rock unit nomenclature is from Dibblee (1973a, fig. 6) and the biostratigraphy is from Warren (this volume).

The large foraminifers, collections 76CB1633, 1634, and 1641 (table 5) are from the Avenal Sandstone. The Avenal Sandstone contains, in addition to the large foraminifers, abundant worm tubes called "*Spiroglyphus*" (*Rotularia*) in several reports. The geographic and geologic position of the sandstone and the fossils collected from it are provided by Brabb, Clark, and Throckmorton (1977, p. 94-103).

[illegible]




FIGURE 17.—Localities where large foraminifera were collected in the San Francisco Bay region.

SVEADAL AREA

In the Sveadal area near Loma Prieta (figs. 17 and 19), Osbun (1975) discovered large foraminifers (table 5, collections 49-6 and 1-1) in limestone boulders in a basal conglomerate of the Temblor Formation of Miocene age. These boulders were presumably derived from the erosion of the limestone in the vicinity of localities 71CB983 and 71CB971 near San Jose (fig. 18).

AGE

The large foraminifers from the Devils Den area

are in rocks correlated by Warren (this volume) on the basis of calcareous nannoplankton with the Penutian Stage of Mallory (1959) and the *Discoaster lodoensis* Zone of Bukry (1973). Both these correlations suggest an early Eocene age (Brabb and others, 1977, fig. 2). These correlations are supported by the occurrence of planktic foraminifers of Zone P7 (lower Eocene) stratigraphically close to the beds with large foraminifers in the San Jose area. On the other hand, the large foraminifers (table 5) have age ranges that are much more suggestive of the middle Eocene. Of course, the large foraminifers could be early Eocene in some areas and middle Eocene in others, but we are in-




FIGURE 18.—Geologic map of the southern San Jose area showing localities where large foraminifers were collected. Bases from Los Gatos and Santa Teresa Hills 7.5-minute quadrangles. Geology generalized from Bailey and Everhart (1964).

clined to view these assemblages as closely related and probably the same age, middle Eocene.




FIGURE 19.—Geologic map and stratigraphic column for Sveadal area. Geology from Osbun (1975).

SYSTEMATIC PALEONTOLOGY

Nummulites willcoxi Heilprin, 1882

Plate 6, figures 6, 7, 10

Operculinoides willcoxi (Heilprin). Cole, 1958, pl. 33, figs. 3-12.

Camerina willcoxi (Heilprin). Cole and Applin, 1964, pl. 4, figs. 9, 14-16.

Nummulites willcoxi Heilprin. Butterlin, 1966, pl. XIV, figs. 1-10.

Nummulites willcoxi Heilprin. Butterlin, 1971, pl. VI, figs. 1-8.

Description. Small lenticular species, diameter to thickness 3.6:0.6 mm; globular in juvenile stage with diameter to thickness 1.6:0.65 mm; tends to become flat by growth from the foot of the spire; more loosely coiled spire in outer parts of the test; thin septa curved at top; diameter of the protoconch approximately 100 μ m.

Young specimens cannot be distinguished from *N. trinitatis* (Nuttall). Older specimens have a somewhat flattened marginal area in contrast to the rounded marginal area of *N. trinitatis*.

Distribution. California, locality 71CB983C and EB633 near San Jose; Florida, Mexico, Caribbean, Panama, and Columbia.

Range. Top of middle Eocene to base of upper Eocene.

Nummulites striatoreticulatus L. Rutten, 1928

Plate 6, figure 9

Camerina striatoreticulata (L. Rutten). Cole, 1958, pl. 32, Figs. 1-16

Nummulites striatoreticulata L. Rutten. Butterlin, 1966, fig. 5.

Description. Globular species from 1 to 3 mm in diameter, sinuous septal filament, more or less granulated, thick spiral laminae loosely coiled. Septa straight and inclined in lower part, curved in upper part. Small protoconch of megalospheric form.

Distribution. California, locality 71CB983C near San Jose; Cuba, Haiti, Mexico, Jamaica, Costa Rica, and Colombia.

Range. Top of middle Eocene to basal upper Eocene.

Nummulites sp.

Description. Small, lenticular, globular forms 1 mm in diameter. Lack the granular form of *N. striatoreticulatus*. More globular than *N. willcoxi*. Unfortunately, they are only observable in thin sections.

Distribution. California, locality 71CB971B near San Jose.

Family AMPHISTEGINIDAE

Genus *Eoconuloides* Cole and Bermudez, 1944*Eoconuloides lopeztrigoi* (D. K. Palmer, 1934)

Plate 6, figures 3, 13

Amphistegina lopeztrigoi D. K. Palmer, 1934, pl. 5, fig. 6-8.*Amphistegina lopeztrigoi* D. K. Palmer. Barker and Grimsdale, 1936, pl. 30, fig. 2; pl. 32, fig. 1-3; pl. 34, fig. 1; pl. 38, fig. 3.*Amphistegina lopeztrigoi* D. K. Palmer. Levin, 1957, pl. 4, fig. 8-9, 13-14.*Amphistegina lopeztrigoi* D. K. Palmer. Butterlin, 1966, pl. XIV, fig. 14.*Amphistegina lopeztrigoi* D. K. Palmer. Butterlin, 1968, pl. III, fig. 8.

Description. Small thin spiral lamina, slightly trochoidal, showing pillars in relief, generally well marked.

Distribution. California, localities EB633 and 71CB971B near San Jose; Caribbean, Columbia, Venezuela.

Range. From the base to the summit of the middle Eocene and basal upper Eocene.

Remarks. Butterlin (1970) showed that *Eoconuloides parvulus* evolved into *Eoconuloides lopeztrigoi* and *Eoconuloides wellsii*.

Eoconuloides wellsii Cole and Bermudez, 1944

Plate 6, figure 4

Description. Test conical with flat base, apex bluntly rounded; peripheral angle subacute; surface smooth. Specimen: height, 0.8 mm to 1.6 mm; diameter of base: 0.96 mm to 1.20 mm. On slightly oblique horizontal sections, the final chambers are subdivided into chamberlets in the peripheral section; axial sections show the trochoid character of test; the spiral wall has differences in thickness, the thicker walls having irregularly developed pillars; the thin peripheral walls appear to lack pillars.

Range. Lower Eocene to middle Eocene.

Amphistegina parvula (Cushman, 1919)

Plate 6, figure 1, 2, 14

Nummulites parvula Cushman, 1919, pl. 4, fig. 3-6.*Amphistegina parvula* (Cushman). Cole, 1958, pl. 25, fig. 17-19.*Amphistegina cf. parvula* (Cushman). Butterlin, 1968, pl. II, fig. 8.*Amphistegina parvula* (Cushman). Butterlin, 1970, pl. 3, fig. 5-6.

Description. *Amphistegina* with slightly asymmetrical axial cut. Has very thick walls and very corroded and barely visible pillars. Septa inclined and curved. In a good equatorial cut, could be confused with a very small nummulite. An axial cut shows the distinction.

Distribution. California, localities EB633 and 76CB971B near San Jose and Osburn 49-6 near Sveadal; Caribbean region (abundant).

Range. Lower Eocene to lower part of upper Eocene.

Remarks. All collections observed are rich in *Amphistegina*. New species are probably present. Uncemented material and separate specimens are necessary for a detailed study.

Family DISCOCYCLINIDAE

The classification of Discocyclinidae is a complex problem for American species. The classifications are numerous and varied: Vaughan and Cole (1941), Bronnimann (1945), Cole (1948), and Caudri (1972). We accept Cole's classification for our work.

Discocyclina (Discocyclina) marginata (Cushman, 1919)*Orthophragmina marginata* Cushman, 1919, pl. 1, fig. 2; pl. 2, fig. 1.*Orthophragmina marginata* Cushman, 1920, pl. IX, fig. 1-2.*Discocyclina (Discocyclina) marginata* (Cushman). Cole and Gravell, 1952, pl. 93, fig. 1-9; pl. 94, fig. 1-8; pl. 95, fig. 7-8.*Discocyclina (Discocyclina) marginata* (Cushman). Cole and Applin, 1964, pl. 10, fig. 1-8.

Description. Lenticular species, inflated, diameter 10-14 mm, equatorial chambers higher than wide ($h/w = 1.5$) and very small. Thirty to forty rows of lateral chambers are very compressed and poorly visible.

Distribution. California, localities 71CB983C, 71CB978B, and EB633 near San Jose; Caribbean region (Cuba, St. Bartholomew), Florida, Georgia.

Range. Middle Eocene.

Remarks. Previously described in California as *D. californica* (Schenck, 1929, pl. 28, fig. 5; pl. 29, fig. 1-3; pl. 30, fig. 2-3), according to Cole and Applin (1964).

Pseudophragmina (Proporocyclina) flintensis (Cushman)

Plate 6, figure 12

Orthophragmina flintensis Cushman, 1917, pl. 40, fig. 1-2.

Orthophragmina flintensis Cushman, 1919, pl. IX, fig. 3-6.

Proporocyclina flintensis (Cushman). Cole, 1958, pl. 50, 51, 52, figs. 1-2.

Pseudophragmina (Pseudophragmina) flintensis (Cushman). Butterlin, 1971, pl. IV, fig. 8.

Proporocyclina flintensis (Cushman). Caudri, 1972, pl. 1, fig. 4.

Description. Test diameter 5 mm, circular, compressed and ornate with granules. Narrow equatorial chambers, radially lengthened to the periphery. Complete radial septa, almost rectilinear. Small and narrow lateral chambers.

Distribution. California, localities 71CB983C, 71CB978B, and EB633 near San Jose, EB611 near Mount Diablo (see also Cole, 1958) and Orocopia Mountains (Cole, 1958); Caribbean (with *Lepidocyclina antillea*), Nicaragua, Georgia, Texas, Florida, Panama.

Range. Top of middle Eocene to upper Eocene.

Pseudophragmina (Proporocyclina) clarki (Cushman)

Plate 6, figure 17

Orthophragmina clarki Cushman, 1920, pl. VII, fig. 4-5.

Discocyclina clarki (Cushman). Schenck, 1929, pl. 27, fig. 1, 2, 5, and text fig. 7.

Discocyclina clarki (Cushman). Keenan, 1932, pl. 4, fig. 1-2.

Pseudophragmina (Proporocyclina) clarki (Cushman). Cole, 1958, pl. 52, fig. 3-11.

Pseudophragmina (Proporocyclina) clarki (Cushman). Cole and Applin, 1964, pl. 9, fig. 1-4.

Description. Circular flat test, 3-6 mm in diameter (larger than that of *Pseudophragmina (Proporocyclina) flintensis*), with pillars and central granules. Large embryonic apparatus. In equatorial section, the length and width of the embryonic apparatus are 0.2 mm. In axial section, the length is 0.2 mm and the height is 0.175 mm. Equatorial chambers 1.5 to 2 times higher than wide, lateral chambers narrow and very elongated.

Distribution. California, localities 71CB983C, EB633, and 71CB971B near San Jose. Type locality: Domengine Creek near Mount Diablo, California; also Oregon, Washington (Berthiaume, 1938), Mexico, Peru, and Florida.

Range. Middle Eocene.

Pseudophragmina (Proporocyclina) teres Cole and Gravell, 1952

Pseudophragmina (Proporocyclina) teres Cole and Gravell, 1952, pl. 101, fig. 1-8.

Pseudophragmina (Proporocyclina) teres Cole and Gravell, 1952. Cole and Applin, 1964, pl. 8, fig. 1-7.

Description. Circular species, flat, inflated in the center, diameter 5-6 mm, with pillars and central granules. Equatorial chambers squared toward the center, elongated toward the periphery (h/w approx. 2). Lateral chambers with thick borders and roofs that are flat and barely visible.

Distribution. California, locality 71CB971B near San Jose; Caribbean, Cuba, Florida.

Range. Middle Eocene.

Remarks. Probably present in other samples.

Genus *Asterocyclina* Gumbel, 1870

Asterocyclina aster (Woodring, 1930)

Plate 6, figure 11

Actinocyclina aster Woodring, 1930, fig. 3-6; pl. 16, fig. 1-4; pl. 17, fig. 1-2.

Description. Very flat species with 5 to 13 radiating enlargements. Diameter 5 to 11 mm.

Distribution. California, locality EB611 near Mount Diablo, EB633, 71CB983C, and 71CB971B near San Jose; Osbun 1-1 and 49-6 near Sveadal. Type locality is in Santa Ynez Range, Santa Barbara County.

Range. Upper Eocene (Woodring, 1930), lower and middle Eocene (Berthiaume, 1938; Cole, 1958), middle Eocene (Cole and Gravell, 1952).

Remarks. Cole (1958) put *A. aster* in synonymy with *A. penonensis* of Cole and Gravell (1952). Butterlin (1970) disagrees with this synonymy and lists several reasons for considering the two forms as separate species. Present in all the samples.

Family CYMBALOPORIDAE

Genus *Eofabiana* Kupper, 1955

Eofabiana grahami Kupper, 1955

Plate 6, figures 8, 16

Eofabiana grahami Kupper, 1955, pl. 19, figs. 1-7.

Description. Conical test, primary (first) chambers poorly developed and spirally arranged, not divided into chamberlets. Apertures on the umbilical side.

Distribution. California and the Caribbean.

Range. Lower to middle Eocene.

LOCALITY DESCRIPTIONS

SAN JOSE AREA

Locality 71CB983C, lat 37°13.44'N., long 121°53.29'W., Los Gatos 7.5-minute quadrangle, 1953 edition. Unnamed sedimentary rocks of Eocene age. This locality is within the same limestone lentil and very close geographically to LSJU 309, from which the holotype of *Discocyclina californica* Schenck, 1929 was obtained. The general geology and a more detailed description of the lithology are provided by Bailey and Everhart (1964, p. 69-71 and pl. 1). Collected by Earl Brabb, 1971.

Locality 71CB971B, lat 37°13.25'N., long 121°47.14'W., Santa Teresa Hills 7.5-minute quadrangle, 1953 edition. Unnamed sedimentary rocks of Eocene age. Float pieces of limestone from a small abandoned limestone quarry roughly 0.6 km east of the portal of the Bernal Mine. Collected by Earl Brabb, 1971.

Locality EB633, lat 37°12.9'N., long 121°47.5'W., Santa Teresa Hills 7.5-minute quadrangle, 1953 edition. Unnamed sedimentary rocks of Eocene age. Float pieces of limestone collected by Edgar Bailey in 1948.

SVEADAL AREA

Locality Osbun 1-1, lat 37°05.47'N., long 121°45.03'W., Loma Prieta 7.5-minute quadrangle, 1955 edition. Fossils from conglomerate at or near the base of the Temblor Formation of Miocene age. From small outcrop on north side of road to Sveadal. Collected by E. D. Osbun, 1971. See geologic map by McLaughlin and others (1971); additional maps and a description of the rocks provided in Osbun (1975).

Locality Osbun 49-3, lat 37°06.27'N., long 121°45.68'W., Loma Prieta 7.5-minute quadrangle. Fossils from conglomerate at or near the base of the Temblor Formation of Miocene age. From nose of hill at about 1160 ft (354 m) elevation. Collected by E. D. Osbun, 1971.

Locality Osbun 49-6, lat 37°06.31'N., long

121°45.5'W., Loma Prieta 7.5-minute quadrangle. Fossils from conglomerate at or near the base of the Temblor Formation of Miocene age. From small knob along ridge crest, elevation about 1170 ft (357 m). Collected by E. D. Osbun, 1971.

MOUNT DIABLO AREA

Locality EB611, lat 37°51.35'N., long 121°56'W., Diablo 7.5-minute quadrangle, 1953 edition. Southwest side of road from Mount Diablo State Park headquarters to Junction Camp. From glauconitic limestone near the base of the Domengine Formation where it rests unconformably on shale of Late Cretaceous age. Mapped as lower siltstone and claystone member of Domengine Sandstone by Brabb, Sonneman, and Switzer (1971). Collected by Earl Brabb, 1963. Not exposed in 1976.

DEVILS DEN AREA

Locality 76CB1633, lat 35°42.4'N., long 120°01.1'W., Sawtooth Ridge 7.5-minute quadrangle, 1961 edition (photorevised 1973). Gredal Shale Member of Kreyenhagen Formation. Along north bank of cut for aqueduct, about 20 m east of dirt access road, about 12 m stratigraphically below base of red bed unit, from 50-cm-thick coarse-grained sandstone bed.

Locality 76CB1634, lat 35°42.3'N., long 121°01.3'W. Gredal Shale Member of Kreyenhagen Formation. Along north bank of cut for aqueduct, about 60 m east of two metal posts and 7 m east of telephone box. Same stratigraphic level as 76CB1633 but from part of section duplicated by faulting.

Locality 76CB1641, lat 35°43.8'N., long 121°01.3'W. Gredal Shale Member of Kreyenhagen Formation. About 600 m N.63 W. from BM611 from sandstone rubble on hillside. Probably from same stratigraphic level as 76CB1633 and 76CB1634, but sandstone at this locality appears to be several meters thick.

Locality 76CB1671, about 4 m stratigraphically below 76CB1634. Gredal Shale Member of Kreyenhagen Formation. Sample collected by A. D. Warren.

EOCENE TO MIOCENE CALCAREOUS PLANKTON FROM THE SANTA CRUZ MOUNTAINS AND NORTHERN SANTA LUCIA RANGE, CALIFORNIA

By RICHARD Z. POORE and DAVID BUKRY

ABSTRACT

Stratigraphically diagnostic calcareous plankton (coccoliths and planktic foraminifers) were recovered from two measured sections (Zayante Creek and San Lorenzo River sections) in the Santa Cruz Mountains, one section (Año Nuevo section) along the adjacent coast, and several samples from the Church Creek Formation in the northern Santa Lucia Range.

Samples from below the uppermost sheared part of the Vaqueros(?) Formation of Hall, Jones, and Brooks (1959) as exposed on the coast at the Año Nuevo section yield coccolith assemblages referable to the upper Oligocene *Sphenolithus cipoensis* Zone and the lower Miocene *Triquetrorhabdulus carinatus* Zone. Samples from the sheared upper part of the Vaqueros(?) Formation contain coccoliths that are indicative of the Miocene *Helicosphaera ampliaperta* Zone or the *Sphenolithus heteromorphus* Zone.

Late Oligocene coccoliths were recovered from the Vaqueros Sandstone at Zayante Creek in the Santa Cruz Mountains. The sparse low-diversity assemblages in most samples could not be assigned to specific zones.

Diverse calcareous plankton assemblages were recovered from samples from the San Lorenzo River section. Coccoliths from the upper part of the Butano Sandstone and lower part of the Twobar Shale Member of the San Lorenzo Formation are referable, respectively, to the *Discoaster bifax* Subzone and *Discoaster saipanensis* Subzone of the middle Eocene *reticulofenestra umbilica* Zone. Planktic foraminifer assemblages from the same samples correlate with middle Eocene Zones P 13 to P 14. Coccoliths from the upper part of the Twobar Shale Member are assigned to the upper Eocene *Discoaster barbadiensis* Zone. Samples from this same interval yield planktic foraminifers assigned to upper Eocene Zones P 15 to P 16. A sample from the overlying Rices Mudstone Member of the San Lorenzo Formation contains a coccolith assemblage referable to the Oligocene *Sphenolithus distentus* Zone. Planktic foraminifer assemblages from this level are assigned to Oligocene zonal interval P 19 - P 20.

Planktic foraminifers in several samples from the Church Creek Formation, northern Santa Lucia Range, are correlated with upper Eocene Zones P 16 to P 17.

INTRODUCTION

A number of stratigraphic sections in and around the Santa Cruz Mountains and the Santa Lucia Range were examined for coccoliths and planktic foraminifers as part of a field conference and meeting held in Menlo Park, California, by the International Commission on Paleogene Stratigraphy. In this report, data from the Zayante Creek and the San Lorenzo River sections of the Santa Cruz Mountains

are presented along with data from the coastal Año Nuevo section (fig. 20). In addition, the occurrence of planktic foraminifers in several samples from the Church Creek Formation in the Church Creek area of the northern Santa Lucia Range are discussed. Brabb, Bukry, and Pierce (1971) and Brabb, Clark, and Throckmorton (1977) give detailed location information for sections and most samples that were examined for this report. Locality data for samples not mentioned in the above reports are given at the end of this report. The zonation of Blow (1969) is employed for planktic foraminifers, and the zonation of Bukry (1975) is used for coccoliths.

Correlation of biostratigraphic zones to subdivisions of the Cenozoic follows Hardenbol and Berggren (1978) for the Eocene and Oligocene, and Ryan and others (1975) for the Miocene.

Acknowledgments.—We thank Earl Brabb, John Barron, and Kristin McDougall for comments concerning this manuscript.




FIGURE 20.—Index map showing general location of sections. Map from Brabb, Bukry, and Pierce (1971). Section A is in the Año Nuevo area; section B, along the San Lorenzo River, and section C, along Zayante Creek.




FIGURE 21.—Generalized stratigraphic column of the Año Nuevo section and location of samples yielding calcareous plankton. Modified from Brabb, Clark, and Throckmorton (1977, fig. 31). Position of benthic foraminiferal stages from McDougall (this volume). Position of Oligocene-Miocene boundary estimated from our samples.

TABLE 6.—Distribution of coccoliths and resultant zone assignments for the Año Nuevo section.

[Species shown by asterisk are considered to be reworked]

Species Sample number	Campylosphaera eodella* Chiasmolithus sp. Coccolithus spp. Cyclicargolithus abisectus C. floridanus Dictyococites bisectus D. scrippsae Discoaster deflandrei D. divaricatus D. exilis D. lodoensis* D. variabilis Discoaster spp. Discolithina multipora D. plana s. ampl. Helicosphaera carteri H. euphratis H. intermedia Helicosphaera spp. Reticulofenestra spp. Sphenolithus belemnos S. conicus S. dissimilis S. heteromorphus S. moriformis S. radians* Tessellolithus sp. Tribrachiatus orthostylus* Triquetrorhabdulus carinatus Trochoaster sp. Zygrhablithus sp.																							Coccolith zone of Bukry (1975)	Age																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
76CB1541B Mf4668	×	×	×	aff.	aff.		×	×	×											×	aff.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				

AÑO NUEVO SECTION

A composite section for the Año Nuevo area with the location of samples yielding stratigraphically diagnostic assemblages of calcareous plankton is shown in figure 21. Coccoliths were found throughout the Vaqueros(?) Formation of Hall, Jones, and Brooks (1959) in this section (table 6). Planktic foraminifers are sparse and poorly preserved in samples Mf4664 and Mf4665, but several species were identified (table 7). Planktic foraminifers are common to abundant in samples Mf4667 and Mf4668 but are so poorly pre-

TABLE 7.—Distribution of planktic foraminifers in samples from the Año Nuevo section

Taxon	Sample number	
	76CB1532 (Mf4665)	76CB1531 (Mf4664)
<i>Globigerina angustumbilicata</i> Bolli	---	x
<i>G. juvenilis</i> Bolli	x	x
<i>G. praebulloides</i> Blow	x	x
<i>G. woodi</i> Jenkins	x	x
<i>G. tripartita</i> Koch	---	x
<i>Globorotalia nana</i> Bolli	x	x
<i>G. munda</i> Jenkins	aff.	aff.
<i>Catapsydrax unicavus</i> Bolli, Loeblich and Tappan	---	x
<i>Globigerinita praestainforthi</i> Blow	aff.	aff.
<i>Globorotaloides suteri</i> Bolli	x	x

served and distorted that no confident identifications were possible. The ranges of several important taxa detected in our samples are shown in figure 22.

The coccoliths *Triquetrorhabdulus carinatus* or *T. sp. cf. T. carinatus* and *Dictyococcites bisectus* in samples Mf1376 through Mf4664 (table 6) are secondary zonal guide taxa that indicate assignment of this interval to the upper Oligocene *Sphenolithus ciperoensis* Zone. The occurrence of *Globorotalia nana* in this interval is compatible with an upper Oligocene assignment. The next higher sample (Mf2188) lacks *Dictyococcites bisectus* and contains *Helicosphaera carteri*, a form restricted to the Miocene and younger strata. We assign this assemblage to the *Triquetrorhabdulus carinatus* Zone and place the Oligocene-Miocene boundary (see fig. 21) between sample Mf2188 and sample Mf4664. Samples Mf2190, Mf4667, and Mf4668 from the sheared section of the Vaqueros(?) Formation, just below the Monterey Formation, contain few to common *Sphenolithus heteromorphus*, which indicates correlation with the lower and middle Miocene *Helicosphaera ampliaperta* Zone or the lower middle Miocene *Sphenolithus heteromorphus* Zone. Thus this sheared zone is near the lower Miocene-middle Miocene boundary. Our age interpretations when compared to the California benthic

SERIES	FORMATION	SAMPLE	TAXA
Pliocene	Purisima Formation		
Miocene	Monterey Formation		
	Sheared section		
Oligocene	Vaqueros(?) Formation		

FIGURE 22.—Ranges of selected calcareous plankton in the Año Nuevo section.

foraminiferal stage determinations for this section (McDougall, this volume; Brabb and others, 1977) suggest that the Zemorrian Stage-Saucesian Stage boundary is within the uppermost Oligocene or lower Miocene and the Saucesian Stage-Relizian Stage boundary is within the lower Miocene or middle Miocene.

ZAYANTE CREEK SECTION

The stratigraphic columns and locations of samples yielding coccoliths from the Zayante Creek section (north limb) are shown on figure 23. Coccolith assemblages from the upper part of the Rices Mudstone Member of the San Lorenzo Formation cannot be assigned to a specific zone or limited zonal interval but are probably of late Oligocene age (fig. 23, table 8). Meager coccolith assemblages from the overlying Vaqueros Sandstone suggest correlation with the upper Oligocene *Sphenolithus distentus* Zone or *Sphenolithus ciperoensis* Zone. An assemblage from the lower part of the Lambert Shale (JC 61-11d) is

tentatively considered to be of Miocene age because with the exception of a few fragmented specimens, *Dictyococcites bisectus* is absent whereas it is fairly common in samples from the underlying Vaqueros Sandstone.

Only sparse, juvenile planktic foraminifers that could not be identified with certainty were found in the Zayante Creek Section.

SAN LORENZO RIVER SECTION

The stratigraphic column for the San Lorenzo River section and locations of samples containing calcareous plankton are shown on figure 24. Data on planktic foraminifers from this section were reported by Poore and Brabb (1977). Coccoliths in samples from the upper part of the Butano Sandstone and the Twobar Shale Member of the San Lorenzo Formation are listed in table 9. Ranges of selected planktic foraminifers and coccoliths in the San Lorenzo River section are shown on figure 25. Poore and Brabb's (1977) rationale for planktic foraminifer zone assignments are briefly summarized below.

The occurrence of *Globorotaloides suteri* in Mf3301 (upper part of Butano Sandstone) and *Globorotalia cerroazulensis cerroazulensis* in Mf1350 and Mf3304 (lower part Twobar Shale Member of San Lorenzo Formation) indicate a zonal assignment no lower than Zone P 13 and Zone P 14 respectively, whereas taxa such as "*Globigerinoides*" *higginsi* and *Truncorotaloides rohri* in these samples indicate an assignment no higher than Zone P 14. The occurrence of *Globigerina angiporoides minima* and *G. praebuloides* in Mf1351 and Mf3305 followed by the occurrence of typical representatives of *G. angiporoides angiporoides* in Mf1352 and Mf3306 indicate assignment of these levels to Zone P 15 to Zone P 16.

Samples from the lower part of the Rices Mudstone Member of the San Lorenzo Formation are assigned to Oligocene zonal interval P 19 - P 20 because of the occurrence of *Globigerina ampliapertura* and associated coccoliths (see below).

After completion of Poore and Brabb's (1977) study, two additional samples containing diagnostic planktic foraminifers were discovered in the Twobar Shale Member (table 10). Planktic foraminifers in sample Mf4756, from the San Lorenzo River section (fig. 24), are referable to middle Eocene Zone P 14. Planktic foraminifers in sample Mf2657, from the upper part of the type Twobar Shale Member in the nearby Kings Creek section, are referable to upper Eocene Zone P 15 or Zone P 16. Thus these data corroborate the study of Poore and Brabb (1977) showing that the Zone P 14 - P 15 boundary (middle Eocene-upper Eocene boundary of Hardenbol and Berggren,

TABLE 8.—Distribution of coccoliths and zone assignments for the Zayante Creek section.

[Taxa shown with asterisk are considered to be reworked]

Taxon	<i>Coccolithus formosus</i> *	<i>C. pelagicus</i> (s. ampl.)	<i>Cyclicargolithus abisectus</i>	<i>C. floridanus</i>	<i>Dictyococcites bisectus</i>	<i>D. scrippsae</i>	<i>Discoaster deflandrei</i>	<i>D. divaricatus</i>	<i>D. lodoensis</i> *	<i>Discolithina plana</i> (s. ampl.)	<i>Discolithina</i> spp.	<i>Helicosphaera euphratis</i>	<i>H. intermedia</i>	<i>H. obliqua</i>	<i>H. perchnielsenae</i>	<i>H. recta</i>	<i>Ishmolithus recurvus</i> *	<i>Reticulofenestra</i> spp.	<i>Sphenolithus dissimilis</i>	<i>S. moriformis</i>	<i>S. radians</i> *	<i>Sphenolithus</i> spp.	<i>Tessellatolithus</i> sp.	<i>Tribrachiatus orthostylus</i> *	<i>Zygrhablithus</i> sp.	Coccolith zone of Bukry (1975)	Age
JC61-11d	X	aff.	X	aff.	X	aff.	X	aff.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	?	Miocene
76CB1592 Mf4677	X	X	X	X	X	X	X	X	X	X	aff.	aff.	aff.	aff.	X	X	X	X	X	X	X	X	X	X	X	<i>Sphenolithus ciperoensis</i> or <i>Sphenolithus distentus</i>	Late Oligocene
76CB1595 Mf4680	aff.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
76CB1601 Mf4681	X	X	aff.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
76CB1603 Mf4683	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
76CB1606 Mf4685	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	?	Late Oligocene?
76CB1607 Mf4686	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		

1978) is within the lower part of the Twobar Shale Member of the San Lorenzo Formation.

Coccoliths in samples from the San Lorenzo River section support the age interpretations derived from planktic foraminifers. The presence of *Reticulofenestra umbilica* in sample Mf3301 and *Dictyococcites bisectus* in sample Mf3304 indicates assignment of these samples respectively to the *Discoaster bifax* Subzone and the *Discoaster saipanensis* Subzone of the middle Eocene *Reticulofenestra umbilica* Zone. The assemblage from the next higher sample (Mf3305) is difficult to assign to a zone, but the occurrence of *Sphenolithus predistentus*, *Pedinocyclus larvalis*, and *Chiasmolithus altus* in Mf3306 suggests correlation of this sample with the upper Eocene *Discoaster barbadiensis* Zone. The sparse assemblage of sample Mf3308 contains *Discoaster saipanensis* and is therefore referable to the same zone. We did not recover age-diagnostic coccolith assemblages from the Rices Mudstone Member of the San Lorenzo Formation in our study; however, Bukry, Brabb, and Vedder (1977) reported an assemblage, containing *Cyclicargolithus abisectus*, from a sample (68CB123) equivalent to our sample Mf3310, that they questionably assign to the Oligocene *Sphenolithus distentus* Zone.

Figure 26 summarizes the relation of lithostratigraphy and biostratigraphic determinations derived from planktic foraminifers, coccoliths, and benthic

foraminifers from the San Lorenzo River section. Planktic foraminifer data indicate that the Zone P 14 - P 15 boundary occurs in the lower part of the Twobar Shale Member of the San Lorenzo Formation, that is, between the levels delineated by samples Mf3304 and Mf3305. Coccolith data show that the boundary between the *Discoaster saipanensis* Subzone of the *Reticulofenestra umbilica* Zone and the *Discoaster barbadiensis* Zone is at approximately the same place, that is, between samples Mf3304 and Mf3306. The two data sets are in good agreement and show that at this location the middle Eocene-upper Eocene boundary coincides with the base of the *Amphimorphina jenkinsi* Zone of the Narizian Stage. As noted above, we did not recover age-diagnostic coccoliths from the Rices Mudstone Member of the San Lorenzo Formation; however, Warren and Newell (1980) report *Discoaster barbadiensis* from the lower (Refugian) part of the Rices Mudstone Member at this section, which suggests an upper Eocene assignment. In addition, other published studies (Brabb and others, 1971; Lipps, 1967a) suggest an upper Eocene assignment for at least part of the Refugian Stage at other localities. Thus the Eocene-Oligocene boundary in the San Lorenzo River section is tentatively placed in the lower part of the Rices at the boundary between the Refugian and the Zemorrian stages. The remaining data from this section indicate an Oligocene




FIGURE 23.—Stratigraphic column for north limb of Zayante Creek section and location of samples yielding calcareous nannofossils. Modified from Brabb, Clark, and Throckmorton (1977, fig. 20). Benthic foraminiferal stage boundaries are from McDougall (this volume). Data on coccoliths and benthic foraminifers from the Lambert Shale are from different samples. No samples were examined from the south limb of the Zayante Creek section for this study.

Continued




FIGURE 23.—Continued.




FIGURE 24.—Stratigraphic column for San Lorenzo River section and location of samples yielding calcareous plankton. Modified from Poore and Brabb (1977, fig. 2). Sample 68CB123 of Bukry, Brabb, and Vedder (1977) is approximately equivalent to sample Mf3310.

[Taxa shown with asterisk are considered to be reworked]

[illegible]

Butano Sandstone (upper part)	San Lorenzo Formation		FORMATION
	Two-bar Shale Member	Rices Mudstone Member (lower part)	MEMBER
			SAMPLE
———— “<i>Globigerinoides</i>” <i>higgins</i> ———— <i>Globorotaloides suteri</i> ———— <i>Globorotalia cerroazulensis cerroazulensis</i> ———— ? <i>Truncorotaloides rohri</i> ———— <i>Globorotaloides carcoselleensis</i> —— — ? <i>Globigerina utilisindex</i> —— — <i>Globigerina angiporoides minima</i> —— — ? <i>Globigerina angiporoides angiporoides</i> ———— <i>Globigerina praebuloides</i> —— — <i>Globigerina ampliapertura</i> —— — <i>Globorotalia gemma</i>			PLANKTIC FORAMINIFERAL SPECIES
P13-14	P14	P15-16 P16 P19 P20	PLANKTONIC FORAMINIFERAL ZONE
———— <i>Reticulofenestra umbilica</i> ———— <i>Dictyococcites bisectus</i> ———— <i>Discoaster nodifer</i> ———— <i>Discoaster saipanensis</i> ———— <i>Daktylethra punctulata</i> —— — ? <i>Sphenolithus predistentus</i> —— — <i>Pedinocyclus larvalis</i> —— — <i>Chiasmolithus altus</i> —— — <i>Cyclicargolithus abisectus</i>			COCCOLITH SPECIES
<i>Discoaster barbadensis</i> zone ? <i>Discoaster saipanensis</i> subzone <i>Discoaster saipanensis</i> subzone <i>Discoaster bifax</i> zone <i>Discoaster barbadensis</i> zone ? <i>Sphenolithus distentus</i> zone			COCCOLITH ZONE OR SUBZONE

FIGURE 25.—Ranges of selected calcareous plankton and resultant zone assignments for the San Lorenzo River section.

assignment for at least the lower part of the *Uvigerina gallowayi* Zone of the Zemorrian Stage.

CHURCH CREEK AREA

Brabb, Bukry, and Pierce (1971) reported coccoliths indicative of the *Discoaster barbadiensis* Zone and benthic foraminifers referable to the Refugian Stage from the Church Creek Formation in the Church Creek area of the Santa Lucia Range (fig. 20). As there are few published records of planktic foram-

inifers in California marine strata assigned to the Refugian Stage, samples from this unit including those reported by Brabb, Bukry, and Pierce (1971) were examined for planktic foraminifers. Most samples are barren of planktic foraminifers or contain poorly preserved, nondiagnostic assemblages. Two samples, however, did yield relatively diverse assemblages (table 11). The assemblages present in samples Mf2616 and Mf2626 are compatible with the upper Eocene assignment derived from coccoliths of this

FORMATION	MEMBER	PLANKTONIC FORAMINIFERAL ZONE OF BLOW (1969)	COCCOLITH ZONE OR SUBZONE OF BUKRY (1975)	BENTHONIC FORAMINIFERAL		EPOCH
				STAGE	ZONE	
San Lorenzo Formation	Rices Mudstone Member (lower part)	XXXXX P19-P20	X ? <i>Sphenolithus distentus</i> zone	Zemorrian	<i>Uvigerina gallowayi</i> zone	Oligocene
				Refugian		
Butano Sandstone (upper part)	Two-bar Shale Member	X P16 X P15-16 X P14	X <i>Discoaster barbadiensis</i> zone X <i>Discoaster barbadiensis</i> X ? <i>Discoaster saipanensis</i> X <i>Discoaster saipanensis</i> subzone	Narizian	<i>Amphimorphina jenkinsi</i> zone	Late Eocene
		X P13-14	X <i>Discoaster bifax</i> zone			Middle

Probable paraconformity

FIGURE 26.—Summary of lithostratigraphy and biostatigraphic determinations for the San Lorenzo River section. Data on benthic foraminifers from McDougall (this volume). The *Valvulineria tumeyensis* Zone and *Uvigerina vicksburgensis* Zone of the Refugian Stage appear to be present, albeit poorly developed (McDougall, this volume). Modified from Poore and Brabb (1977, fig. 3). Sample levels for calcareous plankton are shown by X.

formation. Moreover, the occurrence of *Globorotalia gemma* is significant in that this form ranges no lower than Zone P 16 (Blow, 1969). Thus these assemblages are correlative with Eocene Zones P 16 or P 17.

DISCUSSION

Planktic calcareous microfossils were either absent or very sparse and poorly preserved in most samples from sections we examined for this field conference. Additionally, most zone assignments for planktic

TABLE 10.—Occurrence of planktic foraminifers and zone assignments from samples Mf4756 and Mf2657, from the Twobar Shale Member of the San Lorenzo Formation in the San Lorenzo River and Kings Creek sections

Taxon	Sample number	
	Mf 4756	Mf 2657
<i>Chiloguembelina</i> sp.	x	---
<i>Globigerina angiporoides angiporoides</i> Hornibrook.....	---	x
<i>G. angiporoides minima</i> Jenkins.....	x	x
<i>G. eocaena</i> Gumbel s.l.	x	---
<i>G. praebulloides</i> Blow	---	x
<i>G. praeturritilina</i> Blow and Banner	---	x
<i>G. pseudovenezuelana</i> Blow and Banner	---	x
<i>G. senilis</i> Bandy	x	x
<i>G. tripartita</i> Koch	---	x
<i>G. utilisindex</i> Jenkins and Orr	x	---
<i>Globigerinatheka index</i> (Finlay) s.l.	x	---
<i>G. mexicana</i> (Cushman) s.l.	x	---
<i>Globigerinita martini</i> Blow and Banner s.l.	x	x
" <i>Globigerinoides</i> " <i>higginsii</i> Bolli	x	---
<i>Globorotalia cerroazulensis</i> (Cole)	---	x
<i>Globorotaloides carcoselleensis</i> Toumarkine and Bolli	x	x
<i>G. wilsoni</i> (Cole)	x	x
<i>Gumbelitria columbiana</i> Howe	x	---
<i>Hantkenina</i> sp.	---	x
<i>Planorotalites pseudoscutula</i> Glaessner)	x	---
<i>Pseudohastigerina lillisi</i> (Church)	x	x
<i>P. micra</i> (Cole)	x	x
<i>Truncorotaloides collectea</i> (Finlay)	x	---
<i>T. rohri</i> Bronnimann and Bermudez	x	---
Zone assignment	P14	P15-P16

TABLE 11.—Planktic foraminifers and zone assignments for samples from the Church Creek Formation, Santa Lucia Range.

Taxon	Sample number	
	Mf2616	Mf2626
<i>Globigerina angiporoides angiporoides</i> Hornibrook	x	---
<i>G. a. minima</i> Jenkins	x	---
<i>G. officinalis</i> Subbotina	x	---
<i>G. ouachitaensis</i> Howe and Wallace	x	---
<i>G. praebulloides</i> Blow	x	---
<i>G. pseudovenezuelana</i> Blow and Banner	---	x
<i>G. tripartita</i> Koch	---	x
<i>Globigerinita martini</i> Blow and Banner (s.l.)	x	x
<i>Globorotalia gemma</i> Jenkins	x	x
<i>Globorotaloides suteri</i> Bolli	x	---
Zone assignment	P16 or P17	P16 or P17

foraminifer or coccolith assemblages were based on secondary rather than primary zone indicators. Even though we employed concentration techniques in coccolith preparations, primary zone indicators were still not encountered in most samples. The absence of calcareous plankton in these sections is due to a variety of factors including: unfavorable facies, geographic limits of index taxa, diagenesis, and the deep weathering characteristic of the Coast Ranges. Nonetheless, the record from the San Lorenzo River section is valuable information concerning upper middle Eocene to upper Eocene calcareous plankton from the Pacific coast. Undoubtedly these data will become important in future studies concerning biogeography of calcareous plankton of this region. Similarly, data from the San Lorenzo River section show that the middle Eocene-upper Eocene boundary of international usage is approximated by the base of the *Amphimorphina jenkinsi* Zone of the Narizian Stage and further that in this area, the Narizian Stage-Refugian Stage boundary is in the upper Eocene. Data from the Zayante Creek and Año Nuevo sections, albeit sparse, suggest that the Zemorrian Stage-Saucesian Stage boundary is uppermost Oligocene or lower Miocene on the basis of calcareous plankton.

TAXONOMIC NOTES

Comments in this section are restricted to planktic foraminifers illustrated on plates 7 and 8. A more complete discussion and illustrations of plankton foraminifers from the San Lorenzo River section are presented in Poore and Brabb (1977). Coccoliths identified during the present study are listed at the end of this report along with reference to a publication giving a representative light-microscope illustration.

Chiloguembelina cubensis (Palmer)
Plate 7, figures 1, 2

Gumbelina cubensis Palmer, 1934, p. 235-254, figs. 1-6.

Chiloguembelina cubensis occurs as a minor component in several of the planktic assemblages from the San Lorenzo River section. The lower stratigraphic limit of this taxon is ambiguous, but is probably no lower than Zone P 13. Several workers (Jenkins and Orr, 1972; Berggren and Amdurer, 1973) suggest that the last occurrence of *Chiloguembelina cubensis* can be used to divide Oligocene Zone P 21 into two informal subzones.

Globigerina praeturritilina Blow and Banner
Plate 8, figures 7, 8

Globigerina turritilina praeturritilina Blow and

Banner, 1962, p. 99, pl. XIII, figs. a-c.

The relatively high spire, open umbilicus, and arching aperture of this taxon serve to distinguish it from large generalized middle and late Eocene *Globorotalia*.

Globorotalia sp. aff. *G. munda* Jenkins
Plate 8, figures 5, 6

Globorotalia munda Jenkins, 1966, p. 1121, pl. 13, no. 152-166, text fig. 14, no. 126-133.

This form is sparse in samples Mf4664 and Mf4665 from the Año Nuevo section. Poor preservation and few individuals available for study preclude a more positive identification.

Globorotalia nana Bolli
Plate 8, figures 1-4

Globorotalia opima nana Bolli, 1957, p. 118, pl. 28, fig. 3.

As presently understood, typical *Globorotalia nana* ranges no higher than upper Oligocene Zone P 22 (Blow, 1969; Stainforth and others, 1975). Other similar forms, such as *Globorotalia pseudocontinua* Jenkins, however, range well into the Miocene. Thus care must be taken in making a pre-Miocene assignment based solely on the occurrence of *Globorotalia nana*.

Gümbelitra columbiana Howe
Plate 7, figures 7-9

Gümbelitra columbiana Howe, 1939, p. 62, pl. 8, figs. 12 and 13.

Poorly preserved specimens of this taxon are present in sample Mf4756 from the San Lorenzo River section. *Gümbelitra columbiana* was not detected in the other middle and upper Eocene samples from this section (Poore and Brabb, 1977).

Pseudohastigerina lillisi (Church)
Plate 7, figures 3, 6

Pullenia lillisi Church, 1931, p. 209, pl. A, fig. 10

The concepts of McKeel and Lipps (1975) are followed in identifying this species. Thus the tendency towards loose coiling in mature specimens and the relatively high arched aperture distinguish *Pseudohastigerina lillisi* from *P. micra*.

Pseudohastigerina micra (Cole)
Plate 7, figures 4, 5

Nonion micrus Cole, 1927, p. 22, pl. 5, fig. 12

Pseudohastigerina micra ranges from within basal middle Eocene Zone P 10 through Oligocene zonal interval P 19 - 20. This taxon is fairly common in the Pacific coast rocks of appropriate age and is often

useful as a secondary indicator for detecting the lower Eocene-middle Eocene boundary.

CALCAREOUS NANNOFOSSILS

Some representative coccolith species that are well known from the Gulf Coast, Europe, and Deep Sea Drilling Project cores are illustrated (pl. 9) to show the typical moderate preservation state of northern California assemblages.

LOCALITY DATA

Samples Mf2616 and Mf2626 from the Santa Lucia Range.

Both samples are from the Church Creek Formation in the Chews Ridge 7.5-minute quadrangle, Monterey County, California. Samples were collected by E. E. Brabb in 1969.

Sample Mf2616, field number 69CB651. Lat 36° 16'4" N., long 121° 34'8" W.

Sample Mf2626, field number 69CB642. Lat 36° 16'9" N., long 121° 35'6" W.

Sample taken at creek junction, elevation 2,860 ft (872 m).

ILLUSTRATION REFERENCES FOR IDENTIFIED COCCOLITH SPECIES

- Braarudosphaera bigelowi* (Gran and Braarud)—Bybell, 1975.
Bramletteius serraculoides Gartner—Gartner, 1969.
Campylosphaera dela (Bramlette and Sullivan)—Bramlette and Sullivan, 1961.
C. eodela Bukry and Percival—Bukry and Percival, 1971.
Chiasmolithus altus Bukry and Percival—Bukry and Percival, 1971.
C. expansus (Bramlette and Sullivan)—Bramlette and Sullivan, 1961.
C. grandis (Bramlette and Riedel)—Bramlette and Sullivan, 1961.
C. solitus (Bramlette and Sullivan, 1961).
Coccolithus cribellum (Bramlette and Sullivan)—Bramlette and Sullivan, 1961.
C. formosus Kamptner—Kamptner, 1963.
C. staurion Bramlette and Sullivan—Bramlette and Sullivan, 1961.
Cruciplacolithus tenuis (Stradner) s. ampl.—Martini, 1971.
Cyclicargolithus abisectus (Muller)—Bukry and Percival, 1971.
C. floridanus (Roth and Hay)—Bramlette and Wilcoxon, 1967 (as *C. neogammation*).
C.? luminus (Sullivan) n. comb.—basonym: *Cyclococcolithus luminus* Sullivan, 1965, p. 33, pl. 3, figs. 9a, b.
Cyclococcolithina gammation (Bramlette and Sullivan)—Bukry, 1974.
C.? kingi (Roth)—Gartner, 1971 (as *C. protoannula*).
Daktylethra punctulata Gartner—Gartner and Bukry, 1969.
Dictyococcites bisectus (Hay, Mohler, and Wade)—Bukry and Percival, 1971.
D. scrippsae Bukry and Percival—Bukry and Percival, 1971.
Discoaster barbadiensis Tan—Bramlette and Sullivan, 1961.
D. deflandrei Bramlette and Riedel—Bramlette and Wilcoxon, 1967.

- Discoaster divaricatus* Hay—Hay and others, 1967.
D. elegans Bramlette and Sullivan—Bramlette and Sullivan, 1961.
D. exilis Martini and Bramlette—Martini and Bramlette, 1963.
D. lodoensis Bramlette and Riedel—Bramlette and Sullivan, 1961.
D. nodifer (Bramlette and Riedel)—Bukry, 1974.
D. saipanensis Bramlette and Riedel—Bukry, 1974.
D. tani Bramlette and Riedel—Bramlette and Riedel, 1954.
D. variabilis Martini and Bramlette, 1963.
Discolithina distincta (Bramlette and Sullivan)—Bramlette and Sullivan, 1961.
D. multipora (Kamptner)—Kamptner, 1948.
D. plana (Bramlette and Sullivan)—Bramlette and Sullivan, 1961.
Helicosphaera bramlettei (Muller)—Muller, 1970.
H. carteri (Wallich)—Bramlette and Wilcoxon, 1967.
H. compacta Bramlette and Wilcoxon—Bramlette and Wilcoxon, 1967.
H. euphratis Haq—Haq, 1973.
H. intermedia Martini—Haq, 1973.
H. obliqua Bramlette and Wilcoxon—Bramlette and Wilcoxon, 1967.
H. perch nielsenasae Haq—Haq, 1973.
H. recta Haq—Haq, 1973.
Isthmolithus recurvus Deflandre—Hay and others, 1966.
Lanternithus minutus Stradner—Gartner and Burky, 1969.
Lophodolichus reniformis Bramlette and Sullivan—Bramlette and Sullivan, 1961.
Micrantholithus flos Deflandre—Sullivan, 1964.
Pedinocyclus larvalis (Bukry and Bramlette)—Bukry and Bramlette, 1971.
Pemma angulatum Martini—Martini, 1959.
P. basquensis (Martini) s. ampl.—Bybell, 1975.
Reticulofenestra oamaruensis (Deflandre)—Stradner and Edwards, 1968.
R. reticulata (Gartner and Smith)—Bybell, 1975.
R. umbilica (Levin)—Bybell, 1975.
Rhabdosphaera tenuis Bramlette and Sullivan—Bramlette and Sullivan, 1961.
Sphenolithus belemnus Bramlette and Wilcoxon—Bramlette and Wilcoxon, 1967.
S. conicus Bukry—Bukry, 1971.
S. dissimilis Bukry and Percival—Bukry and Percival, 1971.
S. furcatolithoides Locker—Locker, 1967.
S. heteromorphus Deflandre—Bramlette and Wilcoxon, 1967.
S. moriformis (Bronnimann and Stradner)—Bramlette and Wilcoxon, 1967.
S. predistentus Bramlette and Wilcoxon—Bramlette and Wilcoxon, 1967.
S. pseudoradians Bramlette and Wilcoxon—Bramlette and Wilcoxon, 1967.
S. radians Deflandre—Sullivan, 1965.
S. spiniger Bukry—Bukry, 1971.
Transversopontis pulcheroides (Sullivan)—Sullivan, 1964 (as *Discolithus pulcheroides*).
Tibrachiatius orthostylus Shamrai—Bramlette and Sullivan, 1961 (as *Discoaster tibrachiatius*).
Triquetrorhabdulus carinatus Martini—Bramlette and Wilcoxon, 1967.
Zygolithus dubius Deflandre—Bybell, 1975.
Zygrhablithus bijugatus (Deflandre)—Bybell, 1975.
- Indeterminate or undifferentiated species of the following genera are also tabulated:
- Chiasmolithus* Hay, Mohler, and Wade, 1966.
Coccolithus Schwarz, 1894.
Discolithina Loeblich and Tappan, 1963.
Helicosphaera Kamptner, 1954.
Reticulofenestra Hay, Mohler, and Wade, 1966.
Rhabdosphaera Haeckel, 1894.
Sphenolithus Deflandre, 1952.
Tessellatolithus Haq, 1968.
Thoracosphaera Kamptner, 1927.
Trochoaster Klumpp, 1953.
Zygrhablithus Deflandre, 1959.

UPPER EOCENE TO LOWER MIOCENE BENTHIC FORAMINIFERS FROM THE SANTA CRUZ MOUNTAINS AREA, CALIFORNIA

By KRISTIN MCDougALL

ABSTRACT

Benthic foraminiferal assemblages from three sections (San Lorenzo River, Zayante Creek, and Año Nuevo sections) in the Santa Cruz Mountains area, California, provide a sequence of species that are diagnostic of the middle Tertiary California benthic foraminiferal stages—Narizian through Relizian Stages. The lower Narizian *Bulimina corrugata* Zone is recognized in the San Lorenzo River section by the presence of restricted early Narizian species such as *Amphimorphina becki*, *Cibicides pachyderma*, *Karrerella elongata*, and *Pleurostomellina alternans*. The upper Narizian *Amphimorphina jenkinsi* Zone is recognized in this section by the presence of *Uvigerina garzaensis nudorobusta*,

eggerella elongata, and the first occurrence of many species that range into younger strata. The Refugian Stage is recognized in the strata above a glauconite bed in the Twobar Shale Member of the San Lorenzo Formation. These Refugian faunas contain species typical of deeper water and have a more northern aspect than present in the type area of the Refugian Stage, and prove additional data on the age ranges of the diagnostic Refugian species.

Parts of the Zemorrian Stage are recognized from all three sections examined. The lower Zemorrian *Uvigerina gallowayi* Zone is identified in the San Lorenzo River and Zayante Creek sections by diagnostic species such as *Uvigerina gallowayi*, *Textularia shivelyi*, *Pseudonodosaria gallowayi*, and *Uvigerinella obesa impolita*. The upper Zemorrian *Uvigerinella sparsicostata*

Zone is recognized in the Zayante Creek and Año Nuevo sections by the lowest occurrences of *Siphogenerina nodifera*, *S. mayi*, *S. multicostata*, *Bulimina carnerensis*, and *Uvigerinella sparsicostata*. Saucian faunas identified from the Año Nuevo section include *Baggina robusta*, *Bolivina marginata adelaidana*, and *Dentalina quadrulata*. This section also contains Relizian assemblages, which are identified by the presence of poorly preserved specimens of *Siphogenerina hughesi* and *Valvulineria californica*. The lower Zemorrian-upper Zemorrian and Zemorrian-Saucian boundaries are gradational owing to continuous deposition and unchanged bathymetric conditions; this necessitates some modification of species ranges and diagnostic species defining the stage and zone boundaries.

The infrequent and sporadic occurrences of planktic foraminifers and nannofossils in these sections do not allow definitive calibration of planktic foraminiferal zone boundaries to benthic foraminiferal stage or zone boundaries but do suggest the following correlations: (1) the lower Narizian-upper Narizian boundary corresponds approximately to the middle Eocene-upper Eocene boundary (2) the Refugian Stage is upper Eocene, (3) the Zemorrian Stage is Oligocene, and (4) the Saucian Stage is lower Miocene.

INTRODUCTION

Benthic foraminifers were examined from three sections in the central California Coast Ranges as part of the field conference and meeting held by the International Commission on Paleogene Stratigraphy in 1977. The purpose of this meeting was to correlate the West Coast zonal schemes based on benthic foraminifers to zonal schemes of wider usage

based on nannofossils and planktic foraminifers, and ultimately to correlate the California sequences with the European stages. The purpose of this paper, therefore, is to provide the stage, zonal, and subzonal assignments based on benthic foraminifers.

The impetus for microfossil studies and the development of a zonal scheme was the discovery of oil in California in the early part of the twentieth century. This activity led to the development of a Tertiary zonal scheme based on benthic foraminifers. The definitions and descriptions of these stages and zones are given in Kleinpell (1938), Mallory (1959), and Donnelly (1976—this modified the original definition of Schenck and Kleinpell, 1936). Work since then has been directed toward describing faunas and extending these zonal schemes along the west coast of North America. Extension of these stages outside their type localities has resulted in correlation problems. Planktic foraminiferal nannofossil studies (Steineck and Gibson, 1971; Bukry and others, 1977; Poore, 1976, and this volume) have demonstrated that the benthic foraminiferal stages are commonly time-transgressive from one area to another when compared to zonal schemes based on planktic organisms. Benthic foraminiferal studies should now be directed toward recognizing the problems and causes, and resolving them.

Benthic foraminifers from the Santa Cruz Mountains (San Lorenzo River and Zayante Creek sections, fig. 27) and along the adjacent coast (Año Nuevo section, fig. 27) were analyzed to determine stage and zone boundaries. These sections provide a nearly continuous sequence from the Narizian Stage to the Relizian Stage (fig. 28). The benthic foraminiferal assemblages include species from bathymetric facies different than in the type areas of the stages and zones and thus provide additional information on species age ranges and environmental tolerances. This information suggests that minor revisions are needed in the benthic foraminiferal criteria used to recognize the stages and zones. Planktic microfossil data, though sparse or limited, do suggest that these stages can be correlated with international time scales. The sections examined in this report do not overlap enough in time, however, to consider the problem of time-transgressive stages. As more complex sampling and more detailed work with planktic and benthic organisms is done, factors contributing to or causing the time-transgressive stages can be identified and perhaps resolved.

Acknowledgments—I thank W. V. Sliter, R. Z. Poore, and E. E. Brabb for discussions and suggestions on the manuscript and also Robert Oscarson




FIGURE 27.—General location of sections examined for this study.

and M. G. Murphy for assisting in the preparation of the plates. Special thanks are given to E. E. Brabb for measuring sections and collecting the samples used in this study.

SAN LORENZO RIVER SECTION

The San Lorenzo River section is in the Santa Cruz Mountains northwest of the town of Santa Cruz (fig. 27). Tertiary rock units exposed along this river

section include the Butano Sandstone, San Lorenzo Formation, and Vaqueros Sandstone. New benthic foraminiferal data (table 12) from the San Lorenzo Formation augment and modify (fig. 29) the earlier work by Sullivan (1962). Detailed descriptions of the lithology, structure, and sample localities are given in Sullivan (1962), Poore and Brabb (1977), and Brabb, Clark, and Throckmorton (1977). See table 13 for field numbers and the corresponding sample numbers used in this report.

SERIES	BENTHIC FORAMINIFERAL STAGE	BENTHIC FORAMINIFERAL ZONE	SUBZONE	SAN LORENZO RIVER	ZAYANTE CREEK	AÑO NUEVO
Miocene	Relizian		No subzones			Mf4668 Mf4662
		<i>Uvigerinella obesa impolita</i>				
	Saucesian	<i>Plectofrondicularia miocenica</i>				
		<i>Siphogenerina transversa</i>				
Oligocene	Zemorrrian	<i>Uvigerinella sparsicostata</i>			Mf4671 Mf4677 Mf4679 Mf4681	Mf4664 Mf1374-1376 Mf4665
		<i>Uvigerina gallowayi</i>			Mf4682-4683	
				Mf2700-2701 Mf1366-1363		
Eocene	Refugian	<i>Uvigerina vicksburgensis</i>		Mf1364-1365		
		<i>Valvulineria tumeyensis</i>	<i>Uvigerina atwilli</i>	Mf1360-1362		
			<i>Cibicides haydoni</i>	? Missing ?		
	Narizian	<i>Amphimorphina jenkinsi</i>	No subzones	Mf1355-1357 Mf1350-1352		
		<i>Bulimina corrugata</i>	<i>Uvigerina garzaensis</i>	Mf3304		
			<i>Uvigerina churchi</i>			

FIGURE 28.—Stratigraphic relations and stage and zone assignments of samples examined for this study. Benthic foraminiferal stages and zones were defined by Schenck and Kleinpell (1936), Kleinpell (1939), Mallory (1959), and Donnelly (1976).

Benthonic foraminiferal stage	Narizian							Refugian					Early Zemorrian				
	Early		Late														
Taxon	Mf1304	Mf1350	Mf1351	Mf1352	Mf1355	Mf1356	Mf1357	Mf1360	Mf1361	Mf1362	Mf1364	Mf1365	Mf1366	Mf2700	Mf2701	Mf1367	Mf1368
<i>Amphimorphina becki</i> Mallory	R	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Anomalina garzaensis</i> Cushman and Siegfus	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Asterigerina crassaformis</i> Cushman and Siegfus	R	—	R	—	—	—	—	—	—	—	—	—	?	—	—	—	—
<i>Bathysiphon eocenica</i> Cushman and Hanna	F	F	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Bathysiphon</i> spp.	F	—	—	—	—	—	—	—	—	—	—	—	F	—	—	—	—
<i>Bifarina eleganta</i> (Plummer)	F	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Boldia</i> cf. <i>B. hodgei</i> (Cushman and Schenck)	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Bolivina</i> sp.	R	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Bulimina corrugata</i> Cushman and Siegfus	F	—	—	—	—	F	C	—	—	—	—	—	—	—	—	—	—
<i>Bulimina curtissima</i> Cushman and Siegfus	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Chilostomella</i> sp.	R	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Cibicides pachyderma</i> (Rzehak)	R	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Cibicides</i> spp.	F	—	F	—	—	R	—	—	—	—	—	—	—	—	—	—	—
<i>Cibicides spiropunctatus</i> Galloway and Morrey	A	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Dentalina spinosa ornator</i> Sullivan	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Dorothia cubana</i> (Cushman and Bermudez)	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Eggerella elongata</i> Blaisdell	F	F	?	—	F	C	—	—	—	—	—	—	—	—	—	—	—
<i>Eggerella subconica</i> Parr	F	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
? <i>Eggerella subconica</i> Parr	F	—	F	—	—	F	—	—	—	—	—	F	—	—	—	—	—
<i>Ellipsonodosaria atlantisae hispidula</i> Cushman	—	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Gyroidina orbicularis planata</i> Cushman	—	—	—	—	—	—	—	R	—	—	—	—	—	A	—	R	F
<i>Haplophragmoides</i> spp.	C	—	—	—	F	F	—	F	—	—	F	—	—	—	—	—	—
<i>Karrerella elongata</i> Mallory	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Lenticulina midwayensis</i> (Plummer)	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Lenticulina pseudovortex</i> (Cole)	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Lenticulina</i> spp.	F	R	—	—	—	F	—	C	C	F	C	F	F	—	—	—	—
<i>Marginulina subbullata</i> Hantken	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Marginulina subbullata</i> of Mallory	R	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Martinottiella eocenica</i> Cushman and Bermudez	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Nodosaria</i> cf. <i>N. gyrata</i> Mallory	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Nodosaria longiscata</i> d'Orbigny	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Nodosaria pyrula</i> d'Orbigny	F	R	—	—	—	F	—	—	—	—	—	—	—	—	—	—	—
<i>Nodosaria</i> spp.	F	—	—	—	R	F	F	F	F	—	—	—	—	—	—	—	—
<i>Oridorsalis umbonatus</i> (Reuss)	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Plectina garzaensis</i> Cushman and Siegfus	C	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Pleurostomella alternans</i> Schwager	R	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Praeglobobulimina pupoides</i> (d'Orbigny)	F	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Pseudonodosaria inflata</i> (Bornemann)	R	—	—	—	—	—	—	F	F	—	—	—	F	F	—	—	—
<i>Pullenia quinqueloba</i> (Reuss)	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Quinqueloculina minuta</i> Beck	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Rhabdammina eocenica</i> Cushman and Hanna	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Reophax pilulifer</i> H. B. Brady	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Spiroplectammina richardi</i> Martin	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Uvigerina churchi</i> Cushman and Siegfus	C	R	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Vaginulinopsis saundersi</i> (Hanna and Hanna)	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Anomalina</i> cf. <i>A. umbonata</i> Cushman	—	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Boldia hodgei</i> (Cushman and Schenck)	—	R	—	—	F	—	F	C	C	F	—	—	—	—	—	—	—
<i>Bolivina gracilis</i> Cushman and Applin	—	R	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Bolivina kleinPELLI</i> Beck	—	R	F	—	—	—	—	F	F	—	—	—	—	F	—	—	—
<i>Bolivina scabrata</i> Cushman and Bermudez	—	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Bulimina microcostata</i> Cushman and Parker	—	F	C	—	—	—	—	—	—	—	—	—	—	—	—	—	—
? <i>Buliminella</i> sp.	—	R	—	—	—	—	—	—	—	—	—	—	—	—	—	—	




FIGURE 29.—Age interpretations of San Lorenzo River Section as determined by Sullivan (1962) and this study. Mf localities are from U.S. Geological Survey microfossil locality register, Menlo Park, Calif., and B localities are from University of California, Berkeley, fossil locality register. Stratigraphic column from Poore and Brabb (1977, fig. 2).

[Field numbers with the prefix 76CB and EB were collected by E. E. Brabb. Numbers preceded by JC were collected by J. C. Clark]

Sample number	Field number	Section	Sample number	Field number	Section
Mf1350	75CB1352	San Lorenzo River	Mf2192	JC-68-3	Año Nuevo
Mf1351	76CB1353	Do	Mf2193	JC-68-2	Do
Mf1352	76CB1354	Do	Mf2194	JC-68-4	Do
Mf1355	EB256G	Do	Mf2195	JC-68-1	Do
Mf1356	EB256B	Do	Mf2700	EB395D	San Lorenzo River
Mf1356	EB256K	Do	Mf2701	EB675	Do
Mf1356	EB256A	Do	Mf3294	76CB1451	Do
Mf1361	EB236D	Do	Mf4631	JC-67-5	Año Nuevo
Mf1362	EB256L	Do	Mf4661	76CB1541	Do
Mf1364	EB256N	Do	Mf4664	76CB1531	Do
Mf1365	EB256P	Do	Mf4665	76CB1532	Do
Mf1367	EB676	Do	Mf4667	76CB1521A	Do
Mf1368	EB677	Do	Mf4668	76CB1541B	Do
Mf1374	JC-69-1	Año Nuevo	Mf4671	76CB1562	Zayante Creek
Mf1375	JC-69-2	Do	Mf4677	76CB1592	Do
Mf2188	JC-69-4	Do	Mf4679	76CB1593	Do
Mf2189	JC-69-6	Do	Mf4681	76CB1601	Do
Mf2190	JC-68-4	Do	Mf4682	76CB1602	Do
Mf2191	MC-68-4	Do	Mf4683	76CB1603	Do

The oldest of the samples examined, Mf3304 (fig. 29), is from the lower part of the Twobar Shale Member of the San Lorenzo Formation. This sample was taken from just below sample B2233 (University of California, Berkeley number). Sullivan (1962) considered all the samples in the lower part of the San Lorenzo Formation to be of late Narizian age and diagnostic of the *Amphimorphina jenkinsi* Zone (fig. 29). Sample Mf3304 contains a more diverse and abundant microfauna than that recovered by Sullivan (1962) and suggests instead an early Narizian age. This sample includes species such as *Dorothia cubana* and *Lenticulina midwayensis* (fig. 30), which are usually considered to be diagnostic of the Ulati-




FIGURE 30.—Stratigraphic distribution of selected benthic foraminiferal species diagnostic of Narizian through lower Zemorrian Stages in San Lorenzo River section.

ian Stage. Species restricted to the lower Narizian, or which last occur in the lower Narizian *Bulimina corrugata* Zone include *Amphimorphina becki*, *Anomalina garzaensis*, *Bulimina curtissima*, *Cibicides pachyderma*, *C. spiropunctata*, *Ellipsonodosaria atlantisae hispidula*, *Karreriella elongata*, *Lenticulina pseudo-vortex*, and *Pleurostomella alternans* (fig. 30). The sample also contains many longer ranging species typical of the Narizian and the early Tertiary. There are no species definitive of or restricted to the late Narizian. The upper Narizian Stage assignment by Sullivan (1962) was based on the presence of several specimens he identified as *Amphimorphina jenkinsi* from sample B2231, which is from just above the contact between the Butano Sandstone and the San Lorenzo Formation. The specimen of *Amphimorphina jenkinsi* illustrated by Sullivan (1962, pl. 14, fig. 15) indicates that his identification is tenuous at best. Several specimens from near the top of the Butano Sandstone are identified as *Amphimorphina becki* by me from unpublished benthic foraminiferal data from sample Mf3301. These specimens represent a transitional form between *A. becki* and *A. jenkinsi*, with the peripheral ridges gone and only the basal costae remaining. This may be the form Sullivan (1962) and Schenck (1936) identify as *A. jenkinsi*; however, it is not *Amphimorphina jenkinsi*.

Less than one-third of the species from Mf3304 are also in the overlying sample Mf1350 (fig. 30). All of the species common to both samples are long-ranging species, except *Bifarina elegans* and *Uvigerina churchi*, which are rare in the upper Narizian. In this section both last occur in Mf1350 (fig. 30). The new benthic foraminiferal data indicate that the lower Narizian *Bulimina corrugata* Zone should be extended into the lower part of the Twobar Shale Member of San Lorenzo Formation (fig. 29). The data are insufficient to identify the subzone represented by this assemblage. The lower Narizian-upper Narizian boundary is between samples Mf3304 and Mf1350. Thus, in the San Lorenzo River section, Mf3304 is the stratigraphically highest sample in the lower Narizian *Bulimina corrugata* Zone and Mf1350 is the stratigraphically lowest sample in the upper Narizian *Amphimorphina jenkinsi* Zone.

The next group of samples, Mf1350 through Mf1357 (fig. 30), represents the upper Narizian *Amphimorphina jenkinsi* Zone. Present in these samples are numerous species that are restricted to or last occur in the upper Narizian such as *Bolivina scabrata*, *Bulimina corrugata*, *Bulimina microcostata*, *Eggerella elongata*, *Lenticulina welchi*, *Planularia markleyana*, *Stichocassidulina* cf. *S. thalmani*, and *Uvigerina garzaensis*. Also present are species which

first appear in the upper Narizian, such as *Boldia hodgei*, *Bulimina sculptilis lacinata*, *Plectofrondicularia packardi*, and *Valvulineria tumeyensis*. The late Narizian age for this interval is in agreement with Sullivan (1962).

Overlying Mf1357 is a glauconite bed that marks the base of the Rices Mudstone Member of the San Lorenzo Formation. The contact between the Twobar Shale and Rices Mudstone Members is believed to represent a paraconformity (Poore and Brabb, 1977). Benthic foraminiferal assemblages from the samples, Mf1360-Mf1362, above the glauconite are diagnostic of the lower Refugian. There is an abrupt change in the fauna, and species that continue from the older samples are long-ranging or those whose most common occurrence is in the Refugian, such as *Boldia hodgei* and *Plectofrondicularia packardi*. Species diagnostic of the Refugian Stage that first appear in these samples include *Anomalina californiensis*, *Cibicides elmaensis*, *Karreriella washingtonensis*, *Saracenaria schencki*, *Sigmomorphina schencki*, *Plectofrondicularia packardi multilineata*, and the *Uvigerina cocoaensis* species group (fig. 30). This association of species suggests an early Refugian age (fig. 29); however, the lower Refugian *Cibicides haydoni* Subzone of Donnelly (1976) or the "basal" Refugian faunas of McDougall (1975) appear to be partly or completely missing.

The benthic faunas from the Refugian type section are from only a limited bathymetric facies and are therefore not expected to be characteristic of all Refugian faunas. Schenck and Kleinpell (1936) in the original definition recognized the possibility of problems with paleoecologic facies and designated the fauna of the Lincoln Creek Formation (Lincoln Formation of Weaver, 1912) of Washington as representing a deeper water facies of the Refugian Stage. A similar fauna occurs in the San Lorenzo River section. In the deeper water facies of the Lincoln Creek Formation, *Cibicides elmaensis* and *Uvigerina cocoaensis* occur in the lower Refugian (Rau, 1958, 1966; Donnelly, 1976). The first occurrence of *Uvigerina cocoaensis* in the lower Refugian coincides with the worldwide first occurrence of the species (Lamb, 1964; Boersma, 1974). This first occurrence of *Uvigerina cocoaensis* can only be observed along the west coast of North America where the appropriate bathymetric facies is present. The first occurrence of *Uvigerina cocoaensis* in West Coast sections often records the first occurrence of appropriate bathymetric conditions and not the evolutionary first occurrence of the species. Also diagnostic of the lower Refugian in Washington and Oregon (Rau, 1958, 1966) are the species *Karreriella washingtonensis*, *Sigmomorphina schencki*, *Saracen-*

aria schencki, and *Cibicides haydoni*. The latter species also typifies the lower Refugian *Valvulinera tumeyensis* Zone of California (Donnelly, 1976).

The Refugian faunas also occur in samples Mf1364-Mf1365. Assignment to the upper Refugian *Uvigerina vicksburgensis* Zone is suggested but is not well documented. Diagnostic early Refugian species are not recognized in these samples, but many specimens cannot be identified. Sullivan (1962) reported Refugian faunas throughout samples B2252 to B2254. In B2253 to B2256, specimens that were identified by Sullivan (1962) as *Uvigerina gallowayi* are closer to *Uvigerina jacksonensis* and should be considered part of the *Uvigerina cocoaensis* species group, not *Uvigerina gallowayi*. This is the only new species in these samples, and it is not a good indicator of the upper Refugian Stage.

Placement of the lower and upper boundaries of the Refugian Stage and the species identifications of this report differ from those proposed by Sullivan (1962). Although sample B2251 was placed in the Refugian, it contains only species diagnostic of the Narizian *Amphimorphina jenkinsi* Zone, such as *Eggerella elongata*, *Bulimina microcostata*, *Planularia tolmani*, *Lenticulina welchi* (*Robulus chiranus* of Sullivan, 1962), and *Uvigerina garzaensis nudorobusta*. This sample should be considered Narizian. Only the last species is known to extend into the lower Refugian, and its presence in B2252 is questionable. The bathymetric conditions of B2251 and B2252 are different, and as the glauconite appears to represent a form of unconformity, the presence of the rare *Uvigerina garzaensis nudorobusta* is probably due to reworking. The placement of the boundary at the base of the glauconite, that is, above the highest sample in the Twobar Shale Member and below the lowest sample in the Rices Mudstone Member, coincides with the unconformity, the abrupt faunal change, and the occurrence of diagnostic Refugian species. The abrupt faunal change was also noted by Sullivan (1962) between samples B2251 and B2252.

The Refugian-Zemorrian boundary is placed above sample Mf1365 and below B2255 of Sullivan (1962). Sample Mf1365, which is below B2255, contains a faunal assemblage diagnostic of the Refugian Stage. There are few species in sample B2255 and in the overlying sample, B2256. *Plectofrondicularia packardii* and *P. p. multilineata* are present, but the upper limit of these species is not clearly defined. *Uvigerina jacksonensis* (*Uvigerina gallowayi* of Sullivan, 1962) ranges from late Eocene through the early Oligocene (Lamb, 1964; Boersma, 1974). Also, *Bolivina marginata* occurs in both these samples, and the first occurrence of this species is in the early Zemorrian (Klein-

pell, 1938). The placement of the Refugian-Zemorrian boundary (fig. 28) between Mf1365 and B2255 is compatible with the data.

The Zemorrian Stage is very poorly developed in the samples examined for this study. There is a gradual faunal change in samples Mf1366 through Mf1368, the result of the improved preservation of the faunas and the development of the Zemorrian fauna.

TABLE 14.—Benthic foraminifers from the Zayante Creek section of Brabb, Clark, and Throckmorton (1977, fig. 22)

Benthic foraminiferal stages	Zemor- rian		Sauce- sian(?)			
	Early	Late				
	Sample					
Taxon	Mf4683	Mf4682	Mf4681	Mf4679	Mf4677	Mf4671
<i>Bolivina marginata</i> Cushman	C	F	A	—	—	—
<i>Cibicides pseudoungerianus evolutus</i>	F	C	C	—	—	—
Cushman and Hobson	F	C	A	—	—	—
<i>Fursenkoina</i> cf. <i>F. bramletti</i> (Galloway and Morrey)	F	C	A	—	F	F
<i>Gyroidina orbicularis planata</i> Cushman	F	C	F	—	—	—
<i>Haplophragmoides</i> sp.	F	C	—	—	—	—
<i>Lenticulina simplex</i> (d'Orbigny)	—	—	—	C	C	?
<i>Lenticulina</i> sp.	A	—	—	—	—	F
<i>Nodosaria holserica</i> Schwager	F	F	—	—	—	—
<i>Oridorsalis umbonatus</i> (Reuss)	F	F	—	—	—	—
<i>Plectofrondicularia miocenica</i> Cushman	F	—	—	—	—	—
<i>Plectofrondicularia packardii multilineata</i>	—	—	—	—	—	—
Cushman and Simonson	F	—	—	—	—	—
<i>Plectofrondicularia vaughani</i> Cushman	F	—	F	—	—	—
<i>Pullenia salisburyi</i> Stewart and Stewart	F	F	—	—	—	—
<i>Textularia shivelyi</i> Kleinpell	F	F	—	—	—	—
<i>Textularia</i> sp. of Fairchild and others	F	—	—	—	—	—
<i>Uvigerinella sparsicostata</i> Cushman and Laiming	A	—	—	C	—	—
<i>Verneulina</i> sp. of Fairchild and others	A	C	F	—	—	—
<i>Ammodiscus incertus</i> d'Orbigny	—	—	R	—	—	—
<i>Anomalina californiensis</i> Cushman and Hobson	—	C	F	F	—	—
<i>Bathysiphon eocenica</i> Cushman and Hanna	—	A	R	—	—	—
<i>Buliminella curta</i> Cushman	—	C	A	—	—	—
<i>Cassidulina crassipunctata</i> Cushman and Hobson	—	A	—	—	—	—
<i>Cyclammmina incisa</i> (Stache)	—	A	—	C	A	—
<i>Cyclammmina</i> sp.	—	A	F	—	—	—
<i>?Dentalina quadrulata</i> Cushman and Laiming	—	F	F	—	—	—
<i>Dentalina</i> sp.	—	F	F	—	—	—
<i>Fissurina</i> sp.	—	F	F	R	—	—
<i>Frondicularia tenuissima</i> Hantken	—	F	F	—	—	—
<i>Gaudryina gracilis</i> Cushman and Laiming	—	F	F	—	—	—
<i>Gaudryina triangularis</i> Cushman	—	—	R	—	—	—
<i>Globobulimina pacifica</i> Cushman	—	F	C	R	F	—
<i>?Karrerella</i> sp.	—	F	F	C	—	—
<i>Lagena sulcata</i> (Walker and Jacob)	—	F	F	—	—	—
<i>Lenticulina calcar</i> (Linne)	—	F	F	—	—	—
<i>Lenticulina</i> cf. <i>L. clypeiformis</i> (d'Orbigny)	—	F	—	—	—	—
<i>Lenticulina inornata</i> (d'Orbigny)	—	C	C	A	—	F
<i>Lenticulina mayi</i> (Cushman and Parker)	—	F	—	—	—	—
<i>Lenticulina</i> spp.	—	A	F	—	—	—
<i>Marginulina alazaensis</i> Nuttall	—	F	—	—	—	—
<i>Marginulina dubia</i> Neugeboren	—	F	—	—	—	—
<i>Marginulina subbullata</i> Hantken	—	F	F	—	—	—
<i>Nodosaria longiscata</i> d'Orbigny	—	F	F	—	—	—
<i>Nonionella</i> cf. <i>N. miocenica</i> Cushman	—	F	F	—	—	—
<i>Pseudonodosaria gallowayi</i> (Cushman)	—	F	F	—	—	—
<i>Pseudonodosaria inflata</i> (Bornemann)	—	F	F	A	—	F
<i>Reophax pilulifer</i> H. B. Brady	—	F	A	—	—	—
<i>Saracenaria</i> cf. <i>S. schencki</i> Cushman and Hobson	—	F	A	—	—	—
<i>Siphogenerina nodifera</i> Cushman and Kleinpell	—	F	—	A	A	A
<i>Stilostomella advena</i> (Cushman and Laiming)	—	F	F	—	—	—
<i>Uvigerina auberiana</i> d'Orbigny	—	C	—	—	—	—
<i>Uvigerina gallowayi</i> Cushman	—	R	—	—	—	—
<i>Buliminella subfusiformis</i> Cushman	—	F	—	—	A	—
<i>Stilostomella</i> sp.	—	—	F	—	—	—
<i>Cibicides floridanus</i> (Cushman)	—	—	—	F	—	—
<i>Globocassidulina globosa</i> (Hantken)	—	—	—	R	—	—
<i>Melonis pompilioides</i> (Fichtel and Moll)	—	—	—	R	—	—
<i>Nodosaria parexilis</i> Cushman and Stewart	—	—	—	R	—	—
<i>Sphaeroidina bulloides</i> d'Orbigny	—	—	—	C	—	—
<i>Spiroplectammina</i> sp.	—	—	—	F	—	—
<i>Cibicides</i> sp.	—	—	—	—	R	R
<i>Cyclammmina</i> spp.	—	—	—	—	F	—
<i>?Globobulimina pacifica</i> Cushman	—	—	—	—	F	—
<i>Gyroidina</i> sp.	—	—	—	—	R	—
<i>Marginulina</i> sp.	—	—	—	—	F	—
<i>Marginulina subrecta</i> Franke	—	—	—	—	R	—
<i>Martinottiella patens</i> (Cushman and Laiming)	—	—	—	—	F	—
<i>Orthomorphina rohri</i> (Cushman and Stainforth)	—	—	—	—	F	—
<i>Stilostomella lepidula</i> (Schwager)	—	—	—	—	F	—
<i>Nonionella costifera</i> (Cushman)	—	—	—	—	A	—

TABLE 15.—Benthic foraminifers from the Año Nuevo section

Benthonic foraminiferal stages	Late		Sauce-		Relizian			
	Zemorian		sian					
	Sample							
Taxon	Mf4665	Mf1376	Mf1374	Mf4664	Mf1375	Mf4661	Mf4667	Mf4668
<i>Anomalina californiensis</i> Cushman and Hobson	A	F	—	F	—	—	—	—
<i>Astacolus</i> sp.	F	F	—	F	—	R	—	—
<i>Bulimina carneroensis</i> Cushman and Kleinpell	F	F	—	C	F	—	—	—
<i>Bolivina marginata</i> Cushman	F	F	—	—	—	—	—	—
<i>Cassidulina crassipunctata</i>	—	—	—	—	—	—	—	—
Cushman and Hobson	A	F	F	—	F	—	—	—
<i>Cassidulina margareta</i> Karrer	F	F	—	A	F	—	—	F
<i>Cibicides americanus</i> (Cushman)	R	—	F	—	—	—	—	—
<i>Cibicides</i> cf. <i>C. americanus</i> (Cushman)	R	—	—	—	—	—	—	—
<i>Cibicides floridanus</i> (Cushman)	C	F	—	C	F	—	—	F
<i>Cibicides pseudoungerianus evolutus</i>	—	—	—	—	—	—	—	—
Cushman and Hobson	A	—	F	—	F	—	—	—
<i>Cyclammina cancellata</i> Brady	F	—	—	F	—	—	—	—
<i>Dentalina cooperensis</i> Cushman	R	—	—	F	R	—	—	—
<i>Fursenkoina bramletti</i> (Galloway and Morrey)	F	—	—	F	—	F	—	—
<i>Glandulina laevigata</i> d'Orbigny	R	—	—	—	—	—	—	—
<i>Globobulimina pacifica</i> Cushman	R	—	—	F	F	—	—	—
<i>Gyoidina orbicularis planata</i> Cushman	R	F	—	—	—	—	—	—
<i>Gyroidina soldanii</i> d'Orbigny	A	R	F	F	—	—	—	—
<i>Lagena costata</i> (Williamson)	F	—	—	—	—	—	—	—
<i>Lagena semistriata</i> Williamson	R	—	—	—	—	—	—	—
<i>Lenticulina barbati</i> Cushman	F	—	R	F	—	—	—	—
<i>Lenticulina calcar</i> (Linne)	F	—	—	—	—	—	—	—
<i>Lenticulina</i> cf. <i>L. calcar</i> (Linne)	F	—	R	F	—	—	—	—
<i>Lenticulina</i> cf. <i>L. clypeiformis</i> (d'Orbigny)	F	—	—	F	F	—	—	—
<i>Lenticulina coloratus</i> (Stache)	R	—	—	—	—	—	—	—
<i>Lenticulina pseudorotulata</i> (Asano)	R	—	—	—	—	—	—	—
<i>Lenticulina</i> sp. of Fairchild and others	F	—	—	C	F	—	F	F
<i>Lenticulina</i> spp.	F	—	—	R	R	—	—	—
<i>Marginulina</i> spp.	F	—	F	—	—	—	—	—
<i>Martinottiella patens</i> (Cushman and Laiming)	F	—	—	C	F	—	—	—
<i>Nodosaria paxillilis</i> Cushman and Stewart	F	—	—	—	—	—	—	—
<i>Nodosaria paxillilis sentifera</i>	—	—	—	—	—	—	—	—
Cushman and Parker	F	—	—	—	—	—	—	—
<i>Nonionella incisa kernensis</i> Kleinpell	F	—	—	—	—	—	—	—
<i>Nonionella miocenica</i> Cushman	F	—	—	F	—	—	—	—
<i>Orthomorphina rohri</i> (Cushman and Stainforth)	F	—	—	F	—	—	—	—
<i>Plectofrondicularia miocenica directa</i>	—	—	—	—	—	—	—	—
Cushman and Laiming	C	—	F	F	F	—	—	—
<i>Plectofrondicularia vaughani</i> Cushman	F	—	—	F	F	—	—	—
<i>Pseudonodosaria inflata</i> (Costa)	F	—	—	F	F	—	—	—
<i>Reophax pilulifera</i> Brady	F	—	—	—	—	—	—	—
<i>Reophax</i> spp.	F	—	—	F	—	—	—	—
<i>Siphogenerina nodifera</i> Cushman and Kleinpell	A	C	—	A	F	—	—	—
<i>Stilostomella advena</i> (Cushman and Laiming)	C	—	—	—	—	—	—	—
? <i>Uvigerina gallowayi</i> Cushman	F	—	—	F	—	—	—	—
? <i>Uvigerina</i> sp.	R	—	—	—	—	—	—	—
<i>Verneuilina</i> sp. of Fairchild and others	R	R	—	F	F	—	—	—
<i>Haplophragmoides</i> sp.	R	—	—	—	—	—	—	—
<i>Cibicides americanus crassiseptus</i>	—	—	—	—	—	—	—	—
Cushman and Laiming	F	—	F	F	F	—	—	—
<i>Bathysiphon</i> spp.	—	R	F	—	F	—	—	—
<i>Stilostomella sanctaerucis</i> Kleinpell	—	F	—	R	F	—	—	—
<i>Trochammina</i> spp.	—	—	—	—	—	—	—	—
<i>Ammodiscus</i> cf. <i>A. incertus</i> d'Orbigny	—	—	F	—	—	—	—	—
<i>Bolivina</i> spp.	—	—	F	F	F	F	F	F
<i>Bulimina inflata alligata</i> Cushman and Laiming	—	—	F	F	F	—	—	—
<i>Buliminella curta</i> Cushman	—	—	F	—	F	—	—	—
<i>Cyclammina</i> spp.	—	—	F	—	—	—	—	—
<i>Dentalina quadrulata</i> Cushman and Laiming	—	—	F	F	F	—	—	—
<i>Lenticulina</i> sp. A	—	—	R	F	—	—	—	—
<i>Melonis pompilioides</i> (Fichtel and Moll)	—	—	R	F	F	—	—	—
<i>Nodosaria longiscata</i> d'Orbigny	—	—	F	F	—	—	—	—
<i>Nodosaria pyrula</i> d'Orbigny	—	—	F	—	F	—	—	—
<i>Oridorsalis umbonatus</i> (Reuss)	—	—	F	—	F	—	—	—
<i>Plectofrondicularia californica</i>	—	—	—	—	—	—	—	—
Cushman and Stewart	—	—	F	—	F	—	—	—
<i>Pullenia multiobata</i> Chapman	—	—	R	F	—	—	—	—
<i>Siphogenerina multicostata</i> Cushman and Jarvis	—	—	C	R	—	—	—	—
<i>Stilostomella</i> sp.	—	—	F	—	—	—	—	—
<i>Sphaeroidina bulloides</i> d'Orbigny	—	—	F	C	F	—	—	—
<i>Uvigerina auberiana</i> d'Orbigny	—	—	F	C	F	A	—	F
<i>Baggina robusta</i> Kleinpell	—	—	—	—	—	A	—	F
<i>Bolivina marginata adelaidana</i>	—	—	—	—	—	—	—	—
Cushman and Kleinpell	—	—	—	F	—	—	—	—
<i>R. ...</i>	—	—	—	R	—	—	—	—
<i>R. ...</i>	—	—	—	R	—	—	—	—
<i>C. ...</i>	—	—	—	F	—	—	—	—
<i>Gaudryina gracilis</i> Cushman and Laiming	—	—	—	F	—	—	—	—
<i>Lagena acuticosta</i> (Reuss)	—	—	—	F	—	—	—	—
<i>Lagena hexagona</i> (Williamson)	—	—	—	F	—	—	—	—
<i>Lagena vulgaris</i> Williamson	—	—	—	F	—	—	—	—
<i>Lenticulina pseudocultratus</i> (Cole)	—	—	R	F	—	—	—	—
<i>Lenticulina pseudovortex</i> (Cole)	—	—	—	—	—	—	—	—
<i>Verneuilina advena</i> (Costa)	—	—	R	—	—	—	—	—
<i>P. ...</i>	—	—	—	F	—	—	—	—
<i>P. ...</i>	—	—	—	R	F	C	—	—
<i>P. ...</i>	—	—	—	F	—	F	—	—
<i>Saracenaria</i> cf. <i>S. schencki</i>	—	—	—	—	—	—	—	—
Cushman and Hobson	—	—	—	F	R	—	—	—
<i>Siphogenerina mayi</i> Cushman and Parker	—	—	F	F	—	—	—	—
<i>Stilostomella adolphina</i> (d'Orbigny)	—	—	—	C	F	—	—	—
<i>S. ...</i>	—	—	—	F	—	—	—	—
<i>S. ...</i>	—	—	—	F	—	—	—	—
<i>S. ...</i>	—	—	—	F	—	—	—	—

TABLE 15.—Continued

Benthonic foraminiferal stages	Late Zemorrian			Sauce-sian		Relizian		
	Sample							
Taxon	Mf4665	Mf1376	Mf1374	Mf4664	Mf1375	Mf4661	Mf4667	Mf4668
<i>Bolivina advena</i> Cushman	—	—	—	—	R	—	—	R
<i>Bulimina</i> spp.	—	—	—	—	—	—	—	—
<i>Chilostomella</i> sp.	—	—	—	—	—	—	—	—
<i>Pseudonodosaria gallowayi</i> (Cushman)	—	—	—	—	—	F	—	—
<i>Siphogenerina transversa</i> Cushman	—	—	—	—	—	F	—	—
<i>Cassidulina</i> sp.	—	—	—	—	—	A	—	A
<i>Epistominella</i> spp.	—	—	—	—	—	—	F	—
<i>Gyroidina</i> sp.	—	—	—	—	—	—	F	—
<i>Siphogenerina hughesi</i> Cushman	—	—	—	—	—	—	—	F
<i>Elphidium</i> sp.	—	—	—	—	—	—	—	F
<i>Nodosaria</i> sp.	—	—	—	—	—	—	—	F
<i>Nonionella costifera</i> (Cushman)	—	—	—	—	—	—	—	F
<i>Valvulineria californica</i> variations	—	—	—	—	—	—	—	F

fore the early Zemorrian-late Zemorrian boundary is placed between Mf4682 and Mf4681. The late Zemorrian assemblages are dominated by *Siphogenerina nodifera*, *Uvigerinella sparsicostata*, and several agglutinated species as well as a number of species which first appear in the Zemorrian, including *Lenticulina simplex*, *Buliminella subfusiformis*, *Sphaeroidina bulloides*, and *Cibicides floridanus*.

The last sample, Mf4671, is from the Lambert Shale. Few of the species from the underlying samples occur but *Nonionella costifera* is present. The first occurrence of *Nonionella costifera* and several other species was considered by Kleinpell (1938) to mark the base of the Saucian Stage. As none of the other species occur in Mf4671, it is only questionably assigned to the Saucian Stage.

AÑO NUEVO SECTION

The Año Nuevo section is compiled from strata exposed in the sea cliffs north and east of Año Nuevo Point (fig. 27). Samples Mf1374-Mf1376 (north of Año Nuevo Point) were previously reported on by Clark (in Brabb and others, 1977). Both the Vaqueros(?) and Monterey Formations were sampled; however, only assemblages from the Vaqueros(?) Formation are discussed here (table 15). Localities and sedimentological and structural relations are described in Brabb, Clark, and Throckmorton (1977). See table 13 for field numbers and corresponding Mf numbers used in this report.

The lowest samples in this section, Mf4665 and Mf1376 (fig. 32), are representative of the upper Zemorrian *Uvigerinella sparsicostata* Zone. *Bolivina marginata* and *Siphogenerina nodifera* are still present, but *Uvigerinella sparsicostata* is not. Species diagnostic of the late Zemorrian include *Bulimina carneroensis*, *Cassidulina margareta*, *Nonionella incisa*

kernensis, *Siphogenerina mayi*, and *Siphogenerina multicostata*. These species are present in Mf4664, stratigraphically higher in the section but in association with species diagnostic of the Saucesian Stage such as *Baggina robusta*, *Bolivina marginata adalaidana*, *Bulimina inflata alligata*, and *Dentalina quadrulata*.

Kleinpell (1938) indicated that *Siphogenerina mayi*, though diagnostic of the late Zemorrian, could be found in the early Saucesian. He subsequently recognized a similar range for *Siphogenerina multicostata* (Boris Laiming, oral commun., 1972). The fauna represented in Mf4664 is similar to the assemblages of the San Joaquin Valley (Tipton and others, 1973), where the bathymetric conditions remain unchanged across the Zemorrian-Saucesian boundary. The result is a gradual change in the fauna and a mixing of the Zemorrian and Saucesian species. The boundary is placed at sample Mf4664 with the full realization that the precise location cannot be defined as it is not known when the mixing of faunas occurred. The slight range extension of *Bulimina carneroensis* and *Siphogenerina multicostata* should be noted. Sample Mf1374 is probably also in this zone, but the faunas are less diverse.

Assemblages diagnostic of the Saucesian Stage occur in the sheared section in the upper part of the Vaqueros(?) Formation. Samples Mf4661 and Mf2188-Mf2191 of Clark (*in* Brabb and others, 1977), from this interval, are poorly preserved. Diagnostic species present include many of the Saucesian species discussed earlier and *Siphogenerina transversa*. This species ranges from the Zemorrian and Saucesian into Miocene faunas but is most characteristic of the Saucesian.

Samples Mf4667 and Mf4668, also from the sheared section but higher than Mf4661, contain only siliceous molds of foraminifers. Many of these can be identified to species. Present in this sample are *Siphogenerina hughesi* and *Valvulinaria californica*, which are diagnostic of the Relizian Stage. Since the section is sheared and the faunas are only represented by siliceous molds, interpretations in this interval are highly speculative.

SUMMARY

Correlation of faunal assemblages in the three sections yields a nearly continuous sequence from the lower Narizian Stage to the Relizian Stage (fig. 33). Stage boundaries are recognized by distinctive benthic foraminiferal assemblages associated with the base of each new stage (fig. 34) or by the gradual change in the faunas. Zonal and subzonal boundaries are less easily recognized because faunal changes are subtle and more frequently recognized by changes in the dominance of species.




FIGURE 33.—Correlation of the San Lorenzo River, Zayante Creek, and Año Nuevo sections with California benthic foraminiferal zones and stages. Open triangles, Church Creek section, Santa Lucia Range (Poore and Bukry, this volume). Filled triangles, San Lorenzo River, Zayante Creek, or Año Nuevo sections (Poore and Bukry, this volume).




FIGURE 34.—Ranges of species diagnostic of middle Tertiary benthic foraminiferal stages (Narizian through Relizian Stages) and zones of the Santa Cruz Mountains, California.

The Narizian Stage is represented in the San Lorenzo section by assemblages of the lower and upper Narizian *Bulimina corrugata* and *Amphimorphina jenkinsi* Zones. Recognition of these zones requires no modification of the original description of the zones. The lower Narizian-upper Narizian boundary corresponds to the Zone P 14 - P 15 boundary and the *Discoaster saipanensis* Zone-*Discoaster barbadensis* Zone boundary (Poore and Bukry, this volume). This boundary corresponds to the middle Eocene - upper Eocene boundary.

In the San Lorenzo River section, the Refugian Stage is marked by an abrupt faunal and lithologic change, associated with the onset of shallower water conditions. The extended range of several Refugian species, *Boldia hodgei* and *Plectofrondicularia packardii*, noted by previous workers and in the modified description of the Refugian Stage (Donnelly, 1976), is confirmed in this section. The Refugian faunas are poorly preserved, but they do suggest that additional revisions are needed in the definition of diagnostic Refugian faunas. The Refugian faunas in the San Lorenzo River section represent a deeper bathymetric facies than those in the type area in California. They resemble faunas in the Lincoln Creek Formation of Washington. The recognition of a bathyal Refugian fauna requires the downward range extension of *Cibicides elmaensis* and various members of the *Uvigerina cocoaensis* species group, along with the addition of diagnostic species such as *Karreriella washingtonensis*, *Saracenaria schencki*, and *Sigmomorphina schencki*. Also since the paleoecologic facies present in the San Lorenzo section are different than the facies of the type area (Cañada de Santa Anita), many of the diagnostic late Refugian species do not appear.

Planktic microfossils from the San Lorenzo River section and correlative sections suggest that the Refugian-Zemorian boundary lies between Zone P 17 and Zone P 19 and between the *Discoaster barbadensis* Zone and *Sphenolithus distentus* Zone (Poore and Bukry, this volume). Neither nannofossils nor planktic foraminifers were sufficiently diverse or abundant in this section to locate the boundary more closely.

The Zemorian Stage can be recognized in all three sections. The criteria for recognition of the Zemorian Stage and the Zemorian zones (*Uvigerina gallowayi* and *Uvigerinella sparsicostata*) need little modification from the original description of Kleinpell (1939). Both the early Zemorian-late Zemorian and the Zemorian-Saucian boundaries are marked by the presence in a sample of species of both zones or stages. This mixing of faunas would be expected in

closely sampled intervals and where environmental conditions are relatively consistent across the boundary. Early and late Zemorrian species are present in sample Mf4682 in the Zayante Creek section, and both Zemorrian and Saucian species are present in Mf4664 in the Año Nuevo section.

Planktic foraminifers diagnostic of Zones P 19 and P 20 and nannofossils diagnostic of the *Sphenolithus distentus* Zone are associated with the early Zemorrian. The *Sphenolithus distentus* Zone and *Sphenolithus ciperoensis* Zone and Zone P 22(?) (Poore and Bukry, this volume) are associated with the late Zemorrian. In the Año Nuevo section, upper Oligocene *Sphenolithus ciperoensis* Zone (sample Mf4664) and Miocene *Triquetrorhabdulus carinatus* Zone (sample Mf2188) nannofossils are recognized (Poore and Bukry, this volume). These zone and age assignments coupled with the benthic foraminiferal data support the interpretation that Mf4664 and Mf1376 straddle the Zemorrian-Saucian boundary.

Faunas diagnostic of the Saucian Stage occur in one sample (Año Nuevo section) and questionably in a second (Zayante Creek section). Nannofossils associated with this interval indicate that the Saucian Stage is Miocene (Poore and Bukry, this volume).

Faunas of Relizian age in the Año Nuevo section are correlative with the *Helicopontosphaera ampliata* or *Sphenolithus heteromorphus* Zone (Poore and Bukry, this volume) and therefore are also Miocene.

REFERENCE LIST OF TAXA

- Ammodiscus incertus* (d'Orbigny) = *Operculina incerta* d'Orbigny, 1839, in de la Sagra, 1839, Histoire physique naturelle de l'Île de Cuba: Paris, A. Bertrand, p. 49, p. 6, figs. 16-17.
- Amphimorphina becki* Mallory, 1959, Lower Tertiary biostratigraphy of the California Coast Ranges: Tulsa, Okla., Am. Assoc. Petroleum Geologists, p. 215, pl. 19, figs. 1a-b.
- Anomalina californiensis* Cushman and Hobson, 1935, Cushman Lab. Foram. Research Contr., v. 11, pt. 3, p. 64, pl. 9, figs. 8a-c.
- Anomalina salinasensis* Klempell, 1938, Miocene stratigraphy of California: Tulsa, Okla., Am. Assoc. Petroleum Geologists, p. 377, pl. VIII, figs. 12a-c.
- Anomalina garzaensis* Cushman and Siegfus, 1939, Cushman Lab. Foram. Research Contr., v. 15, pt. 2, p. 32, pl. 7, figs. 3a-c.
- Asterigerina crassaformis* Cushman and Siegfus, 1935, Cushman Lab. Foram. Research Contr., v. 11, pt. 4, p. 94, pl. 14, fig. 10—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 282, pl. 20, figs. 3a, b, c.
- Baggina robusta* Klempell 1938, Miocene stratigraphy of California: Tulsa, Okla., Am. Assoc. Petroleum Geologists, p. 325, pl. XI, figs. 8a-c.
- Bathysiphon eocenica* Cushman and Hanna, 1927, California Acad. Sci. Proc., ser. 4, v. 16, no. 8, p. 210, pl. 13, figs. 2, 3—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 249, pl. 1, figs. 2, 3.
- Bifarina eleganta* (Plummer) = *Siphogenerina eleganta* Plummer, 1926, Texas Univ. Bull. 2644, p. 126, pl. 8, figs. 1a-c.—*Bifarina eleganta* (Plummer) of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 276, pl. 15, figs. 11a, b.
- Boldia hodgei* (Cushman and Schenck) = *Cibicides hodgei* Cushman and Schenck, 1928, California Univ. Pubs., Dept. Geol. Sci. Bull., v. 17, no. 9, p. 315, pl. 45, figs. 3-5.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 286, pl. 23, figs. 8a, b, c.—*Cibicides cushmani* Nuttall of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 287, pl. 23, figs. 7a, b, c.
- Bolivina advena* Cushman, 1925, Cushman Lab. Foram. Research Contr., v. 1, pt. 2, p. 29, pl. 5, figs. 1a-b.
- Bolivina gracilis* Cushman and Applin, 1926, Am. Assoc. Petroleum Geologists Bull., v. 10, pt. 1, p. 167, pl. 7, figs. 1, 2.
- Bolivina klempelli* Beck, 1943, Jour. Paleontology, v. 17, no. 6, p. 606, pl. 107, figs. 3, 9.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 276, pl. 15, figs. 6a, b.
- Bolivina marginata* Cushman, 1925, Cushman Lab. Foram. Research Contr., v. 1, pt. 2, p. 30, pl. 5, figs. 5a-b.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 276, pl. 15, figs. 9a, b.
- Bolivina scabrata* Cushman and Bermudez, 1936, Cushman Lab. Foram. Research Contr., v. 12, pt. 2, p. 29, pl. 5, figs. 11-12.
- Bolivinoidea mexicanus* (Cole) = *Proropus mexicanus* Cole, 1927, Bulls. Am. Paleontology, v. 14, pt. 51, p. 26, pl. 1, fig. 19.
- Bulimina carneroensis* Cushman and Klempell, 1934, Cushman Lab. Foram. Research Contr., v. 10, pt. 1, p. 5, pl. 1, figs. 12a, b.
- Bulimina corrugata* Cushman and Siegfus, 1935, Cushman Lab. Foram. Research Contr., v. 11, pt. 4, p. 92-93, pl. 14, figs. 7a-b.
- Bulimina curtissima* Cushman and Siegfus, 1935, Cushman Lab. Foram. Research Contr., v. 11, pt. 4, p. 93, pl. 14, figs. 9a, b.
- Bulimina inflata alligata* Cushman and Laiming, 1931, Jour. Paleontology, v. 5, no. 2, p. 107, pl. 11, figs. 17a-b.
- Bulimina microcostata* Cushman and Parker, 1936, Cushman Lab. Foram. Research Contr., v. 12, pt. 2, p. 39, pl. 7, fig. 2.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 274, pl. 14, figs. 6a, b.
- Bulimina sculptilis lacinata* Cushman and Parker, 1937, Cushman Lab. Foram. Research Contr., v. 13, pt. 2, p. 46-53, pls. 5-6.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 274, pl. 14, figs. 8a, b.
- Buliminella curta* Cushman, Lab. Foram. Research Contr., v. 1, pt. 2, p. 33, pl. 29, fig. 4.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 273, pl. 14, figs. 1a, b.
- Buliminella elegantissima* (d'Orbigny) = *Bulimina elegantissima* d'Orbigny, 1839, Voyage dans l'Amerique Meridionale, v. 5, pt. 5, pl. 51, pl. 7, figs. 13, 14.
- Buliminella subfusiformis* Cushman, 1925, Cushman Lab. Foram. Research Contr., v. 1, pt. 2, p. 33, pl. 5, fig. 12.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 273, pl. 14, figs. 2a, b, 3.
- Cassidulina crassipunctata* Cushman and Hobson, 1935, Cushman Lab. Foram. Research Contr., v. 11, pt. 3, p. 63, pl. 9, fig. 10.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 282, pl. 20, figs. 6a, b.
- Cassidulina diversa* Cushman and Stone, 1949, Cushman Lab. Foram. Research Contr., v. 25, pt. 3, p. 56.
- Cassidulina margareta* Karrer, 1877, K. K. Geol. Reichsanst., Abh., Wien, Bd. 9, p. 386.
- Cassidulinoides californiensis* Bramlette, 1951, in Woodring and Bramlette, 1951, U.S. Geol. Survey Prof. Paper 222, p. 61.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 283, pl. 20, figs. 9a, b.
- Cibicides americanus* (Cushman) = *Truncatulina americana* Cushman, 1918, U.S. Geol. Survey Bull. 676, p. 63.

- Cibicides americanus crassiseptus* Cushman and Laiming, 1931, Jour. Paleontology, v. 5, no. 2, p. 119, pl. 14, figs. 14a-c.
- Cibicides elmaensis* Rau, 1948, Jour. Paleontology, v. 22, no. 2, p. 173, pl. 31, figs. 18-26.
- Cibicides falconensis* Renz, 1948, Geol. Soc. America Bull., v. 32, p. 128, pl. 11, figs. 6a-c, 7.
- Cibicides floridanus* (Cushman) = *Truncatulina floridana* Cushman, 1918, U.S. Geol. Survey Bull. 676, p. 62.
- Cibicides haydoni* (Cushman and Schenck) = *Planulina haydoni* Cushman and Schenck, 1928, California Univ. Pubs., Dept. Geol. Sci. Bull., v. 17, no. 9, p. 316, pl. 45, figs. 7a-c.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 286, pl. 23, figs. 1a, b, c, 2.
- Cibicides pachyderma* (Rzehak) = *Truncatulina pachyderma* Rzehak, 1885, Naturf. Ver. Brunn, Verh., Brunn, Bd. 24, (1885), p. 87.
- Cibicides pseudoungerianus evolutus* Cushman and Hobson, 1935, Cushman Lab. Foram. Research Contr., v. 11, no. 3, p. 64, pl. 9, fig. 11.
- Cibicides spiropunctatus* Galloway and Morrey, 1931, Jour. Paleontology, v. 5, p. 4, p. 346, pl. 39, fig. 7.
- Cyclammina cancellata obesa* Cushman and Laiming, 1931, Jour. Paleontology v. 5, no. 2, p. 94, pl. 9, figs. 10a, b.
- Cyclammina incisa* (Stache) = *Haplophragmium incisum* Stache, 1865, Novora Exped. 1857-1859, Wien. Osterreich. Geol. Theil. Bd. 1, Abt. 2, p. 165.
- Cyclammina pacifica* Blaisdell, 1943, Jour. Paleontology, v. 17, no. 6, p. 591, pl. 98, figs. 2-3.
- Dentalina cocoaensis* (Cushman) = *Nodosaria cocoaensis* Cushman, 1925, Cushman Lab. Foram. Research Contr., v. 1, pt. 3, p. 66.
- Dentalina cooperensis* Cushman, 1933, Cushman Lab. Foram. Research Contr., v. 9, pt. 1, p. 8, pl. 1, fig. 17.
- Dentalina jacksonensis* Cushman and Applin, 1926, Am. Assoc. Petroleum Geologists Bull., v. 10, no. 2, p. 170, pl. 7, figs. 14-16.
- Dentalina quadrolata* Cushman and Laiming, ms. Cushman and Parker, 1931, Cushman Lab. Foram. Research Contr., v. 7, pt. 1, p. 3-4, pl. 1, figs. 9-11.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 264, pl. 9, figs. 13, 14a, b.
- Dentalina spinosa ornatior* Smith, 1957, California Univ. Pubs. Geol. Sci., v. 32, no. 3, p. 165.
- Dorothia cubana* (Cushman and Bermudez) = *Gaudryina cubana* Cushman and Bermudez, 1936, Cushman Lab. Foram. Research Contr., v. 12, pt. 3, p. 56, pl. 10, figs. 2, 10, 11.
- Eggerella elongata* Blaisdell, 1965, Cushman Found. Foram. Research Contr., v. 16, no. 1, p. 27, pl. 2, figs. 1-3.
- Eggerella subconica* Parr, 1950, British and New Zealand Antarctic Research Exped. 1929-1931, Repts. Adelaide, ser. B, v. 5, pt. 6, p. 281, pl. 5, figs. 22a, b.
- Fronicularia tenuissima* Hantken, 1875, K. Ungar. Geol. Anst., Mitt. Jahrb., Bd. 4, Heft. 1, p. 43.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 267, pl. 11, fig. 6.
- Fursenkoina bramletti* (Galloway and Morrey) = *Virgulina bramletti* Galloway and Morrey, 1929, Bull. Am. Paleontology, v. 15, no. 55, p. 37, pl. 5, figs. 14a, b.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 275, pl. 15, figs. 1a, b, 2.
- Fursenkoina californiensis* (Cushman) = *Virgulina californiensis* Cushman, 1925, Cushman Lab. Foram. Research Contr., v. 1, pt. 2, p. 32, pl. 5, figs. 11a-c.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 275, pl. 15, figs. 3, 4a, b.
- Fursenkoina dibollensis* (Cushman and Applin) = *Virgulina dibollensis* Cushman and Applin, 1926, Am. Assoc. Petroleum Geologists Bull., v. 10, pt. 1, no. 2, p. 168.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 275, pl. 15, figs. 5a, b.
- Gaudryina triangularis* Cushman, 1911, U.S. Natl. Mus. Bull. 71, p. 65.
- Glandulina laevigata* (d'Orbigny) = *Nodosaria (Glandulina) laevigata* d'Orbigny, 1826, Ann. Sci. Nat. Paris, ser. 1, tome 7, p. 252.
- Globobulimina pacifica* Cushman, 1927, Cushman Lab. Foram. Research Contr., v. 3, pt. 2, p. 67, pl. 14, fig. 12.
- Globocassidulina globosa* (Hantken) = *Cassidulina globosa* Hantken, 1875, K. Ungar. Geol. Anst. Mitt. Jahrb., Bd. 4, Heft. 1, p. 64, pl. 16, figs. 2a-b.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 283, pl. 20, figs. 5a, b.
- Guttulina frankei* Cushman and Ozawa, 1930, U.S. Natl. Mus. Proc., v. 77, no. 2829, art. 6, p. 28, pl. 30, figs. 17, 18.
- Guttulina irregularis* (d'Orbigny) = *Globulina irregularis* d'Orbigny, 1846, Foraminifères fossiles du bassin tertiaire de Vienne (Autriche): Paris, Gide et. Cie., p. 226, pl. 13, figs. 9, 10.—*Guttulina irregularis* (d'Orbigny) of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 267, pl. 11, figs. 8a, b, c.
- Gyroidina condoni* (Cushman and Schenck) = *Eponides condoni* Cushman and Schenck, 1928, California Univ. Pubs., Dept. Geol. Sci., Bull., v. 17, no. 9, p. 313, pl. 44, figs. 6, 7a-c.—*Gyroidina condoni* (Cushman and Schenck) of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 280, pl. 18, figs. 2a, b, c.
- Gyroidina orbicularis planata* Cushman, 1935, U.S. Geol. Survey Prof. Paper 181, p. 45, pl. 66, figs. 4-6.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 280, pl. 18, figs. 1a, b, c.
- Gyroidina soldanii* d'Orbigny, 1926, Ann. Sci. Nat. Paris, ser. 1, tome 7, p. 45, pl. 18, fig. 3.
- Haplophragmoides deflata* Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 251, pl. 1, figs. 11, 12.
- Karrerella elongata* Mallory, 1959, Lower Tertiary biostratigraphy of the California Coast Ranges: Tulsa, Okla., Am. Assoc. Petroleum Geologists, p. 127, pl. 5, figs. 4a-c.
- Karrerella washingtonensis* Rau, 1948, Jour. Paleontology, v. 22, pt. 2, p. 158, pl. 27, figs. 5, 6.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 255, pl. 4, figs. 3a, b, 4a, b.
- Lagena acuticosta* Reuss, 1862, Akad. Wiss. Sitz., v. 44, pt. 1, 1861, p. 303, pl. 1, fig. 4.
- Lagena becki* Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 266, pl. 10, figs. 16a, b.
- Lagena costata* (Williamson) = *Entosolenia costata* Williamson, 1858, Ann. Mag. Nat. History (London), ser. 2, v. 1, p. 9, pl. 1, fig. 18.
- Lagena hexagona* (Williamson) = *Entosolenia squamosa* (Montagu) var. *hexagona* Williamson, 1848, Ann. Mag. Nat. History (London), ser. 2, v. 1, p. 20, pl. 2, figs. 9a-b.
- Lagena semistriata* Williamson, 1848, Ann. Mag. Nat. History (London), ser. 2, v. 1, p. 14, pl. 1, figs. 9-10.
- Lagena sulcata* (Walker and Jacob) = *Serpula sulcata* Walker and Jacob, 1798, in Kanmacher, 1798, Adam's essays on the microscope, ed. 2, London, p. 634.
- Lagena vulgaris* Williamson, 1858, Royal Soc. London, p. 3, pl. 1, figs. 5, 5a.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 267, pl. 10, figs. 12a, b.
- Lenticulina barbati* (Cushman and Hobson) = *Robulus barbati* Cushman and Hobson, 1935, Cushman Lab. Foram. Research Contr., v. 11, pt. 3, p. 57, pl. 8, figs. 9a-b.
- Lenticulina calcar* (Linne) = *Nautilus calcar* Linne, 1758, Systema naturae, ed. 10: Holmiae, Suecia, impensis L. Salvii, tomus 1, p. 709.
- Lenticulina chehalisensis* (Rau) = *Robulus chehalisensis* Rau, 1948, Jour. Paleontology, v. 22, no. 2, p. 162, pl. 29, figs. 14-15.

- Lenticulina* cf. *L. clypeiformis* d'Orbigny, Kleinpell, 1938, Miocene stratigraphy of California: Tulsa, Okla., Am. Assoc. Petroleum Geologists, p. 197, pl. 3, fig. 7.
- Lenticulina coaledensis* (Detling) = *Robulus coaledensis* Detling, 1946, Jour. Paleontology, v. 20, pt. 4, p. 353, pl. 48, fig. 1.
- Lenticulina cultratus* (Montfort) = *Robulus cultratus* Montfort, 1808, Conchyliologie systematique et classification methodique des coquilles: Paris, France, tome 1, p. 215.
- Lenticulina inornata* (d'Orbigny) = *Robulus inornata* d'Orbigny, 1846, Foraminiferes fossiles du bassin tertiaire de Vienne: Paris, France, Gide et Cie., p. 102, pl. 4, figs. 25, 26.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 258, pl. 5, figs. 5a, b.
- Lenticulina insuetus* (Cushman and Stainforth) = *Robulus insuetus* Cushman and Stainforth, 1947, Cushman Lab. Foram. Research Contr., v. 23, pt. 4, p. 77, pl. 17, fig. 1.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 259, pl. 5, figs. 6, 7a, b.
- Lenticulina midwayensis* (Plummer) = *Cristellaria midwayensis* Plummer, 1926, Texas Univ. Bull. 2644, p. 95, pl. 13, fig. 5.—(not *Robulus midwayensis* of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 259, pl. 6, figs. 3a, b).
- Lenticulina pseudocultratus* (Cole) = *Robulus pseudocultratus* Cole, 1927, Bulls. Am. Paleontology, v. 14, no. 51, p. 19.
- Lenticulina pseudorotulata* (Asano) = *Robulus pseudorotulatus* Asano, 1938, Tohoku Imp. Univ., Sci. Repts., ser. 2, v. 19, no. 2, p. 201.
- Lenticulina pseudovortex* (Cole) = *Robulus pseudovortex* Cole, 1927, Bull. Am. Paleontology, v. 14, no. 1, p. 19.
- Lenticulina simplex* (d'Orbigny) = *Cristellaria simplex* (d'Orbigny, 1846, Foraminiferes fossiles du bassin Tertiaire du Vienne: Paris, France, Gide et Cie., p. 85.—*Robulus simplex* d'Orbigny of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 260, pl. 5, figs. 8a, b).
- Lenticulina* sp. of Fairchild and others, 1969, California Univ. Pubs. Geol. Sci., v. 81, p. 44, pl. 7, figs. 9a, b.
- Lenticulina welchi* (Church) = *Robulus welchi* Church, 1931, Mining in California, v. 27, no. 2, p. 212, pl. C, figs. 13-14.—*Robulus chiranus* Cushman and Stone of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 257, pl. 7, figs. 1, 2a, b.
- Marginulina adunca* (Costa) = *Glandulina adunca* Costa, 1856, Accad. Pontaniana Napoli, Atti., w. 7, fasc. 2, p. 128, pl. 11, figs. 24a, A.C.
- Marginulina alazaensis* Nuttall, 1932, Jour. Paleontology, v. 6, p. 13, pl. 3, figs. 3, 7.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 261, pl. 8, figs. 7a, b.
- Marginulina dubia* Neugeboren, 1851, Siebenb. ver. Naturw. Hermannstadt, verh. Mitt., Hermannstadt, Ungarn, Jahrg. 2, no. 7, p. 120.
- Marginulina exima* Neugeboren, 1851, Siebenb. ver. Naturw. Hermannstadt, verh. Mitt., Hermannstadt, Ungarn, Jahrg. 2, no. 8, p. 129.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 262, pl. 8, figs. 9a, b.
- Marginulina subbullata* Hantken, 1875, Hungary, K. Ungar. Geol. Anst., Mitt. Jahrb., Bd. 4, Heft 1, p. 46.
- Marginulina subrecta* Franke, 1927, Danmarks, Geol. Undersogelse Skr., Raekke 2, Nr. 46, p. 19.
- Martinottiella eocenica* Cushman and Bermudez, 1937, Cushman Lab. Foram. Research Contr., v. 13, pt. 1, p. 6, pl. 1, figs. 27-28.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 255, pl. 4, figs. 6a, b, c.
- Martinottiella patens* (Cushman and Laiming) = *Clavulina patens* Cushman and Laiming, 1931, Jour. Paleontology, v. 5, no. 2, p. 96, pl. 10, figs. 2a-b.
- Melonis pompilioides* (Fichtel and Moll) = *Nautilus pompilioides* Fichtel and Moll, 1798, Testacea microscopica aliaque minuta ex generibus Argonata et Nautilus, Wien, p. 31.
- Nodosarella atlantisae hispidula* (Cushman) = *Ellipsonodosaria atlantisae hispidula* Cushman, 1939, Cushman Lab. Foram. Research Contr., v. 15, pt. 3, p. 70, pl. 12, figs. 3, 4.
- Nodosaria boffalorae* Martinotti, 1924, Soc. Italiana Sci. Nat. e Museo Civico Storia Nat. Milano Atti, v. 62, fasc. 3-4, p. 333.
- Nodosaria holserica* Schwäger, 1866, Novara Exped. 1857-1859, Wien, Geol. Theil, Bd. 2, Abt. 2, p. 221.
- Nodosaria longiscata* d'Orbigny, 1846, Foraminiferes fossiles du bassin tertiaire du Vienne: Paris, France, Gide et Cie., p. 32, pl. 1, figs. 10-12.
- Nodosaria parexilis sentifera* Cushman and Parker, 1931, Cushman Lab. Foram. Research Contr., v. 7, pt. 1, p. 6, pl. 1, fig. 16.
- Nodosaria pyrula* d'Orbigny, 1826, Ann. Sci. Nat. Paris, ser. 1, tome 7, p. 253.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 265, pl. 10, figs. 3a-b.
- Nonion halkyardi* Cushman, 1936, Cushman Lab. Foram. Research Contr., v. 12, pt. 3, p. 63, pl. 12, fig. 1.
- Nonionella costifera* (Cushman) = *Nonionina costifera* Cushman, 1926, Cushman Lab. Foram. Research Contr., v. 1, no. 4, p. 90, pl. 13, figs. 2a-c.
- Nonionella incisa* (Cushman) = *Nonionina incisa* Cushman, 1926, Cushman Lab. Foram. Research Contr., v. 1, no. 4, p. 90-91, pl. 13, figs. 3a-c.
- Nonionella incisa kernensis* (Kleinpell) = *Nonion incisum* (Cushman) var. *kernensis* Kleinpell, 1938, Miocene stratigraphy of California: Tulsa, Okla., Am. Assoc. Petroleum Geologists, p. 232.
- Nonionella miocenica* Cushman, 1926, Cushman Lab. Foram. Research Contr., v. 1, no. 4, p. 91, pl. 13, figs. 4a-c.
- Oridorsalis umbonatus* (Reuss) = *Rotalina umbonata* Reuss, 1851, Deutsch. Geol. Gesell. Zeitschr., Bd. 3, p. 75, pl. 5, figs. 35.—*Eponides umbonatus* (Reuss) of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 281, pl. 18, figs. 7a, b, c.
- Orthomorphina rohri* (Cushman and Stainforth) = *Nodogenerina rohri* (Cushman and Stainforth, 1945, Cushman Lab. Foram. Research Contr., Spec. Pub. 14, p. 39, pl. 5, fig. 26.
- Planularia markleyana* Church, 1941, California State Mineralogist, v. 27, no. 2, p. 208, pl. A, fig. 6, pl. 8, figs. 1, 10.
- Plectina garzaensis* Cushman and Siegfus, 1935, Cushman Lab. Foram. Research Contr., v. 11, no. 6, p. 92, pl. 14, figs. 3, 4.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 255, pl. 3, figs. 8a, b, c, 9.
- Plectofrondicularia californica* Cushman and Stewart, 1926, Cushman Lab. Foram. Research Contr., v. 2, pt. 2, p. 39, pl. 6, figs. 9-11.
- Plectofrondicularia miocenica* Cushman, 1926, Cushman Lab. Foram. Research Contr., v. 2, pt. 3, p. 58, pl. 7, figs. 10, 11, pl. 8, figs. 11, 12.
- Plectofrondicularia miocenica directa* Cushman and Laiming, 1931, Jour. Paleontology, v. 5, no. 2, p. 105, pl. 11, fig. 12.
- Plectofrondicularia packardi* Cushman and Schenck, 1928, California Univ. Pubs., Dept. Geol. Sci. Bull., v. 17, no. 9, p. 311, pl. 43, figs. 14, 15.—*Plectofrondicularia packardi packardi* Cushman and Schenck of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 270, pl. 12, figs. 11, 12.
- Plectofrondicularia packardi multilineata* Cushman and Simonson, 1944, Jour. Paleontology, v. 18, p. 197, pl. 32, figs. 2-4.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 270, pl. 12, fig. 10.
- Plectofrondicularia vaughani* Cushman, 1927, Cushman Lab. Foram. Research Contr., v. 3, p. 2, p. 112, pl. 23, fig. 3.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p.

- 271, pl. 12, fig. 7.
- Pleurostomella alternans* Schwager, 1866, Novara Exped. 1857-1859, Wien, Geol. Theil, Bd. 2, Abt. 2, p. 238.
- Praeglobobulimina ovata* (d'Orbigny) = *Bulimina ovata* d'Orbigny, 1846, Foraminifères fossiles du bassin tertiaire de Vienne: Paris, France, Gide et Cie., p. 186, pl. 11, figs. 13-14.
- Praeglobobulimina pupoides* d'Orbigny = *Bulimina pupoides* d'Orbigny, 1846, Foraminifères fossiles du bassin tertiaire de Vienne: Paris, France, Gide et Cie., p. 185, pl. 11, figs. 11, 12.
- Pseudonodosaria gallowayi* (Cushman) = *Pseudoglandulina gallowayi* Cushman, 1929, Cushman Lab. Foram. Research Contr., v. 5, pt. 4, no. 84, p. 87.
- Pseudonodosaria inflata* (Costa) = *Glandulina inflata* Costa, 1853, Foraminifer: Napoli, Italia, p. 14, pl. 4, fig. 1.
- Pullenia miocenica* Kleinpell, 1938, Miocene stratigraphy of California: Tulsa, Okla., Am. Assoc. Petroleum Geologists, p. 338, pl. 14, fig. 6.
- Pullenia multilobata* Chapman, 1900, California Acad. Sci. Proc., Geol., v. 1, p. 253, pl. 30, figs. 7, 7a.
- Pullenia quinqueloba* (Reuss) = *Nonionina quinqueloba* Reuss, Deutsch. Geol. Gesell. Zeitschr., v. 3, 1851, p. 71, pl. 5, fig. 31.
- Pullenia salisburyi* Stewart and Stewart, 1930, Jour. Paleontology, v. 4, no. 1, p. 72, pl. 8, fig. 2.
- Quinqueloculina imperialis* Hanna and Hanna, 1924, Washington Univ. Pub. Geology, v. 1, p. 58, pl. 13.
- Quinqueloculina minuta* Beck, 1943, Jour. Paleontology, v. 17, no. 6, p. 593, pl. 99, figs. 5-7.
- Reophax pilulifera* Brady, 1884, Repts. Sci. Results Explor. Voyage H.M.S. Challenger, Zoology, v. 9, p. 292, pl. 30, figs. 18-20.
- Rhabdammina eocenica* Cushman and Hanna, 1927, California Acad. Sci. Proc., ser. 4, v. 16, no. 8, p. 209, pl. 13, fig. 23, p. 57, pl. 8, fig. 11.
- Saracenaria schencki* Cushman and Hobson, 1935, Cushman Lab. Foram. Research Contr., v. 11, pt. 3, p. 57, pl. 8, fig. 11.
- Sigmomorphina schencki* Cushman and Ozawa, 1930, U.S. Natl. Mus. Proc., v. 77, no. 2829, art. 6, p. 133, pl. 35, figs. 6a, b.
- Siphogenerina hughesi* Cushman, 1925, Cushman Lab. Foram. Research Contr., v. 1, pt. 2, p. 36, pl. 7, figs. 4a-b.
- Siphogenerina mayi* Cushman and Parker, 1931, Cushman Lab. Foram. Research Contr., v. 7, pt. 1, p. 10-11, pl. 2, figs. 7a-b.
- Siphogenerina multicostata* Cushman and Jarvis, 1929, Cushman Lab. Foram. Research Contr., v. 5, pt. 4, p. 95, pl. 13, fig. 38.
- Siphogenerina nodifera* Cushman and Kleinpell, 1934, Cushman Lab. Foram. Research Contr., v. 10, pt. 1, p. 13, pl. 2, figs. 15, 16.
- Siphogenerina transversa* Cushman = *Siphogenerina raphanus* var. *transversa* Cushman, 1918, U.S. Natl. Mus. Bull. 103, p. 64.
- Sphaeroidina bulloides* d'Orbigny, 1826, Ann. Sci. Nat. Paris, ser. 1, tome 7, p. 267.
- Spiroloculina texana* Cushman and Ellisor, 1944, Cushman Lab. Foram. Research Contr., v. 20, pt. 3, p. 51, pl. 8, figs. 14-15.—*Spiroloculina wilcoxensis* Cushman and Garrett of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 257, pl. 4, fig. 9.
- Spiroplectammina directa* (Cushman and Siegfus) = *Spiroplectoides directa* Cushman and Siegfus, 1939, Cushman Lab. Foram. Research Contr., v. 15, pt. 2, p. 26, pl. 6, figs. 7-8.
- Spiroplectammina richardi* Martin, 1943, Stanford Univ. Pub. Geol. Sci., v. 3, no. 3, p. 14, pl. 5, figs. 3a-b.
- Stichocassidulina* cf. *S. thalmani* Stone, Mallory, 1959, Lower Tertiary biostratigraphy of the California Coast Ranges: Tulsa, Okla., Am. Assoc. Petroleum Geologists, p. 226, pl. 19, figs. 9a-c.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 283, pl. 20, figs. 8a, b.
- Stilostomella adolphina* (d'Orbigny) = *Dentalina adolphina* d'Orbigny, 1846, Foraminifères fossiles du bassin tertiaire de Vienne: Paris, France, Gide et Cie., p. 51.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 272, pl. 13, figs. 17a, b.
- Stilostomella advena* (Cushman and Laiming) = *Nodogenerina advena* Cushman and Laiming, 1931, Jour. Paleontology, v. 5, no. 2, p. 106, pl. 11, figs. 19a-b.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 272, pl. 13, figs. 20, 21.
- Stilostomella lepidula* (Schwager) = *Nodosaria lepidula* Schwager, 1866, Novara Exped. 1857-1859, Wien, Österreich., Geol. Theil, Bd. 2, Abt. 2, p. 210, pl. 15, figs. 27-28.
- Stilostomella sanctaerucis* Kleinpell, 1938, Miocene stratigraphy of California: Tulsa, Okla., Am. Assoc. Petroleum Geologists, p. 246, pl. 4, fig. 22.
- Stilostomella wegmanni* (Cole) = *Nodogenerina wegmanni* (Cole), Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 273, pl. 13, figs. 22, 23.
- Textularia shivelyi* Kleinpell, 1938, Miocene Stratigraphy of California: Tulsa Okla., Am. Assoc. Petroleum Geologists, p. 190, pl. 1, figs. 5, 9.
- Textularia* sp. of Fairchild and others, 1969, California Univ. Pubs. Geol. Sci., v. 81, p. 32, pl. 2, fig. 1.
- Trifarina hannai* (Beck) = *Angulogerina hannai* Beck, 1943, Jour. Paleontology, v. 17, no. 6, p. 607-608, pl. 108, figs. 26, 28.
- Trifarina wilcoxensis* (Cushman and Ponton) = *Virgulina wilcoxensis* Cushman and Ponton, 1932, Cushman Lab. Foram. Research Contr., v. 8, pt. 3, p. 67, pl. 8, fig. 22.
- Uvigerina auferiana* d'Orbigny, 1839, in de la Sagra, Ramon, Histoire physique et naturelle de l'île de Cuba: Paris, A. Bertrand, p. 106 (plates published separately).
- Uvigerina churchi* Cushman and Siegfus, 1939, Cushman Lab. Foram. Research Contr., v. 15, no. 2, p. 29, pl. 6, fig. 16.
- Uvigerina cocoaensis* Cushman, 1925, Cushman Lab. Foram. Research Contr., v. 2, pt. 2, p. 68, pl. 10, fig. 12.
- Uvigerina elongata* Cole, 1927, Bulls. Am. Paleontology, v. 14, pt. 51, p. 26, pl. 4, figs. 2, 3.
- Uvigerina gallowayi* Cushman, 1929, Cushman Lab. Foram. Research Contr., v. 5, pt. 4, p. 94-95, pl. 13, figs. 33-34.—not *Uvigerina gallowayi* of Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 277, pl. 16, figs. 9a, b.
- Uvigerina garzaensis* Cushman and Siegfus, 1939, Cushman Lab. Foram. Research Contr., v. 15, pt. 2, p. 28, pl. 6, figs. 15a, b.
- Uvigerina hispidocostata* Cushman and Todd, 1945, Cushman Lab. Foram. Research Spec. Pub. no. 15, p. 51, pl. 7, figs. 27, 31.
- Uvigerinella obesa impolita* Cushman and Laiming, 1931, Jour. Paleontology, v. 5, no. 2, p. 111, pl. 12, figs. 11a-b.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 276, pl. 16, figs. 10a, b.
- Uvigerinella sparsicostata* Cushman and Laiming, 1931, Jour. Paleontology, v. 5, no. 2, p. 112, pl. 12, figs. 2a-b.
- Vaginulinopsis saundersi* (Hanna and Hanna) = *Cristellaria saundersi* Hanna and Hanna, 1924, Washington Univ. Pub. Geology, v. 1, no. 4, p. 61, pl. 13, figs. 5, 6.
- Valvulineria araucana* (d'Orbigny) = *Rosalina araucana* d'Orbigny, 1839, Voyage dans l'Amerique meridionale, v. 5, pt. 5, p. 44.
- Valvulineria californica* Cushman, 1926, Cushman Lab. Foram. Research Contr., v. 2, no. 3, p. 60, pl. 9, figs. 1a-c.
- Valvulineria tumeyensis* Cushman and Simonson, 1944, Jour. Paleontology, v. 18, p. 201, pl. 33, figs. 13-14.—Sullivan, 1962, California Univ. Pubs. Geol. Sci., v. 37, no. 4, p. 280, pl. 17, figs. 3a, b, c.
- Verneuilina* sp. of Fairchild and others, 1969, California Univ. Pubs. Geol. Sci., v. 81, p. 33, pl. 3, figs. 9a, b.

REFERENCES CITED

- Aubert, Jane, and Berggren, W. A., 1976, Paleocene benthic foraminiferal biostratigraphy and paleoecology of Tunisia: Bulletin Centre Recherche, Pau-SNPA, France, v. 10, no. 2, p. 279-469, 12 pls.
- Bailey, E. H., and Everhart, D. L., 1964, Geology and quicksilver deposits of the New Almaden District, Santa Clara County, California: U.S. Geological Survey Professional Paper 360, 206 p.
- Bandy, O. L., 1972, Late Paleogene-Neogene planktonic biostratigraphy and its geologic implications, California: Society of Economic Paleontologists and Mineralogists, 47th Meeting, Pacific Section, Bakersfield, Calif., Mar. 9-10, Proceeding, p. 37-51.
- Bandy, O. L. 1972, Upper Paleogene planktonic correlations, Santa Barbara area, California in Guidebook to central Santa Ynez Mountains: American Association of Petroleum Geologists, Pacific Section, p. 55-58.
- Bandy, O. L., Morin, R. W., and Wright, R. C., 1969, Definition of the *Catpsydrax stainforthi* zone in the Sautesian Stage, California: Nature, v. 222, no. 5192, p. 468-469.
- Barker, R. W., and Grimsdale, T. F., 1936, A contribution to the phylogeny of Orbitoidal Foraminifera with a description of new forms from the Eocene of Mexico: Journal of Paleontology, v. 10, p. 231-247.
- Barron, J. A., 1976, Marine diatom and silicoflagellate biostratigraphy of the type Delmontian Stage and the type *Bolivina obliqua* Zone, California: U.S. Geological Survey Journal of Research, v. 4, no. 3, p. 339-351.
- Berger, W. H., 1972, Deep sea carbonates: Dissolution facies and age-depth constancy: Nature, v. 236, p. 392-395.
- Berggren, W. A., 1969, Cenozoic chronostratigraphy, planktonic foraminiferal zonation and the radiometric time scale: Nature, v. 224, p. 1072-1075.
- Berggren, W. A., 1972, A Cenozoic time scale—Some implications for regional geology and paleobiogeography: Lethaia, v. 5, no. 2, p. 195-215.
- Berggren, W. A., and Amdurer, M., 1973, Late Paleogene (Oligocene) and Neogene planktonic foraminiferal biostratigraphy of the Atlantic Ocean (Lat. 30° N. to Lat. 30° S.): Rivista Italiana di Paleontologia e Stratigrafia, v. 79, p. 337-391.
- Berggren, W. A., and Aubert, Jane, 1975, Paleocene benthonic foraminiferal biostratigraphy, paleobiogeography and paleoecology of Atlantic-Tethyan regions: Midway type fauna: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 18, p. 73-192, 19 pls.
- 1976, Eocene benthonic foraminiferal biostratigraphy and paleobathymetry of Orphan Knoll (Labrador Sea): Micropaleontology, v. 22, no. 3, p. 327-346.
- Berry, K. D., 1964, Tertiary micropaleontology, north side of Mt. Diablo, in Guidebook to the Mount Diablo field trip, 1964: Davis, Calif., Geological Society of Sacramento, p. 48-53.
- Berthiaume, S. A., 1938, Stratigraphy and foraminiferal fauna of the Meganos and Vacaville formations (Eocene) of California: Cornell University, Ithaca, N. Y., Ph.D. thesis.
- Blow, W. H., 1969, Late middle Eocene to Recent planktonic foraminiferal biostratigraphy: First International Conference on Planktonic Microfossils, Geneva 1967, Proceedings, p. 199-422.
- Blow, W. H., and Banner, F. T., 1962, The mid-Tertiary (upper Eocene-Aquitania) *Globigerinacea*, in Eames, F. E., and Banner, F. T., eds., Fundamentals of mid-Tertiary stratigraphical correlation: Cambridge, Cambridge University Press, p. 61-151.
- Boersma, A., 1974, Time-space distribution of *Uvigerina*: A Tertiary benthonic foraminiferal genus: Brown University, Providence, R. I., Ph.D. thesis, 252 p.
- Boersma, A., and Shackelton, N. J., 1977, Tertiary oxygen and carbon isotope stratigraphy, site 357 (mid-latitude South Atlantic): Initial Reports of the Deep Sea Drilling Project, v. 39, p. 911-924.
- Bolli, H. M., 1957, Planktonic Foraminifera from the Oligocene-Miocene Cipero and Lengua Formations of Trinidad, British West Indies: United States National Museum Bulletin, v. 215, p. 97-124.
- Brabb, E. E., Bukry, David, and Pierce, R. L., 1971, Eocene (Refugian) nannoplankton in the Church Creek Formation near Monterey, central California: U.S. Geological Survey Professional Paper 750-C, p. C44-C47.
- Brabb, E. E., Clark, J. C., and Throckmorton, C. K., 1977, Measured sections of Paleogene rocks from the California Coast Ranges: U.S. Geological Survey Open-File Report 77-714, 113 p.
- Brabb, E. E., Sonneman, H. S., and Switzer, J. R., Jr., 1971, Preliminary geologic map of the Mount Diablo-Byron area, Contra Costa, Alameda, and San Joaquin Counties, California: U.S. Geological Survey Open-File Map.
- Bramlette, M. N., and Riedel, W. R., 1954, Stratigraphic value of discoasters and some other microfossils related to recent coccolithophores: Journal of Paleontology, v. 28, p. 385-403.
- Bramlette, M. N., and Sullivan, F. R., 1961, Coccolithophorids and related nannoplankton of the Early Tertiary in California: Micropaleontology, v. 7, no. 2, p. 129-188.
- Bramlette, M. N., and Wilcoxon, J. A., 1967, Middle Tertiary calcareous nannoplankton of the Cipero Section, Trinidad, West Indies: Tulane Studies in Geology, v. 5, p. 93-131.
- Bronnimann, Paul, 1945, Zur Frage der verwandtschaftlichen Beziehungen zwischen *Discocyclina* ss. and *Asterocyclina*: Eclogae Geologicae Helvetiae, v. 38, p. 579-615.
- Bukry, David, 1971, Cenozoic calcareous nannofossils from the Pacific Ocean: San Diego Society of Natural History Transactions, v. 16, p. 313-327.
- 1973, Low-latitude coccolith biostratigraphic zonation: Initial Reports of the Deep Sea Drilling Project, v. 15, p. 685-703.
- 1974, Coccoliths as paleosalinity indicators—evidence from Black Sea: American Association of Petroleum Geologists Memoir 20, p. 353-363.
- 1975, Coccolith and silicoflagellate stratigraphy, north-western Pacific Ocean, Deep Sea Drilling Project Leg 32: Initial Reports of the Deep Sea Drilling Project, v. 32, p. 677-711.
- 1978, Biostratigraphy of Cenozoic marine sediment by calcareous nannofossils: Micropaleontology, v. 24, no. 1, p. 44-60.
- Bukry, D., Brabb, E. E., and Vedder, J. G., 1977, Correlation of Tertiary nannoplankton assemblages from the Coast and Peninsular Ranges of California: Memoir Segundo Congreso Latinoamericano de Geologia, v. 3, Venezuela Boletín de Geologia Publicacion Especial no. 7, p. 1461-1483.
- Bukry, David, and Bramlette, M. N., 1971, Validation of *Pedinocylus* and *Quinquerhabdus*, new calcareous nannoplankton genera: Tulane Studies in Geology and Paleontology, v. 8, p. 122.
- Bukry, David, and Percival, S. F., Jr., 1971, New Tertiary calcareous nannofossils: Tulane Studies in Geology and Paleontology, v. 8, p. 123-146.
- Butterlin, Jacques, 1966, Macroforaminifères de l'Éocène de Colombie: France Geological Society Bulletin, v. 7, no. 8, p. 627-629.
- Butterlin, Jacques, 1968, *Asterocyclina Cruzei*, espèce nouvelle de l'Éocène moyen de Colombie: Eclogae Geologicae Helvetiae, v. 61, no. 1, p. 325-326.

- , 1970, Macroforaminiferos y edad de la Formacion Punta Mosquito (Grupo Punta Carnero), de la isla de Margarita, Venezuela: *Asociacion Venezolana de Geologia, Minería y Petroleo Boletín Information*, v. 13, no. 10, p. 273-317.
- , 1971, Contribution à la connaissance du Paléogène marin du nord-ouest de la Colombie, basée sur les Macroforaminifères: *Eclogae Geologicae Helveticae*, v. 64, no. 1, p. 13-27.
- Bybell, L. M., 1975, Middle Eocene calcareous nannofossils at Little Stave Creek Alabama: *Tulane Studies in Geology and Paleontology*, v. 11, p. 177-252.
- Caudri, C. M. B., 1972, Systematics of the American Discocyclinas: *Eclogae Geologicae Helveticae*, v. 65, no. 1, p. 211-219.
- Church, C. C., 1931, Foraminifera of the Kreyenhagen shale: *Mining in California*, v. 27, no. 2, p. 202-213.
- Ciffelli, Richard, 1951, Eocene Foraminifera from the Point of Rocks area, California: Berkeley, University of California, M.A. thesis, 95 p.
- Clark, B. L., 1930, Tectonics of the Coast Ranges of middle California: *Geological Society of America Bulletin*, v. 41, p. 747-828.
- Clark, J. C., 1966, Tertiary stratigraphy of the Felton-Santa Cruz area, Santa Cruz Mountains, California: Stanford University, Stanford, Calif., Ph.D. thesis, 179 p.
- Colburn, I. P., 1961, The tectonic history of Mt. Diablo, California: Stanford University, Stanford, Calif., Ph.D. thesis, 276 p.
- Cole, W. S., 1927, A foraminiferal fauna from the Guayabal formation in Mexico: *Bulletin of American Paleontology*, v. 14, no. 51, 46 p.
- , 1948, Family Discocyclinidae, in J. A. Cushman, Foraminifera, their classification and economic use: Harvard University Press, 4th edit., 1950, p. 367.
- , 1958, Names of and variation in certain American larger Foraminifera particularly the camerinids: *American Paleontology Bulletin*, v. 38, no. 173, p. 261-284.
- Cole, W. S., and Applin, E. R., 1964, Problems of the geographic and stratigraphic distribution of American middle Eocene larger Foraminifera: *Bulletin of American Paleontology*, v. 47, no. 212, 48 p.
- Cole, W. S., and Bermudez, P. J., 1944, New Foraminiferal genera from the Cuban Middle Eocene: *American Paleontology Bulletin*, v. 28, no. 13, p. 333-350.
- Cole, W. S., and Gravell, D. W., 1952, Middle Eocene Foraminifera from Penon Seep, Matanzas Province, Cuba: *Journal of Paleontology*, v. 26, no. 5, p. 708-727.
- Cook, T. D., 1950, Eocene Foraminifera from the Devils Den area, California: California University, Berkeley, M.A. thesis, 116 p.
- Cushman J. A., 1977, Orbitoid Foraminifera of the genus *Orthophragmina* from Georgia and Florida: U.S. Geological Survey professional Paper 108, p. 115-118.
- , 1919, Fossil foraminifera from the West Indies: Carnegie Institution of Washington Publication No. 291, 217 p.
- , 1920, The American species of *Orthophragmina* and *Lepidocyclina*: U.S. Geological Survey Professional Paper 125-D, p. 39-108.
- Cushman, J. A., and Bermudez, P. J., 1936, New genera and species of foraminifera from the Eocene of Cuba: Contributions from the Cushman Laboratory for Foraminiferal Research, v. 12, pt. 2, no. 169, p. 29, pl. 5, figs. 11-12.
- , 1937, Further new species of foraminifera from the Eocene of Cuba: Contributions from the Cushman Laboratory for Foraminiferal Research, v. 13, pt. 1, no. 180, p. 6, pl. 1, figs. 27-28.
- Deflandre, G., 1942, Possibilités morphogenétiques comparées du calcaire et de la silice, a propos d'un nouveau type de microfossil calcaire de structure complexe, *Lithostromatol perdurum* new genus, new species: *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences*, v. 214, p. 917-919.
- , 1952, Classe des Coccolithophorides, in Grasse, P. P. ed., *Traite de Zoologie*: v. 1, pt. 1, p. 466.
- , 1959, Sur les nannofossiles calcaires et leur systématique: *Revue de Micropaléontologie*, v. 2, p. 127-152.
- Dibblee, T. W., Jr., 1973a, Stratigraphy of the southern Coast Ranges near the San Andreas fault from Cholame to Maricopa, California: U.S. Geological Survey Professional Paper 764, 45 p.
- , 1973b, Preliminary geologic map of the Gilroy quadrangle, Santa Clara County, California: U.S. Geological Survey Open-File Report, scale 1:24,000.
- , 1973c, Preliminary geologic map of the Gilroy Hot Springs quadrangle, Santa Clara County, California: U.S. Geological Survey Open-File Report, scale 1:24,000.
- Dibblee, T. W., Jr., 1977, Sedimentology and diastrophism during Oligocene time relative to the San Andreas fault system: Short Course presented to San Joaquin Geological Society, Bakersfield, Calif., Feb. 1977, p. 99-108.
- Donnelly, A. T., 1976, The Refugian Stage of the California Tertiary: Foraminiferal zonation, geologic history and correlations with the Pacific Northwest: Santa Barbara, University of California, Ph.D. thesis, 316 p.
- Douglas, R. G., 1973, Benthonic foraminiferal biostratigraphy in the central North Pacific, Leg 17: Initial Reports of the Deep Sea Drilling Project, v. 17, p. 607-671.
- Foreman, H. P., 1968, Upper Maestrichtian Radiolaria of California: London, The Palaeontological Association, Special Papers in Palaeontology, no. 3, 82 p.
- Funnell, B. M., 1964, The Tertiary Period, in The Phanerozoic time-scale, a symposium: *Quarterly Journal of Geological Society of London* supp., v. 120, p. 179-191.
- Gartner, S., Jr., 1969, Correlation of Neogene planktonic foraminifer and calcareous nannofossil zones: Gulf Coast Association of Geological Societies Transactions, v. 19, p. 535-599.
- , 1971, Calcareous nannofossils from the JOIDES Blake Plateau cores and revision of Paleogene nannofossil zonation: *Tulane Studies in Geology and Paleontology*, v. 8, p. 101-121.
- Gartner, S., Jr., and Bukry, David, 1969, Tertiary holococcoliths: *Journal of Paleontology*, v. 43, p. 1213-1221.
- Gibson, J. M., 1976, Distribution of planktonic foraminifera and calcareous nannoplankton, Late Cretaceous and early Paleogene, Santa Ynez Mountains, California: *Journal of Foraminiferal Research*, v. 6, no. 2, p. 87-106.
- Gibson, J., and Steinbeck, P. L., 1972, Age and correlation of the Ulatisian and Narizian Stages, California; Reply: *Geological Society of America Bulletin*, v. 83, no. 7, p. 2225-2232.
- Goudkoff, P. R., 1945, Stratigraphic relations of Upper Cretaceous in Great Valley California: *American Association of Petroleum Geologists Bulletin*, v. 29, no. 7, p. 956-1007.
- Haeckel, E. H. P. A., 1984, Systematische Phylogenie der Protisten und Pflanzen: Berlin, Reimer, 400 p.
- Hall, C. A., Jr., Jones, D. L., and Brooks, S. A., 1959, Pigeon Point Formation of Late Cretaceous age, San Mateo County, California: *American Association of Petroleum Geologists Bulletin*, v. 43, no. 12, p. 2855-2859.
- Haq, U. Z. B., 1968, Studies on upper Eocene calcareous nannoplankton from northwest Germany: *Stockholm Contributions in Geology*, v. 18, no. 2, p. 13-74.
- , 1973, Evolutionary trends in the Cenozoic coccolithophore genus *Helicopontosphaera*: *Micropaleontology*, v. 19, p. 32-52.
- Hardenbol, J., and Berggren, W. A., 1982, A new paleogene numerical time scale: Tulsa, Okla., American Association of Petroleum Geologists Studies in Geology No. 6. [in press].
- Hay, W. W., Mohler, H. P., Roth, P. H., Schmidt, R. R., and

- Boudereaux, J. E., 1967, Calcareous nannoplankton zonation of the Cenozoic of the Gulf Coast and Caribbean-Antillean area and transoceanic correlation: Gulf Coast Association of Geological Societies Transactions, v. 17, p. 428-480.
- Hay, W. W., Mohler, H. P., and Wade, M. E., 1966, Calcareous nannofossils from Nal'chik (northwest Caucasus): *Eclogae Geologicae Helvetiae*, v. 59, p. 379-400.
- Hill, M. L., and Dibblee, T. W., Jr., 1953, San Andreas, Garlock, and Big Pine faults, California—A study of the character, history, and tectonic significance of their displacement: Geological Society of America Bulletin, v. 64, p. 443-458.
- Hillebrandt, A. von, 1962, Das Paleozan und seine Foraminiferen fauna im Becken von Reichenhall und Salzburg: Abhandlungen der Mathematisch-Naturwissenschaftliche Klasse, Bavarian Academy of Sciences, new series no. 108, 182 p.
- Hinds, N. E. A., 1952, Evolution of the California landscape: California Division of Mines Bulletin 158, 240 p.
- Hornaday, G. R., and Phillips, F. J., 1972, Paleogene correlations, Santa Barbara area, California; An alternative view, in Guidebook, central Santa Ynez Mountains, Santa Barbara County, California: American Association of Petroleum Geologists Pacific Section and Society of Economic Paleontologists and Mineralogists Spring Field Trip, p. 59-71.
- Howe, H. V., 1939, Cook Mountain Eocene Foraminifera: Louisiana Geological Survey Geological Bulletin 14, 122 p.
- Israelsky, M. C., 1951, Foraminifera of the Lodo Formation, central California; Part 1, Arenaceous Foraminifera: U.S. Geological Survey Professional Paper 240-A, 29 p.
- , 1955, Foraminifera of the Lodo Formation central California, Part 2, Calcareous Foraminifera (Miliolidae and Lagenidae): U.S. Geological Survey Professional Paper 240-B, p. 31-77.
- Jenkins, D. G., 1966, Planktonic foraminiferal zones and new taxa from the Danian to Lower Miocene of New Zealand: New Zealand Journal of Geology and Geophysics v. 8, p. 1088-1126.
- Jenkins, D. G., and Orr, W. N., 1972, Planktonic foraminiferal biostratigraphy of the eastern equatorial Pacific—DSDP Leg 9: Initial Reports of the Deep Sea Drilling Project, v. 9, p. 1059-1193.
- Kamptner, E., 1927, Beitrag zur Kenntnis adriatischer coccolithophoriden: Archiv für Protistenkunde, v. 58, p. 173-184.
- , 1948, Coccolithen aus dem Torton des Inneralpinen Wiener Beckens: Österreichische Akademie der Wissenschaften Sitzungsberichte, v. 157, p. 1-16.
- , 1954, Untersuchungen über Feinbau der Coccolithen: Archiv für Protistenkunde, v. 100, no. 1, 90 p.
- , 1963, Coccolithineen-Skelettreste aus Tiefseeablagerungen des Pazifischen Ozeans: Annal Natur und Museum Wien, v. 66, p. 139-204.
- Keenan, M. F., 1932, The Eocene Sierra Blanca limestone at the type locality in Santa Barbara County, California: San Diego Society of Natural History Transactions, v. 7, no. 8, p. 53-84.
- Kleinpell, R. M., 1938, Miocene stratigraphy of California: Tulsa, Okla., American Association of Petroleum Geologists, 450 p.
- Kleinpell, R. M., and Weaver, D. W., 1963, Oligocene biostratigraphy of the Santa Barbara embayment, California—Pt. 3, Systematic catalogue and register of localities: California University Publications in Geological Science, v. 43, p. 165-250.
- Klump, B., 1953, Beitrag zur Kenntnis der Mikrofossilien des Mittleren und Oberen Eozan: Palaeontographica, v. 103A, p. 337-406.
- Kupper, Klaus, 1955, Upper Cretaceous Foraminifera from the "Franciscan Series" New Almaden District, California: Cushman Foundation for Foraminiferal Research Contributions, v. 6, pt. 3, p. 112-118.
- Laiming, Boris, 1943, Eocene foraminiferal correlations in California: California Division of Mines Bulletin 118, p. 193-198.
- Lamb, J., 1964, The stratigraphic occurrence and relationships of some mid-Tertiary Uvigerinas and Siphogenerinas: Micropaleontology, v. 10, no. 1, p. 457-476.
- Laughton, A. S., Berggren, W. A., and others, 1972, Initial reports of the Deep Sea Drilling Project, v. 12: U.S. Government Printing Office, 1243 p.
- Levin, H. L., 1957, Micropaleontology of the Oldsmar limestone (Eocene) of Florida: Micropaleontology, v. 3, no. 2, p. 137-154.
- Lipps, J. H., 1964, Oligocene in California: Nature, v. 208, p. 885-886.
- , 1966, Cenozoic planktonic foraminifera. I. Wall structure, classification and phylogeny of genera. II. California mid-Cenozoic biostratigraphy and zoogeography: Los Angeles, University of California, Ph.D. thesis, 271 p.
- , 1967a, Planktonic foraminifera, intercontinental correlation and age of California mid-Cenozoic microfaunal stages: Journal of Paleontology, v. 41, no. 4, p. 994-999.
- , 1967b, Miocene calcareous plankton, Reliz Canyon, California, in Gabilan Range and adjacent San Andreas fault: American Association of Petroleum Geologists and Society of Economic Paleontologists and Mineralogists, Pacific Sections, Guidebook, p. 54-60.
- Lipps, J. H., and Kalisky, Maurice, 1972, California Oligo-Miocene calcareous nannoplankton biostratigraphy and paleoecology, in Stinmeyer, E. H., and Church, C. C., eds., The Pacific Coast Miocene biostratigraphic symposium: Society of Economic Paleontologists and Mineralogists, Pacific Section, Bakersfield, Calif., p. 239-254.
- Locker, S., 1967, Neue coccolithophoriden (Flagellata) aus dem Alttertiär Norddeutschlands: Geologie, v. 16, p. 361-365.
- Loeblich, A. R., Jr., 1958, Danian Stage of Paleocene in California: American Association of Petroleum Geologists Bulletin, v. 42, p. 2260-2261.
- Loeblich, A. R., Jr., and Tappan, Helen, 1963, Type fixation and validation of certain calcareous nannoplankton genera: Proceedings of the Biological Society of Washington, v. 76, p. 191-196.
- Lohmann, G. P., and Tjalsma, R. C., 1975, Geographic and bathymetric distribution of paleocene deep water benthic foraminifera from the western Atlantic Ocean (abs.), in Benthonics '75: Halifax, Dalhousie University, p. 27.
- Mallory, V. S., 1959, Lower Tertiary biostratigraphy of the California Coast Ranges: Tulsa, Okla., American Association of Petroleum Geologists, 416 p.
- , 1970, Lower Tertiary Foraminifera from the Media Agua Creek drainage area, Kern County, California: Thomas Burke Memorial Washington State Museum Research Report 2, 211 p.
- Martin, Lewis, 1964, Upper Cretaceous and Lower Tertiary Foraminifera from Fresno County, California: Jahrbuch der Geologische Bundesanstalt, V. 9, 128 p.
- Martin, L. T., 1943, Eocene Foraminifera from the type Lodo formation, Fresno County, California: Stanford, Calif., Stanford University Publications in the Geological Sciences, v. 3, no. 3, p. 91-125.
- Martini, E., 1959, *Pemma angulatum* and *Micrantholithus basquensis*, zwei neue Coccolithophoriden-Arten aus dem Eozan: Senckenbergiana Lethaea, v. 40, p. 415-419.
- , 1971, Standard Tertiary and Quaternary calcareous nannoplankton zonation: International Conference on Planktonic Microfossils, 2d, Rome 1970, Proceedings, p. 739-785.
- Martini, E., and Bramlette, M. N., 1963, Calcareous nannoplankton from the experimental Mohole drilling: Journal of Paleontology, v. 37, p. 845-856.

- McDougall, Kristin, 1975, The microfauna of the type section of the Keasey Formation of northwestern Oregon: American Association of Petroleum Geologists, Society of Economic Geologists, Pacific Sections, 1974 Paleogene Symposium and Selected Technical Papers, p. 343-359.
- McKeel, D. R., and Lipps, J. H., 1975, Eocene and Oligocene planktonic foraminifera from the central and southern Oregon Coast Range: *Journal of Foraminiferal Research*, v. 5, no. 4, p. 249-269.
- McKeel, D. R., and Morin, R. W., 1979, Stratigraphic significance of *Planorotalites indiscriminata* (Mallory, 1959): *Journal of Foraminiferal Research*, v. 9, no. 3, p. 246-249.
- McLaughlin, R. J., Simoni, T. R., Osburn, E. D., and Bauer, P. G., 1971, Preliminary map of the Loma Prieta-Mount Madonna area, Santa Clara and Santa Cruz Counties, California: U.S. Geological Survey Open-File Report, scale 1:24,000.
- Miller, C., 1970, Nannoplankton aus dem Mittel-Oligozan von Norddeutschland und Belgien [Nannoplankton from the middle Oligocene of northern Germany and Belgium]: *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen* 135, p. 82-101.
- Nelson, R. N., and Schenck, H. G., 1930, Eocene algae and stellate Orbitoids from the Santa Ynez Range, California [abs.]: *Geological Society of America Bulletin*, v. 42, p. 371.
- Nilsen, T. H., 1977, Introduction to late Mesozoic and Cenozoic sedimentation and tectonics in California: San Joaquin Geological Society Short Course, Bakersfield, Calif., Feb., 1977, p. 7-19.
- Nilsen, T. H., and Clarke, S. H., Jr., 1975, Sedimentation and tectonics in the early Tertiary continental borderland of central California: U.S. Geological Survey Professional Paper 925, 64 p.
- Osburn, E. D., 1975, Geology of the Sveadal area, southern Santa Cruz Mountains, California: San Jose State University, M.S. thesis, 156 p.
- Palmer, K. W., 1929, Orbitolites from the Eocene of Oregon: *Bulletins of American Paleontology*, v. 10, no. 40, p. 13, 14.
- , 1934, Some large fossil Foraminifera from Cuba: *Mémoires de la Société d'histoire naturelle de Cuba*, v. 8, no. 4, p. 235-254.
- Pierce, R. L., 1972, Reevaluation of the late Miocene biostratigraphy of California; summary of evidence: Society of Economic Mineralogists and Paleontologists, Pacific Section, Pacific Coast Miocene Biostratigraphic Symposium Proceedings, p. 334-340.
- Poore, R. Z., 1976, Microfossil correlation of California Lower Tertiary sections: A comparison: U.S. Geological Survey Professional Paper 743-F, p. F1-F8.
- Poore, R. Z., and Brabb, E. E., 1977, Eocene and Oligocene planktonic Foraminifera from the upper Butano Sandstone and type San Lorenzo Formation, Santa Cruz Mountains, California: *Journal of Foraminiferal Research*, v. 7, no. 4, p. 249-272.
- Rau, W. W., 1958, Stratigraphy and foraminiferal zonation in some new Tertiary rocks of southwestern Washington: U.S. Geological Survey Oil and Gas Investigation Chart OC-57.
- , 1966, Stratigraphy and Foraminifera of the Satsop River area, southern Olympic Peninsula, Washington: Washington Division of Mines and Geology Bulletin 53, p. 1-66.
- Ryan, W. B. F., Cita, M. B., Rawson, M. D., Burckle, L. H., and Saito, Tsunemasa, 1975, A paleomagnetic assignment of Neogene stage boundaries and the development of isochronous datum planes between the Mediterranean, the Pacific and Indian Oceans in order to investigate the response of the world ocean to the Mediterranean "Salinity crisis": *Rivista Italiana di Paleontologia e Stratigrafia*, v. 80, p. 631-688.
- Schenck, H. G., 1929, *Discocyclina* in California: *Transactions of the San Diego Society of Natural History*, v. 5, no. 14, p. 211-240.
- , 1933, New records of *Discocyclina* in the California Eocene: *Micropaleontology Bulletin*, v. 4, no. 2, p. 72-79.
- Schenck, H. G., and Childs, T. S., Jr., 1942, Significance of *Lepidocyclina* (*Lepidocyclina*) *californica*, new species, in the Vaqueros Formation (Tertiary), California: Stanford University Publications in the Geological Sciences, v. 3, no. 2, 59 p.
- Schenck, H. G., and Klempell, R. M., 1936, Refugian stage of Pacific Coast Tertiary: *American Association of Petroleum Geologists Bulletin*, v. 20, no. 2, p. 215-225.
- Schmidt, R. R., 1970, Planktonic Foraminifera from the Lower Tertiary of California: University of California, Los Angeles, Ph.D. thesis, 339 p.
- , 1975, Upper Paleocene-Middle Eocene planktonic biostratigraphy from the Great Valley of California and adjacent areas, and correlation to the West Coast Microfaunal Stages, in Paleogene symposium and related papers: American Association of Petroleum Geologists, Society of Economic Paleontologists and Mineralogists, and Society of Economic Geologists, Pacific Section annual meeting, p. 439-455.
- Schoellhamer, J. E., and Kinney, D. M., 1953, Geology of portions of Tumey and Panoche Hills, Fresno County, California: U.S. Geological Survey Oil and Gas Investigations Map OM-128.
- Schwarz, E. H. L., 1894, Coccoliths: *Annals and Magazine of Natural History*, ser. 6, no. 14, p. 341-346.
- Sclater, J. G., Anderson, R. N., and Bell, N. L., 1971, Elevation of ridges and evolution of the central eastern Pacific: *Journal of Geophysical Research*, v. 76, p. 7888-7915.
- Smith, B. Y., 1957, Lower Tertiary Foraminifera from Contra Costa County, California: California University Publications in Geological Science, v. 32, p. 127-242.
- Smith, J. T., 1975, Age, correlation, and possible Tethyan affinities of mollusks from the Lodo Formation of Fresno County, California: society of Economic Paleontologists and Mineralogists, Pacific Section, Annual Convention, Long Beach, Proceedings, p. 464-483.
- Smith, Michael, 1974 Lower Tertiary planktonic biostratigraphy, California aqueduct section, northwestern Kern County, California: Fresno Calif., Fresno State College, M.A. thesis, 43 p.
- Steineck, P. L., and Gibson, J. M., 1971, Age and correlation of the Eocene Ulatian and Narizian stages, California: *Geological Society of America Bulletin*, v. 82, no. 2, p. 477-480.
- Steinforth, R. M., Lamb, J. L., Luterbacher, H., Beard, J. H., and Jeffords, R. M., 1975, Cenozoic planktonic foraminiferal zonation and characteristics of index forms: University of Kansas Paleontological Contributions, Art. 62, 425 p.
- Stinemeyer, E. H., 1976, The biostratigraphy of the Paleocene-Eocene Lodo Formation in the vicinity of Salt Creek, Fresno County, California: Geological guidebook for the 1974 fall field trip of the Pacific Section, Society of Economic Paleontologists and Mineralogists, p. 69-84.
- Stradner, H., and Edwards, A. R., 1968, Electron Microscopic studies on upper Eocene coccoliths from the Oamaru Diatomite, New Zealand: *Geologische Bundesanstalt Jahrbuch*, v. 13, p. 1-66.
- Sullivan, F. R., 1962, Foraminifera from the type section of the San Lorenzo Formation, Santa Cruz County, California: California University Publications in Geological Science, v. 37, no. 4, p. 233-335.
- , 1964, Lower Tertiary nannoplankton from the California Coast Ranges—part I, Paleocene: California University Publications in Geological Science, v. 44, no. 3, p. 163-228.

- 1965, Lower Tertiary nannoplankton from the California Coast Ranges—part II, Eocene: California University Publications in Geological Science, v. 53, p. 1-75.
- Tan, S. H., 1927, Discoasteridae incertae sedis: Koninklijk Nederlandsch Akademie van Wetenschappen Proceedings, v. 30, no. 3, p. 411-419.
- Tipton, Ann, 1980, Foraminiferal zonation of the Refugian Stage, latest Eocene of California, in Sliter, W. V., ed., Studies in marine micropaleontology and paleoecology: A memorial volume to Orville L. Bandy: Cushman Foundation for Foraminiferal Research Special Publication 19, p. 258-277.
- Tipton, Ann, Kleinpell, R. M., and Weaver, D. W., 1973, Oligocene biostratigraphy, San Joaquin Valley, California: California University Publications in Geological Science, v. 105, 81 p.
- Tjalsma, R. C., and Lohmann, G. P., 1975, Taxonomy and diversity of Paleocene deep water benthonic foraminifera from the West Atlantic [abs.], in Benthonics '75: Halifax, Dalhousie University, p. 48.
- Vail, P., Mitchum, R. M., Jr., and Thompson, S., II, 1977, Seismic stratigraphy and global changes of sea level, part 4: Global cycles of relative changes of sea level, in Payton C. E., ed., Seismic stratigraphy—applications to hydrocarbon exploration. American Association of Petroleum Geologists Memoir 26, p. 83-97.
- Vaughan, T. W., and Cole, W. S., 1941, Preliminary report on the Cretaceous and Tertiary larger Foraminifera of Trinidad, British West Indies: Geological Society of America Special Paper 30, 130 p.
- Warren, A. D., 1977, Nannoplankton biostratigraphy of the Devils Den Aqueduct and type Lodo Sections, western San Joaquin Valley, California. International Committee Paleogeographic Stratigraphical Field Conference Paleogene of California and Mexico, 13 p.
- 1980, Calcareous nannoplankton biostratigraphy of Cenozoic marine stages in California: American Association of Petroleum Geologists Studies in Geology No. 11, p. 60-69.
- Warren, A. D., and Newell, J. H., 1976, Nannoplankton biostratigraphy of the Upper Sacate and Gaviota Formations, Arroyo el Bulito, Santa Barbara County, California [abs.]: Pacific Sections, American Association of Petroleum Geologists, Society of Economic Geologists, Society of Economic Paleontologists and Mineralogists Annual Meeting, San Francisco, 1976, Program, p. 19-20.
- Warren, A. D., and Newell, S. H., 1980, Plankton biostratigraphy of the Refugian and adjoining stages of the Pacific Coast Tertiary, in Sliter, W. V., ed., Studies in marine micropaleontology and paleoecology: A memorial volume to Orville L. Bandy: Cushman Foundation for Foraminiferal Research Special Publication 19, p. 233-251.
- Watson, E. A., 1941, Stratigraphic occurrence of *Discocyclina* in the Eocene of California: Stanford University, Stanford, Calif., M.A. thesis, 65 p.
- Weaver, C. E., and others, 1944, Correlation of the marine Cenozoic formations of western North America: Geological Society of America Bulletin, v. 55, no. 5, p. 569-598.
- White, R. T., 1938, Eocene Lodo Formation and Cerros Member of California: Geological Society of America Proceedings, 1937, p. 256-257.
- 1940, Eocene Yokut sandstone north of Coalinga, California: American Association of Petroleum Geologists Bulletin, v. 24, p. 1722-1751.
- Wise, S. W. Jr. and Constans, R. E., 1975, Mid-Eocene planktonic correlations: Northern Italy-Jamaica, West Indies, Gulf Coast Association of Geological Societies Transactions, v. 26, p. 144-155.
- Woodring, W. P., 1930, Upper Eocene orbitoid Foraminifera from the western Santa Ynez Range, California, and their stratigraphic significance: San Diego Society of Natural History Transactions, v. 6, no. 4, p. 145-170.

INDEX

[Italic numbers indicate principal reference]

A

abbreviata, *Tarritilina*, pl. 2
tubulifera, *Angulogerina*, 17
abisectus, *Cyclicargolithus*, 51, 53, 60
acarinata, *Acarinina*, 19, 42
Acarinina acarinata, 19, 42
coalingensis, 19, 42
densa, 7, 12
mckannai, 19
soldadoensis, 42
wilcoxensis, 19
Acknowledgments, 1, 5, 22, 41, 49, 62
Actinocyclus aster, 47
Actinocythereis, 12
acuta, *Pleurostomella*, 11, 16
acuticosta, *Lagena*, 11, 71, 75; pl. 11
acutus, *Anomalinoidea*, 10, 16, 19; pl. 5
adamus, *Zygodiscus*, 31
adelaidana, *Bolivina marginata*, 71, 72
adolphina, *Dentalina*, 77
Nodogenerina, 17
Stilostomella, 64, 71, 77
adunca, *Glandulina*, 76
Marginulina, 11, 65, 71, 76
advena, *Bolivina*, 71, 74
Nodogenerina, 77
Nodosarella, 16; pl. 4
Stilostomella, 64, 69, 71, 77; pl. 13
californica, *Trifarina*, 28; pl. 3
aequa, *Morozovella*, 19
affinis, *Nodosaria*, 11
Age, foraminifers, large, 44
alabamensis, *Uvigerina*, 11
Alabamina wilcoxensis californica, 16, 22
sp., 10
alatalimbatus, *Lenticulina*, 10
alazaensis, *Marginulina*, 65, 69, 76
cubensis, *Pleurostomella*, 17
alleni, *Cibicidoides*, 7, 17
alligata, *Bulimina inflata*, 71, 72, 74; pl. 14
allomorphinoides, *Valvulineria*, 11
Almgren, A. A., cited, 42
alternans, *Pleurostomella*, 64, 68, 77; pl. 16
altus, *Chiasmolithus*, 53, 56, 60
Micrantholithus, 56; pl. 9
americana, *Truncatulina*, 74
americanus, *Cibicides*, 71, 74
crassiseptus, *Cibicides*, 71, 75; pl. 15
Ammobaculites cubensis, 17
sp., 17
Ammodiscus incertus, 69, 71, 74
pennyi, 10, 16; pl. 1
sp., 7
Amphimorphina becki, 64, 68, 74; pl. 13
californica, 16
ignota, 11, 16; pl. 2
jenkinsi, 68
jenkinsi Zone, 53, 57, 59, 67, 68, 69, 73
Amphipyndax plousios, 28
Amphistegina, 46
lopeztrigoi, 46
parvula, 42, 46; pl. 6
sp., 42
Amphisteginidae, 46
ampliapertura, *Globigerina*, 52
anarrhogus, *Sphenolithus*, 31
angiporoides, *Globigerina angiporoides*, 52, 59
angiporoides, *Globigerina*, 52, 59
minima, *Globigerina*, 52, 59
angulatum, *Pemma*, 38, 56, 61

Angulogerina abbreviata tubulifera, 17
cuneata, 10
hannai, 77
wilcoxensis, 10, 17, 28; pl. 3
angusta, *Pullenia quinqueloba*, 17
angustumilicata, *Globigerina*, 51
Año Nuevo area, 51
Año Nuevo section, 49, 51, 59, 60, 71, 74
faunal assemblages, 74
foraminifers, benthic, 71
plankton, calcareous, 51
Anomalina californiensis, 65, 68, 69, 70, 71, 74; pl. 17
crassisepta, 12, 22
dorri, 28
aragonensis, 22
garzaensis, 17, 64, 68, 74; pl. 17
salinasensis, 71, 74; pl. 17
umbonata, 64
Anomalinoidea, 7
acutus, 10, 16, 19; pl. 5
aragonensis, 10, 12, 16; pl. 5
crassiseptus, 11, 13, 17; pl. 5
danica rubiginosa, 7
garzaensis, pl. 5
judas, 10
keeni, 10, 17; pl. 5
regina, 16
rubiginosus, 19
welleri, 10, 16, 19
ansata, *Nonionella*, 17
antillea, *Lepidocyclus*, 47
antipoda, *Lenticulina*, 17
antipodium, *Lenticulina*, 11
applinae, *Loxostomoides*, 11; pl. 3
aproximata, *Dentalina*, 11
aragonensis, *Anomalina dorri*, 22
Anomalinoidea, 10, 12, 16; pl. 5
Aragonia, 10, 16; pl. 2
Globorotalia, 28
Morozovella, 19
Aragonia aragonensis, 11, 16; pl. 2
araucana, *Rosalina*, 77
Valvulineria, 71, 77; pl. 15
arenasensis, *Karrerella*, 16
Argilloecia, 12
arundinea, *Nodosaria*, 10, 16
aspera, *Clavulinoides*, 16
asperuliformis, *Vaginulinopsis*, 11, 22; pl. 2
Astacolus sp., 71
aster, *Actinocyclus*, 47
Asterocyclus, 42, 47; pl. 6
Asterigerina crassaformis, 11, 12, 13, 17, 22, 64, 74
Asterocyclus, 47
aster, 42, 47; pl. 6
distribution, 42
penonensis, 47
astroporus, *Markalius*, 31
asymbatos, *Stichomitra*, 28
atlantisae hispidula, *Ellipsonodosaria*, 64, 68, 76
hispidula, *Nodosarella*, 76; pl. 16
aubieriana, *Uvigerina*, 65, 69, 70, 71, 77; pl. 14
Aubert, Jane, cited, 28, 41
Avenal Sandstone Member, Kreyenhagen Formation,
Devils Den Aqueduct section, 22
Kreyenhagen Formation, Devils Den aqueduct
section, age, 29
Devils Den aqueduct section, Penutian Stage, 28
foraminifers, large, 41
Media Agua Creek area, 32
age, 33
worm tubes, 41

B

Baggatella californica, 17; pl. 4
Baggina robusta, 71, 72, 74; pl. 15
barbadiensis, *Discoaster*, 30, 38, 53, 56, 60; pl. 9
barbati, *Lenticulina*, 71, 75; pl. 11
Robulus, 75
basiplanata, *Dentalina*, 10, 17
basquensis, *Micrantholithus*, 56
Pemma, 61
Bathysiphon, 7, 12, 13, 21
eoecenic, 64, 69, 74
eoecenicus, 11, 13, 16, 19
sp. B Israelsky, 10
spp., 64, 71
beatus, *Cibicidoides*, 10, 11, 16; pl. 5
beccariiiformis, *Gavelinella*, 19
becki, *Amphimorphina*, 64, 68, 74; pl. 13
Lagena, 65, 75
belemnos, *Sphenolithus*, 51, 61
Benthic, defined, 1
Benthic foraminifers, 1, 5
age assignments, 36
Año Nuevo section, 62
central Coast Ranges, California, 22
Devils Den aqueduct section, 22
Eocene zonations, California, 41, 61
Lodo Gulch area, 22
Media Agua Creek area, 22
San Joaquin Valley, 22
San Lorenzo River section, 62, 63, 64, 65
Santa Cruz Mountains, 62
area, California, 61
Zayante Creek section, 62, 69
Benthonic, defined, 1
Benthonic foraminiferal zones, 1, 22
Año Nuevo section, 62, 71
central ranges of California, 4
Devils Den aqueduct area, 13, 22, 44
Locatelli Formation, 19, 21
Lodo Formation, 29
Lodo Gulch area, 22, 29
Mallory, 5, 22, 28, 33
Media Agua Creek, 22
Paleogene of California, summary, 36
Rices Mudstone Member, 68
San Joaquin Valley, 22, 33
San Lorenzo Formation, 67, 68
San Lorenzo River section, 62, 63, 73
Santa Cruz Mountains, 61
Twobar Shale Member, 67, 68
Vaqueros Formation, 71
Zayante Creek section, 62, 70
Berggren, W. A., cited, 28, 41, 42
bidens, *Chiasmolithus*, 30
Bifarina elegans, 68
elegans, 11, 64, 74; pls. 2, 15
nuttalli, 11, 16, 28; pl. 2
vicksburgensis, 10
bigelowi, *Braarudosphaera*, 56, 60
bijugatus, *Zygrhablithus*, 31, 38, 56, 61
binodosus, *Discoaster*, 30, 38
Biostratigraphy, foraminiferal, 7, 13
nannofossils, calcareous, 38
bisectus, *Coccolithus*, 38
Dictyococcites, 30, 51, 52, 53, 56; 60; pl. 9
Blow, W. H., cited, 49
boffalorae, *Nodosaria*, 76
Boldia hodgei, 64, 68, 74; pl. 17
stratigraphic range, 73
Bolivina advena, 71, 74

Bolvinia—Continued

- crenulata*, 19
explicata californica, 11
gracilis, 64, 74
huxeri, 16; pl. 3
kleinPELLI, 17, 64, 74; pl. 13
marginata, 65, 69, 70, 71, 74; pls. 10, 13
adelaidana, 71, 72
pisciformis, 17
scabrata, 64, 68, 74; pl. 13
 sp., 11, 64; pl. 13
 spp., 71
Bolivina *mexicanus*, 65, 74; pl. 13
boweri, *Subbotina*, 12
Braurudosphaera bigelowi, 56, 60
discula, 30
 Brabb, E. E., cited, 1, 5, 22, 28, 39, 49, 52, 58, 63, 71
 Bracheux Sandstone, Paris Basin, 7
bradburyi, *Bulimina*, 10, 16, 19; pl. 2
bramlettei, *Helicosphaera*, 56, 61
Virgulina, 75
Bramletteius serraculoides, 56, 60; pl. 9
bramletti, *Fursenkoina*, 65, 69, 71, 75; pl. 16
brevispira, *Turritilina*, 10, 16
 Bukry, David, cited, 1, 22, 28, 29, 33, 36, 38, 49, 58
Bulimina, 7
callahani, 10, 12, 13, 16; pl. 3
carneroensis, 71, 74; pls. 13, 14
 stratigraphic range, 72
consanguina, 17
corrugata, 11, 22, 64, 68, 71, 74; pl. 14
 Zone, 68, 73
curtissima, 10, 16, 64, 68, 74; pls. 3, 14
debilis, 11, 28
elegantissima, 74
excavata, 17, 28
impedens, 10, 17; pl. 3
inflata, 65, 70
alligata, 71, 72, 74; pl. 14
instabilis, 10, 16
macilenta, 10, 16; pl. 3
microcostata, 64, 68, 69, 74; pl. 14
orphanensis, 12, 13
ovata, 10, 77
pupoides, 16, 77
pyrula, 16
sculptilis lacinata, 65, 68, 74; pl. 14
semicostata lacrima, 17
trihedra, 17
trinitatis, 10, 12, 16; pl. 3
truncanella, 17
whitei, 11, 16; pl. 3
 sp., 10, 17
 spp., 71
Buliminella bradburyi, 10, 16, 19; pl. 2
curta, 69, 71, 74
elegantissima, 71, 74; pl. 13
grata, 12
convoluta, 10, 16, 22; pl. 2
subfusiformis, 69, 71, 74
 sp., 64
 Bulitian Stage, 28, 29, 33, 36
 age, 29, 33, 36
 benthonic foraminiferal zones, 33
 type section, 33
bulloides, *Sphaeroidina*, 69, 71, 77; pl. 13
 Burdigalian fauna, Europe, 1
 Butano Sandstone, Bear Creek area, 38
 Bear Creek area, age, 39
 coccoliths, 52
 foraminifers, benthic, 68
 benthic, agglutinated, 19, 21
 calcareous, 19, 21
 planktic, 19, 21
 Lompico area, 5, 19, 38
 age, 38
 nannofossils, 38
 paleobathymetry, 19, 21
 San Lorenzo River section, 63, 68
 Santa Cruz Mountains, 5, 13, 19, 21

C

- calathus*, *Chiphragmalithus*, 30
calcar, *Lenticulina*, 69, 71, 75; pl. 11
Nautilus, 75
 California, basement rocks, 4
 Eocene rocks, 4
 marine strata, 58
 Paleogene benthonic foraminiferal stages, 22
 Paleogene, summary, 36
 paleogeography, 4
 provincial stages, 5, 38
 tectonic evolution during Mesozoic and Cenozoic time, 4
 Tertiary, map, 4
 California Tertiary stages, 1, 5
 Año Nuevo section, 52, 59, 62, 72, 74
 Avenal Sandstone, 28
 benthonic foraminiferal stages, 1, 4, 22, 29, 36, 41, 62, 72
 Bulitian Stage, 28, 29, 33, 36
 Cheneyan Stage, 36, 38
 Church Creek area, 58
 Delmontian Stage, 1
 Devils Den aqueduct area, 22, 36, 44
 Kleinpell, 1, 72, 73
 Lambert Shale, 71
 Lodo Formation, type, 29, 32, 33, 36
 Lodo Gulch area, 29, 36
 Mallory, 1, 5, 22, 28, 29, 33
 Martinez Stage, 7
 Media Agua Creek area, 33, 36
 Mohnian Stage, 1
 nannoplankton zones, 1, 22, 28, 29, 36
 Narizian boundary, 73
 Narizian-Refugian boundary, 59
 Narizian Stage, 28, 33, 36, 42, 53, 59, 62, 67, 68, 69, 70, 72, 73
 Penutian Stage, 28, 29, 33, 36, 44
 planktonic foraminiferal zones, 1, 22, 36, 41, 59
 Refugian Stage, 53, 58, 59, 68, 69, 70, 73
 Refugian-Zemorrian boundary, 53, 69, 73
 Relizian Stage, 52, 62, 72, 74
 San Lorenzo River section, 53, 59, 62, 67, 68, 70, 73
 Santa Cruz Mountains, 62
 Saucian-Relizian boundary, 52
 Saucian Stage, 52, 59, 71, 72, 74
 Twobar Shale Member, 68
 Ulatisian-Narizian boundary, 29
 Ulatisian Stage, 29, 33, 36, 42, 67, 68
 Ynezian-Bulitian boundary, 29, 33
 Ynezian Stage, 28, 29, 33, 36
 Zayante Creek section, 59, 62, 70, 74
 Zemorrian boundary, 70, 71, 73
 Zemorrian Stage, 36, 38, 52, 53, 58, 59, 69, 70, 71, 72, 73
 Zemorrian-Saucian boundary, 52, 59, 72, 73, 74
californica, *Atabamina wilcoxensis*, 16, 22
Amphimorphina, 16
Baggatella, pl. 4
Bolvinia explicata, 11
Discocyclina, 46, 48
Globorotalia, 28
Plectofrondicularia, 71, 76
Siticosigmollina, 10, 17, 19; pl. 1
Trifarina advena, 10, 17, 28; pl. 3
Valvulineria, 71, 72, 77
californicus, *Chiasmolithus*, 30, 38
Clavulinoides, 10, 16; pl. 1
californiensis, *Anomalina*, 65, 68, 69, 70, 71, 74; pl. 17
Cassidulinoides, 65, 70, 74; pl. 13
Fursenkoina, 64, 75; pl. 16
Virgulina, 75
callahani, *Bulimina*, 10, 12, 13, 16; pl. 3
Camerina striatoreticulata, 45
Camerina willcoxi, 45
Campylosphaera dela, 30, 56, 60
eodela, 30, 51, 60
 Cañada de Santa Anita, 73
cancellata, *Cyclammina*, 71
obesa, *Cyclammina*, 75; pl. 10
 Canoas Siltstone Member, Kreyenhagen Shale, age, 36
 Kreyenhagen Shale, Garza Creek, 36
 Oil City, 36
carcoselleensis, *Globorotalia*, 59
carinatus, *Triquetrorhabdulus*, 51, 61
carneroensis, *Bulimina*, 71, 74; pls. 13, 14
carteri, *Helicosphaera*, 51, 61
Cassidulina crassipunctata, 65, 69, 70, 71, 74; pl. 16
diversa, 64, 74; pl. 16
globosa, 11, 16, 75
margareta, 71, 74
 sp., 71
 spp., 64
Cassidulinoides californiensis, 65, 70, 74; pl. 13
Catapsydrax unicavus, 51
Caucasina schencki, 22
cavus, *Coccolithus*, 30
 Cenozoic formations, western North America, correlation, 1
cerroazulensis, *Globorotalia*, 59
Globorotalia cerroazulensis, 52
cerroazulensis, *Globorotalia*, 52
charoides, *Glomospira*, 17, 19
chehalisensis, *Lenticulina*, 65, 75
Robulus, 75
 Cheneyan Stage, age, 36, 38
Chiasmolithus, 61
altus, 53, 56, 60
bidens, 30
californicus, 30, 38
consuetus, 30, 38
expansus, 30, 38, 56, 60
gigas, 30, 38
 Zone, 31
grandis, 30, 56, 60
solitus, 30, 56, 60
 sp., 51
childsi, *Valvulineria*, 10, 16
Chiloguembelina cubensis, 59; pl. 7
midwayensis subcylindrica, 28
 sp., 59
Chilostomella sp., 64, 71
Chilostomelloides cylindroides, 10
 sp., 17
Chiphragmalithus calathus, 30
chirans, *Robulus*, 69, 76
 Church Creek area, 49, 58
 plankton, calcareous, 58
 Santa Lucia Range, 49, 59
 Church Creek Formation, Church Creek area, 49, 58
 Church Creek area, age, 59
 coccoliths, 58
 foraminifers, planktic, 49, 58, 59
 Santa Lucia Range, 49, 58
 zone assignments, 59
churchi, *Uvigerina*, 64, 68, 77; pl. 14
Cibicides americanus, 71, 74
americanus crassiseptus, 71, 75; pl. 15
cocoaensis, 11
cushmani, 74
elmaensis, 65, 68, 73, 75
 stratigraphic range, 73
falconensis, 64, 75
felix, 11
floridanus, 69, 71, 75; pl. 15
grimsdalei, 17
haydoni, 65, 69, 75; pl. 15
 Subzone, Donnelly, lower Refugian, 68
hodgei, 74
kernensis, 11, 17, 28
laimingi, 22
martinezensis, 11, 16
pachyderma, 17, 64, 68, 75
praecursorius, 11
pseudoungerianus evolutus, 65, 69, 71, 75; pl. 16
spirospunctata, 68
spirospunctatus, 11, 22, 64, 75; pl. 16
 sp., of Mallory, 16
 sp. B, 17
 sp., 69

Cibicides—Continued

- spp., 64, 71
Cibicidina mauricensis, 17
Cibicidoides, 7
 alleni, 7, 17
 beatus, 10, 11, 16; pl. 5
 cushmani, 17
 eponidiformis, 10, 16
 fortunatus, 10, 13, 16; pl. 5
 grimsdalei, pl. 5
 martinezensis, 19
 pseudowuellerstorfi, 16
 spiropunctatus, 16
 whitei, 10, 16, 19; pl. 5
 Claiborne Group, Gulf Coast, 42
 Clark, J. C., cited, 1, 19, 22, 39, 49, 63, 71
 Clark, Joseph, cited, 5
clarki, *Cyclammina*, 17; pl. 1
 Discocyclina, 47
 Orthophragmina, 47
 Pseudophragmina, 42
 (*Proporocyclina*), 47; pl. 6
Clavulina patens, 76
Clavulinoides, 7
 aspera, 16
 californicus, 10, 16; pl. 1
 sp., 7
clypeiformis, *Lenticulina*, 69, 71, 76; pl. 11
coaledensis, *Lenticulina*, 65, 76
 Robulus, 76
coalingensis, *Acarinina*, 19, 42
 Gaudryina, 17; pl. 1
 jacksonensis, 10, 28
 Haplophragmoides, 10, 16
 Coast Ranges, Calif., 2
 foraminifers, large, Eocene, 41
 stratigraphy, 1
 Coccolith assemblages, Año Nuevo section, 51
 Church Creek area, 58
 northern California, 60
 San Lorenzo River section, 52, 53
 Zayante Creek section, 52
 zone assignments, 51, 59
 Coccolith, 49, 51
 distribution, 51
 sample preparation, 59
 sp., Deep Sea Drilling Project cores, 60
 Europe, 60
 Gulf Coast, 60
 illustration references, 60
 indeterminate, 61
 undifferentiated, 61
 zonation, 49
 Coccolith zones, Año Nuevo section, 51, 74
 Bukry, 51, 53, 56
 Church Creek area, 58
 Church Creek Formation, 58
 Lambert Shale, 52
 Rices Mudstone Member, 52, 53
 San Lorenzo Formation, 52, 53
 San Lorenzo River section, 53, 73
 Santa Cruz Mountains, 49
 Santa Lucia Range, 49
 Two-bar Shale Member, 53
 Vaqueros Formation, 51
 Vaqueros Sandstone, 52
 Zayante Creek section, 52, 74
Coccolithus, 61
 bisectus, 38
 cavus, 30
 crassus, 29, 30
 cribellum, 30, 56, 60
 copelagicus, 30, 38
 formosus, 53, 56, 60; pl. 9
 magnicrassus, 30
 pelagicus, 30, 38, 53
 staurion, 56, 60
 subdistichus, 30
 spp., 51, 56
cocoaensis, *Cibicides*, 11

cocoaensis—Continued

- Dentalina*, 65, 75
Nodosaria, 75
Uvigerina, 68, 69, 73, 77
 Code of Stratigraphic Nomenclature, 5, 39
 Colburn, I. P., cited, 42
 Cole, W. S., cited, 46
colei, *Tritaxilina*, 11, 13, 16; pl. 1
collactea, *Truncorotaloides*, 59
coloratus, *Lenticulina*, 71
columbiana, *Gümbelitra*, 59, 60; pl. 7
communis, *Dentalina*, 11
compacta, *Helicospaera*, 56, 61
concinus, *Heliorthus*, 38
condoni, *Eponides*, 75
 Gyroidina, 64, 75
conicus, *Sphenolithus*, 51, 61
Conotrochammina spp., 17
consanguinea, *Bulimina*, 17
constricta, *Nodosarella*, 16
consuetus, *Chiasmolithus*, 30, 38
contortus, *Trochamminoides*, 16
convergens, *Lenticulina*, 17
convoluta, *Buliminella grata*, 22; pl. 2
cooperensis, *Dentalina*, 71, 75; pl. 11
 Corals, 1
 Correlation chart, standard, based on benthic foraminifers, 1
 standard, based on mollusks, echinoids, and corals, 1
 for marine Cenozoic formations of western North America, 1
 Correlation, summary, faunal assemblages, 72
corrugata, *Bulimina*, 11, 22, 64, 68, 71, 74; pl. 14
costata, *Entosolenia*, 75
 Lagena, 65, 71, 75
costifera, *Nonionella*, 69, 71, 76
 Nonionina, 76
crassaformis, *Asterigerina*, 11, 12, 13, 17, 22, 64, 74
crassipunctata, *Cassidulina*, 65, 69, 70, 71, 74; pl. 16
crassisepta, *Anomalina*, 12, 22
crassiseptus, *Anomalinoidea*, 11, 13, 17; pl. 5
 Cibicides americanus, 71, 75; pl. 15
crassus, *Coccolithus*, 29, 30
craticulus, *Toweius*, 31
crebra, *Rhabdosphaera*, 31
crenulata, *Bolivina*, 19
cribellum, *Coccolithus*, 30, 56, 60
Cribrastomoides sp., 17
cristata, *Nannotetrina*, 38
Cristellaria midwayensis, 76
 saundersi, 77
 simplex, 76
crowleri, *Lagena*, 11
cruciformis, *Discoaster*, 30
Cruciplacolithus staurion, 30
 tenuis, 56, 60
 Zone, 36
cubana, *Dorothia*, 64, 67, 75; pl. 10
 Gaudryina, 75
cubensis, *Ammobaculites*, 17
 Chiloguembelina, 59; pl. 7
 Dorothia, 17
 Gumbelina, 59
 Plectina, 13, 17
 Pleurostomella alazanensis, 17
cultratus, *Lenticulina*, 76
 Robulus, 76
cuneata, *Angulogerina*, 10
curta, *Buliminella*, 69, 71, 74
 Vulvulina, 10, 16
curtissima, *Bulimina*, 10, 16, 64, 68, 74; pls. 3, 14
curvatura, *Marginulina*, 10
cushmani, *Cibicides*, 74
 Cibicidoides, 17
Cyclammina, 7, 12, 19
 cancellata, 71
 obesa, 75; pl. 10
 clarki, 17; pl. 1
 incisa, 17, 69, 75; pl. 1

Cyclammina—Continued

- pacifica*, 65, 75
simiensis, 11, 13, 16, 19; pl. 1
 sp., 69
 spp., 65, 69, 71
Cyclicargolithus abisectus, 51, 53, 60
 floridanus, 30, 38, 51, 53, 56, 60
 luminus, 56, 60
 neogammation, 60
Cyclococcolithina formosa, 30
 gammation, 30, 56, 60
 kingi, 56, 60
 protoannula, 60
 pseudogammation, 30
Cyclococcolithus formosus, 38
 inversus, 38
 luminus, 60
Cyclolithella inflexa, 38
cylindroides, *Chilostomelloides*, 10
 Cymbaloporidae, 47
 Cytherella, 12

 D

Daktylethra punctulata, 56, 60; pl. 9
danica rubiginosa, *Anomalinoidea*, 7
danvillensis, *Pseudoparrella*, 64
debilis, *Bulimina*, 11, 28
 Deep Sea Drilling Project sites, 12
 Definition, Lodo Formation, 39
deflandrei, *Discoaster*, 30, 51, 53, 60; pl. 9
deflata, *Haplophragmoides*, 65, 75; pl. 10
dela, *Campylosphaera*, 30, 56, 60
delicatus, *Discoaster*, 38
deliciae, *Nodosaria*, 10, 16
 Delmontian Stage of Kleinpell, 1
densa, *Acarinina*, 7, 12
Dentalina adolphina, 77
 approximata, 11
 basiplanata, 10, 17
 cocoaensis, 65, 75
 communis, 11
 cooperensis, 71, 75; pl. 11
 jacksonensis, 11, 16, 65, 75
 multilineata, 11
 quadralata, 69, 71, 72, 75; pl. 11
 soluta, 65
 spinosa, 10, 16
 ornatior, 64, 75; pl. 11
 substrigata, 10
 wilcoensis, 11
 sp., 69, 71; pl. 11
 spp., 10
 Devils Den, Calif., 2, 4
 Devils Den area, 4, 41, 44
 aqueduct section, 13, 21, 22
 age, 36
 Bulitian Stage, 28
 conclusions, 29
 foraminiferal biostratigraphy, 13
 foraminifers, benthic, 7
 benthic, stages, 22, 28, 29
 planktic, 21
 nannoplankton, 28
 Narizian Stage, 22, 28
 paleobathymetry, 13
 Penutian Stage, 28
 rock units, 22
 Ulatian Stage, 28
 Ynezian Stage, 28
 comparison, 33
 foraminifers, large, 41, 44
 Kern County, Calif., 5, 13, 22
 locality descriptions, 48
 stratigraphy, 41
 diastypus, *Discoaster*, 30
 Dibblee, T. W., Jr., cited, 4, 33, 36, 41
dibollensis, *Fursenkoina*, 64, 75; pl. 16
Virgulina, 75

dictyoda, *Reticulofenestra*, 31, 38
Dictyococites bisectus, 30, 51, 52, 53, 56, 60; pl. 9
hesslandii, 38
scrippsae, 30, 51, 53, 56, 60
Dictyomitra multicosata, 28
directa, *Plectofrondicularia miocenica*, 71, 76; pl. 12
Spiroplectamina, 16, 65, 77; pls. 1, 10
Spiroplectoides, 77
Discoaster barbadiensis, 30, 38, 53, 56, 60; pl. 9
barbadiensis Zone, 31, 53, 56, 58, 73
/Helicopontosphaera reticulata Zone, 31
bifaz Subzone, 53, 56
binodosus, 30, 38
Zone, 31
cruciformis, 30
deflandrei, 30, 51, 53, 60; pl. 9
delicatus, 38
diastypus, 30
Zone, 28, 29, 31, 33
boundary, 29
distinctus, 30, 38
divaricatus, 51, 53, 61
elegans, 30, 56, 61
exilis, 51, 61
gemmifer, 30
germanicus, 30
helianthus, 30
lenticularis, 30
lodoensis, 30, 38, 51, 53, 56, 61
Zone, 13, 29, 31, 33, 44
mirus, 30, 38
mohleri Zone, 29, 32, 33
multiradiatus, 30
Zone, 29, 31, 33
nobilis, 30
Zone, 29, 33
nodifer, 30, 56, 61; pl. 9
nonaradiatus, 30
saipanensis, 30, 38, 53, 56, 61; pl. 9
Subzone, 53, 56
Zone-*Discoaster barbadiensis* Zone boundary, 73
strictus, 30
Zone, 31
sublodoensis, 29, 31, 33
Zone, 13, 29, 31, 33
tani, 31, 56, 61
ornatus, 31
tribrachiat, 61
variabilis, 51, 61
wemmelensis, 31
spp., 51
Discoasteroides kuepperi, 31
kuepperi Zone, 31
megastypus, 31
Discoasters, Paleocene, 38
Discocyclina californica, 46, 48
californica, holotype, 42
clarki, 47
sp., 42
(*Discocyclina*) *marginata*, 42, 46
(*Discocyclina*) *marginata*, *Discocyclina*, 42, 46
Discocyclinidae, 46
classification, 46
Discolithina, 61
distincta, 61
multiplora, 51, 61
plana, 30, 31, 51, 53, 61
spp., 53
Discolithus distinctus, 56
pulchroides, 61
spp., 56
discula, *Braarudosphaera*, 30
Discussion, Lodo Formation, 39
plankton, calcareous, 59
dissimilis, *Sphenolithus*, 51, 53, 61
distincta, *Discolithina*, 61
distinctus, *Discoaster*, 30, 38
Discolithus, 56
divaricatus, *Discoaster*, 51, 53, 61
diversa, *Cassidulina*, 64, 74; pl. 16

Domengine Formation, age, 36
foraminifers, large, 42
Mount Diablo area, age, 42
Oil City, 36
Domengine Sandstone, 7
Lodo Gulch, 39
age, 36
Donnelly, A. T., cited, 62, 68
dorfi, *Eponides*, 17
Dorothia, 7
cubana, 64, 67, 75; pl. 10
cubensis, 17
excentrica, 10, 11
germanica, 17; pl. 1
principiensis, 17
sp., 11
dorri, *Anomalina*, 28
aragonensis, *Anomalina*, 22
dubia, *Marginulina*, 69, 76
dubius, *Neochiastozygus*, 39
Neococcolithes, 31
Zygodolithus, 56, 61; pl. 9
E
Early Zemorrian-late Zemorrian boundary, 70, 71, 73
echinata, *Vaginulinopsis*, 16
Echinoids, 1
eggerella elongata, 64, 68, 69, 75; pl. 10
subconica, 64, 75; pl. 10
sp., 11
eggeri, *Haplophragmoides*, 13, 16
elegans, *Bifarina*, 68
Discoaster, 30, 56, 61
eleganta, *Bifarina*, 11, 64, 74; pls. 2, 15
Siphogenerina, 74
elegantissima, *Bulimina*, 74
Buliminella, 71, 74; pl. 13
Ellipsolithus macellus, 31
Ellipsonodosaria atlantisae hispidula, 64, 68, 76
elmaensis, *Cibicides*, 65, 68, 73, 75
elongata, *eggerella*, 64, 68, 69, 75; pl. 10
Hyperammina, 19
Karrerella, 11, 64, 68, 75; pl. 11
Uvigerina, 7, 11, 17, 64, 77; pls. 3, 14
Elphidium sp., 71
eminens, *Toweius*, 31
entaster, *Micrantholithus*, 31
Entosolenia costata, 75
squamosa hexagona, 75
eocaena, *Globigerina*, 59
Eocene foraminifers, large, Coast Ranges, Calif., 41
Eocene rocks, California, 4
Eocene stages of Mallory, 1
Eocene-Oligocene boundary, San Lorenzo River section, 63
eocenica, *Bathysiphon*, 64, 69, 74
Gonatosphaera, 16
Hoeglundina, 19
Martinottiella, 16, 64, 76; pls. 1, 11
Pullenia, 10, 16
Rhabdammina, 10, 13, 16, 19, 64, 77; pl. 10
eocenicus, *Bathysiphon*, 13, 19
Eoconuloides, 46
lopeztrigoi, 42, 46; pl. 6
parvulus, 46
wellsi, 42, 46; pl. 6
eodela, *Campylosphaera*, 30, 51, 60
Eofabiana, 47
grahami, 42, 47; pl. 6
eopelagicus, *Coccolithus*, 30, 38
epigona-lata, *Rzehakina*, 19
Epistomina partechiana, 10, 17
Epistominella spp., 71
Eponides condoni, 75
dorfi, 17
lodenensis, 11, 28
minima, 17
primus, 7, 11, 16
umbonatus, 76
sp., 10, 16, 17, 65

eponidiformis, *Cibicoides*, 10, 16
Ericsonia subpertusa, 38
euphratis, *Helicosphaera*, 51, 53, 61
Europe, Oligocene, 1
Eustatic sea-level change, identified by seismic stratigraphic methods, 21
evolutus, *Cibicides pseudoungerianus*, 65, 69, 71, 75; pl. 16
ewaldi, *Nodosaria*, 10, 16
excavata, *Bulimina*, 17, 28
excavatus, *Haplophragmoides*, 11, 13, 17, 19
excentrica, *Dorothia*, 10, 11
exilis, *Discoaster*, 51, 61
exima, *marginulina*, 65, 76
expansus, *Chiasmolithus*, 30, 38, 56, 60
explicata californica, *Bolivina*, 11
F
falconensis, *Cibicides*, 64, 75
fallax, *Karrerella*, 11
Fasciculolithus involutus, 31
mitreus, 31
schaubi, 31
tympaniformis, 31, 38
Zone, 38
Faunal assemblages, correlation, summary, 72
Faunas, Coast Range sections, 1
Coast Ranges, California, 2
Paleogene, 1
Faulting, Paleogene sequences, 4
felix, *Cibicides*, 11
Field Conference on the Paleogene, of the International Commission on Paleogene Stratigraphy, 1, 5, 49, 62
Field numbers, key, 67
Fissurina sp., 69
flintensis, *Orthophragmina*, 47
Proporocyclina, 47
Pseudophragmina (*Proporocyclina*), 42, 46, 47; pl. 6
(*Pseudophragmina*), 47
floridana, *Truncatulina*, 75
floridanus, *Cibicides*, 69, 71, 75; pl. 15
Cyclicargolithus, 30, 38, 51, 53, 56, 60
Florilus, 12
florinense, 7
florinensis, 11
Florilus-Eponides-Uvigerina assemblage, 13
florinense, *Florilus*, 7
florinensis, *Florilus*, 11
flos, *Micrantholithus*, 56, 61
Flysch assemblage, classic, characteristics, 19
Flysch faunas, Paleogene, Carpathians, 21
Paleogene, North Atlantic, 21
Foraminiferal biostratigraphy, 7, 13
Foraminiferal data, synthesis, 36
Foraminiferal stages of Mallory, 5, 29
Foraminiferal zones, 36
benthic, 62
planktic, 36, 62
Foraminifers, Angola, 21
Aquitaine Basin, 21
benthic, 1, 7, 12, 36, 41, 61
age assignments, 36
Año Nuevo section, 62
agglutinated, 7, 12, 19, 21
assemblages, 7, 72
calcareous, 7, 19
California Coast Ranges, 22, 41, 62
correlate West Coast zonal schemes, 62
Church Creek area, 58
Devils Den section, 7
Eocene, 41, 61
Lodo Gulch section, 7, 19
Miocene, 61
provincial stages, California, 1, 36
Refugian Stage, 58
San Lorenzo River section, 62, 63, 64, 65

Foraminifers—Continued

- benthic—Continued
 - Santa Cruz Mountains area, California, 61
 - Tertiary zonal scheme, 62
 - vertical distribution, 22
 - Zayante Creek section, 62, 69
- cosmopolitan, 21
 - Tunisia, 12
- faunas, Coast Range sections, 1
- flysch faunas, Paleogene, 12
- large, age, 42, 44
 - Arenal Sandstone, 41
 - Coast Ranges, California, 41
 - correlations, 44
 - deposition, 41
 - Devils Den area, 41, 44
 - distribution, 41
 - Domengine Formation, 42
 - Eocene, Coast Ranges, California, 41
 - evolution, 41
 - Miocene, 61
 - Mount Diablo area, 41, 42
 - Pacific Coast assemblages, 41
 - San Jose area, 42
 - Sveadal area, 44
 - Temblor Formation, 44
- Lodo Formation, type section, 7, 40
- Midway Group, 12
- Orphan Knoll, 12, 21
- Pacific affinities, 21
- planktic, 7, 13, 19, 36, 44, 49, 58, 62
 - Año Nuevo section, 51
 - Butano Sandstone, 19
 - Church Creek Formation, 49, 58, 59
 - Devils Den aqueduct section, 13
 - Locatelli Formation, 19
 - Lodo Formation, type, 7, 13
 - Lodo Gulch area, 13
 - marine strata, California, 58
 - occurrence, 59
 - Refugian Stage, 58
 - San Lorenzo River section, 52, 59
 - taxonomic notes, 59
 - Zayante Creek section, 52
 - zonation, 7, 13, 41, 49, 52, 59
 - Zones P19 and P20, 74
- siliceous molds, 72
- formosa*, *Cyclococcolithina*, 30
- Morozovella*, 42
- formosus*, *Coccolithus*, 53, 56, 60; pl. 9
- Cyclococcolithus*, 38
- fortunatus*, *Cibicides*, 10, 13, 16; pl. 5
- Franciscan basement rocks, 4
 - relation to granitic basement rocks, 4
- frankei*, *Guttulina*, 65, 75
- Fronicularia naheolensis*, 11
- tenuissima*, 69, 75
- frontosa*, *Subbotina*, 7, 12
- fulgens*, *Nannotetrina*, 31
- furcatolithoides*, *Sphenolithus*, 31, 56, 61; pl. 9
- Fursenkoina bramletti*, 65, 69, 71, 75; pl. 16
- californiensis*, 64, 75; pl. 16
- dibollensis*, 64, 75; pl. 16
- zetina indirecta*, 11, 17

G

- gallowayi*, *Pseudoglandulina*, 77
- Pseudonodosaria*, 69, 70, 71, 77; pl. 12
- Uvigerina*, 69, 70, 71, 77; pl. 14
- gamma*, *Cyclococcolithina*, 30, 56, 60
- gardnerae*, *Marginulina*, 11
- garzaensis*, *Anomalina*, 17, 64, 68, 74; pl. 17
- Anomalinoides*, pl. 5
- Plectina*, 64, 76; pl. 11
- Plectofronicularia*, 17
- Uvigerina*, 65, 68, 77; pl. 14
- nudorobusta*, *Uvigerina*, 69
- Gaudryina*, 19
- coalingensis*, 17; pl. 1
- cubana*, 75

Gaudryina—Continued

- gracilis*, 69, 70, 71; pl. 10
- jacksonensis coalingensis*, 10, 28
- laevigata*, 10, 16; pl. 1
- macrocamerata*, 10
- triangularis*, 69, 75; pl. 10
- Gavelinella beccariiiformis*, 19
- gemma*, *Globorotalia*, 59
- gemmifer*, *Discoaster*, 30
- germanica*, *Dorothia*, 17; pl. 1
- germanicus*, *Discoaster*, 30
- gibba globosa*, *Globulina*, 17
- gigas*, *Chiasmolithus*, 30, 38
- glabra*, *Marginulina*, 11
- Glandulina adunca*, 76
- inflata*, 77
- laevigata*, 71, 75
- sp., 11
- (*Glandulina*) *laevigata*, *Nodosaria*, 75
- Glauconite*, 7, 68, 69
- potassium-argon date, Lodo Formation, 7
- Globigerina*, 60
- ampliapertura*, 52
- angiporoides angiporoides*, 52, 59
- minima*, 52, 59
- angustiumbilicata*, 51
- eocaena*, 59
- juvenilis*, 51
- minima*, 59
- officinalis*, 59
- ouachitaensis*, 59
- praebuloides*, 51, 52, 59
- praeturritilina*, 59; pl. 8
- pseudovenezuelana*, 59
- senilis*, 59
- tripartita*, 51, 59
- turritilina praeturritilina*, 59
- utilisindex*, 59
- woodi*, 51
- Globigerinatheka index*, 59
- mexicana*, 59
- globigeriniiformis*, *Trochammina*, 16
- Globigerinita martini*, 59
- præstainforthi*, 51
- Globigerinoides higginsii*, 52, 59
- Globobulimina pacifica*, 65, 69, 71, 75; pl. 14
- sp., 10
- Globocassidulina globosa*, 65, 69, 75; pl. 4
- Globorotalia aragonensis*, 28
- californica*, 28
- cerroazulensis*, 59
- cerroazulensis*, 52
- gemma*, 59
- marginodentata*, 28
- munda*, 51, 60; pl. 8
- nana*, 51, 60; pl. 8
- opima nana*, 60
- pseudochapmani*, 28
- pseudocontinua*, 60
- sp., 60; pl. 8
- Globorotaloides carcoselleensis*, 59
- suteri*, 51, 52, 59
- wilsoni*, 59
- globosa*, *Cassidulina*, 11, 16, 75
- Globocassidulina*, 65, 69, 75; pl. 4
- Globulina gibba*, 17
- Globulina gibba globosa*, 17
- irregularis*, 75
- lacrima*, 17
- Glomospira charoides*, 17, 19
- Goesella* sp., 11
- Gonatosphaera eocenica*, 16
- sp., 17
- Goudkoff, P. R., cited, 36
- gracilis*, *Bolivina*, 64, 74
- Gaudryina*, 69, 70, 71; pl. 10
- Siphonodosaria*, 17
- gracilis*, *Morozovella*, 42
- grahami*, *Eofabiana*, 42, 47; pl. 6
- grandis*, *Chiasmolithus*, 30, 56; 60

grata, *Buliminella*, 12

- convoluta*, *Buliminella*, 10, 16, 22; pl. 2
- Great Valley, Paleogene, 4
- Great Valley sequence, 2
- Gredal Shale Member, Kreyenhagen Formation, 5, 13, 21, 22, 33
- Kreyenhagen Formation, age, 29
- Devils Den aqueduct section, 13, 22, 33, 36
- age, 29, 36
- time-stratigraphic relations, 33
- Devils Den area, 5, 19, 30, 48
- fauna, 19
- cosmopolitan, 21
- Pacific affinities, 21
- Media Agua Creek area, 33, 36
- Media Agua Creek section, boundary, 33
- age, 33, 36
- correlation, 36
- San Joaquin Valley, 33
- time-stratigraphic relations, 33
- samples, 5
- gredalensis*, *Pleurostomella*, 11
- grimsdalei*, *Cibicides*, 17
- Cibicides*, pl. 5
- Gumbelitra columbiana*, 59, 60; pl. 7
- Guttulina frankei*, 65, 75
- irregularis*, 65, 75
- gyrata*, *Nodosaria*, 10, 64; pl. 11
- Gyroidina condoni*, 64, 75
- orbicularis obliquata*, 17
- planata*, 11, 64, 69, 71; 75; pl. 17
- soldanii*, 65, 71, 75
- octocamerata*, 10, 17
- sp., 69, 71
- Gyroidinoides subangulata*, 16
- gyroscapulum*, *Lenticulina*, 10

H

- halkyardi*, *Nonion*, 65, 76
- hannai*, *Angulogerina*, 77
- Trifarina*, 65, 77
- hantkeni*, *Saracenaria*, 65
- Hantkenina* sp., 59
- Haplophragmium incisum*, 75
- Haplophragmoides*, 7
- coalingensis*, 10, 16
- deflata*, 65, 75; pl. 10
- eggeri*, 13, 16
- excavatus*, 11, 13, 17, 19
- nonionelloides*, 10
- obliquicameratus*, 19
- sp., 13, 69, 71; pl. 10
- sp. 1, 65; pl. 10
- spp., 10, 16, 64
- haydoni*, *Cibicides*, 65, 69, 75; pl. 15
- Planulina*, 75
- helianthus*, *Discoaster*, 30
- Helicopontosphaera ampliapertura* Zone, 74
- lophota*, 31
- seminulum*, 31
- Helicosphaera*, 61
- ampliapertura* Zone, 51
- bramlettei*, 56, 61
- carteri*, 51, 61
- compacta*, 56, 61
- euphratis*, 51, 53, 61
- intermedia*, 51, 53, 61
- lophota*, 56
- obliqua*, 53, 61
- perch nielsenasae*, 53, 61
- recta*, 53, 61
- seminulum*, 38, 56
- spp., 51
- Heliolithus riedeli* Zone, 7
- Heliothrus concinnus*, 38
- hesslandii*, *Dictyococcites*, 38
- heteromorphus*, *Sphenolithus*, 51, 61; pl. 9
- hezagona*, *Entosolenia squamosa*, 75
- Lagena*, 65, 71, 75

Hiatus, between Moreno Shale and type Lodo Formation, 29
 between Panoche Formation and Lodo Formation, 33
higginsii, *Globigerinoides*, 52, 59
hillae, *Reticulofenestra*, 31
hispidocostata, *Uvigerina*, 77
hispidula, *Ellipsonodosaria atlantisae*, 64, 68, 76
Nodosarella atlantisae, 76; pl. 16
hodgei, *Boldia*, 64, 68, 74; pl. 17
Cibicides, 74
Hoeglundina eocenica, 19
 sp., 11
holserica, *Nodosaria*, 69, 70, 76
hughesi, *Siphogenerina*, 71, 72, 77
humeri, *Bolivina*, 16; pl. 3
Hyperammina elongata, 19

I

ignota, *Amphimorphina*, 11, 16; pl. 2
Nodosarella, 16
impedens, *Bulimina*, 10, 17; pl. 3
imperialis, *Quinqueloculina*, 65, 77
Impiaster obscurus, 38
impolita, *Uvigerinella obesa*, 71, 77
inaequispira, *Subbotina*, 7, 12
incerta, *Operculina*, 74
incertus, *Ammodiscus*, 69, 71, 74
incisa, *Cyclammina*, 17, 69, 75; pl. 1
Nonionella, 76
Nonionina, 76
kernensis, *Nonionella*, 71, 76
incisum, *Haplophragmium*, 75
kernensis, *Nonion*, 76
index, *Globigerinatheka*, 59
indirecta, *Fursenkoina zetina*, 11, 17
indiscriminata, *Planorotalites*, 28
Valvulineria, 28
inflata, *Bulimina*, 65, 70
alligata, *Bulimina*, 71, 72, 74; pl. 14
Glandulina, 77
Pseudonodosaria, 64, 69, 71, 77
Rhabdosphaera, 29, 31
inflexa, *Cyclolithella*, 38
inornata, *Lenticulina*, 11, 65, 69, 76
Robulus, 76
instabilis, *Bulimina*, 10, 16
insuetus, *Lenticulina*, 64, 76; pls. 11, 12
Robulus, 76
intermedia, *Helicosphaera*, 51, 53, 61
 International Stratigraphic Guide, 5, 39
 International Commission on Paleogene Stratigraphy, 1, 5, 49, 62
 Introduction, 1, 5, 22, 41, 49, 62
inversus, *Cyclococcolithus*, 38
involutus, *Fasciculithus*, 31
irregularis, *Globulina*, 75
Guttulina, 65, 75
 Israelsky, M. C., cited, 5, 29, 39
 sample collection, 5, 7
 samples, Lodo Formation, type area, 7, 39
 section A, 39, 40
 section B, 39, 40
Isthmolithus recurvus, 31, 53, 61
recurvus to *Coccolithus subdistichus* zone, 31

J, K

jacksonensis, *Dentalina*, 11, 16, 65, 75
Siphonina, 11, 17; pl. 4
Uvigerina, 69
coalingensis, *Gaudryina*, 10, 28
jenkinsii, *Amphimorphina*, 68
juvenilis, *Globigerina*, 51
judas, *Anomalinoidea*, 10
Karrerella fallax, 11
Karrerella, 7
arenasensis, 16
elongata, 11, 64, 68, 75; pl. 11

Karrerella—Continued
mediaguensis, 13, 16, 19; pl. 1
monumentensis, 17; pl. 1
washingtonensis, 65, 68, 73, 75; pl. 11
 sp., 69
keeni, *Anomalinoidea*, 10, 17; pl. 5
kelleyi, *Vaginulinopsis*, 10, 16
kernensis, *Cibicides*, 11, 17, 28
Nonion incisum, 76
Nonionella incisa, 71, 76
kerni, *Plectofrondicularia*, 10; pl. 2
Vaginulinopsis mexicana, 17; pl. 2
kingi, *Cyclococcolithina*, 56, 60
 Kinney, D. M., cited, 39, 40
 Kleinpell, R. M., cited, 1, 36, 62, 68, 71, 72, 73
kleinpelli, *Bolivina*, 17, 64, 74; pl. 13
kressenbergensis, *Nodogenerina*, 11, 17
 Kreyenhagen Formation, Avenal Sandstone Member, 22, 28, 29, 32, 33, 41
 Canoas Siltstone Member, 36
 Gredal Shale Member, 5, 13, 19, 21, 22, 29, 30, 33, 36, 48
 Point of Rocks Sandstone Member, 5, 22, 29, 33, 36
 Krieh, 12
kuepperi, *Discoasteroides*, 31

L

La Honda basin, 4
lacinata, *Bulimina sculptilis*, 65, 68, 74; pl. 14
lacrima, *Bulimina semicostata*, 17
Globulina, 17
laevigata, *Gaudryina*, 10, 16; pl. 1
Glandulina, 71, 75
Nodosaria (Glandulina), 75
Lagena acuticosta, 11, 71, 75; pl. 11
becki, 65, 75
costata, 65, 71, 75
crowleri, 11
hexagona, 65, 71, 75
semistriata, 65, 71, 75; pl. 11
substriata, 16
sulcata, 69, 75
vulgaris, 65, 71, 75
 Laiming, B zones, 42
laimingi, *Cibicides*, 22
Lenticulina, 28
 Lambert Shale, coccoliths, 52
 Sauceian Stage, 71
 Zayante Creek section, 52, 70, 71
lamposa, *Spiroloculina*, 11, 16
Lanternithus minutus, 56, 61; pl. 9
larvalis, *Pedinocyclus*, 53, 56, 61
latejugata, *Nodosaria*, 10, 16, 28
Lenticulina, 7
alatolimbatus, 10
antipoda, 17
antipodium, 11
barbati, 71, 75; pl. 11
calcar, 69, 71, 75; pl. 11
chehalisensis, 65, 75
clypeiformis, 69, 71, 76; pl. 11
coaledensis, 65, 76
coloratus, 71
convergens, 17
cultratus, 76
gyroscaprum, 10
inornata, 11, 65, 69, 76
insuetus, 64, 76; pls. 11, 12
laimingi, 28
mayi, 69
midwayensis, 10, 64, 67, 76; pl. 12
pseudocultrata, 10
pseudocultratus, 71, 76; pl. 12
pseudorotulata, 65, 71, 76
pseudovortex, 10, 16, 64, 68, 71, 76; pl. 12
rosetta, 11
rotulata, 10
simplex, 69, 71, 76
terryi, 10, 16
theta, 10, 17

Lenticulina—Continued
turbinata, 11, 17
ulatisensis, 16, 28; pl. 2
vortex, 10, 16
weaveri, 10
welchi, 17, 65, 68, 69, 76; pl. 12
 sp., Fairchild and others, 71, 76
 sp. A, 71
 sp., pl. 12
 spp., 10, 13, 16, 64, 69, 71; pl. 12
lenticularis, *Discoaster*, 30
Lepidocyclina, 1
antillea, 47
lepidula, *Nodogenerina*, 11, 17
Nodosaria, 77
Stilostomella, 65, 69, 71, 77
tiltisi, *Pseudohastigerina*, 59, 60; pl. 7
Pullenia, 60
linaperta, *Subbotina*, 7, 12, 19
 Lincoln Creek Formation, fauna, 68
 foraminifers, benthic, 68, 69
 Washington, 73
 Lincoln Formation, 68
 Lipps, J. H., cited, 60
Lithostromation perdurum, 38, 56
 Locality data, 60
 Locality descriptions, 48
 Locatelli Formation, foraminifers, benthic, agglutinated, 19, 21
 foraminifers, benthic, calcareous, 19, 21
 planktic, 19, 21
 Lompico area, 13
 nannofossils, 38
 paleobathymetry, 21
 Santa Cruz Mountains, 5, 13, 19, 21
 Smith Grade area, 38
 Smith Grade-Empire Grade area, 5, 13
 Lodo Formation, 21, 22, 32, 33, 36, 39
 definition, 39
 Devils Den aqueduct section, 5, 7, 13, 22, 29
 age, 29, 36
 correlation, 13
 foraminiferal biostratigraphy, 13
 foraminifers, benthic, 13, 21
 planktic, 13, 21
 paleobathymetry, 13, 21
 Devils Den area, 5, 7, 19, 28, 36
 fauna, cosmopolitan, 21
 Pacific affinities, 21
 foraminiferal biostratigraphy, 7
 foraminifers, benthic, 19
 benthic, agglutinated, 7, 10, 11, 19, 21
 calcareous, 7, 10, 11, 19
 zones, 7, 10, 22
 planktic, 7, 13, 19
 zones, 7
 reference sections, 40
 type section, 7, 10, 11, 12, 40
 formation boundary, time-transgressive, 36
 glauconite, 7
 Lodo Gulch area, 5, 13, 19, 22, 28, 29, 36, 39, 40
 age, 29, 36
 foraminiferal biostratigraphy, 7
 paleobathymetry, 12
 stratigraphic relations, 29
 Lodo Gulch section, 7, 39, 40
 Media Agua Creek area, 28, 32, 36
 age, 33, 36
 stratigraphic relations, 33
 Media Agua Creek section, 36
 Midway-type fauna, 13
 molluscan fauna, 7
 nannoplankton, 13, 28, 29, 30, 31, 32, 36
 ostracode faunas, 12
 paleobathymetric history, 12
 paleobathymetry, 12, 19, 21
 reference section, 39, 40
 principal, 40
 Israelsky samples, 5
 San Joaquin Valley, 5

- Lodo Formation—Continued
 Santa Cruz Mountains, 19
 section A, Brabb, 39, 40
 Israelsky, 39, 40
 section B, Brabb, 39, 40
 Israelsky, 39, 40
 section C, Brabb, 39, 40
 Brabb, Clark, and Throckmorton, 39, 40
 section I-X, White, 39, 40
 stratigraphic relations, 29, 39
 Tumey Hills, Calif., 39
 type area, 39
 type locality, 5, 28, 29, 36, 39
 type section, 5, 7, 10, 11, 12, 13, 19, 29, 32, 39, 40
 age, 7, 19, 32, 36
 correlation, 7, 36
 Lodo Gulch area, 5
 Midway-type fauna, 7, 12
 sample materials, 5, 40
 stratigraphic relations, 7, 29
lodoensis, *Discoaster*, 30, 38, 51, 53, 56, 61
Eponides, 11, 28
Uvigerina, 28
myriamiae, *Uvigerina*, 11, 16; pl. 3
 Lompico area, 13
 Lompico area, Santa Cruz County, Calif., 5
 Santa Cruz Mountains, 13, 19
longiscata, *Nodosaria*, 64, 69, 71, 76
lopetrigoi, *Amphistegina*, 46
Eoconuloides, 42, 46; pl. 6
Lophodolichus mochlophorus, 31
nascens, 31
reniformis, 31, 56, 61
lophota, *Helicopontosphaera*, 31
Helicosphaera, 56
 Lower Eocene-middle Eocene boundary, 60
 Lower Miocene-middle Miocene boundary, 51
 Lower Narizian, *Bulimina corrugata* Zone, 68
 Lower Narizian-upper Narizian boundary, 68, 73
Loxocochea, 12
Loxostomoides applinae, 11; pl. 3
luminus, *Cyclicargolithus*, 56, 60
Cyclococcolithus, 60
- M
macellus, *Ellipsolichus*, 31
macilenta, *Bulimina*, 10, 16; pl. 3
macrocamerata, *Gaudryina*, 10
magnicrassus, *Coccolithus*, 30
 Mallory, V. S., cited, 1, 5, 29, 33, 36, 42, 62
margareta, *Cassidulina*, 71, 74
marginata, *Bolivina*, 65, 69, 70, 71, 74; pls. 10, 13
Orthophragmina, 46
Discoecyclina (*Discoecyclina*), 42, 46
adelaidana, *Bolivina*, 71, 72
marginodentata, *Globorotalia*, 28
Marginulina adunca, 11, 65, 71, 76
alazaensis, 65, 69, 76
curvatura, 10
dubia, 69, 76
exima, 65, 76
gardnerae, 11
glabra, 11
sischoae, 11
subbullata, 10, 16, 64, 69, 76; pl. 12
subrecta, 69, 76
 sp., 69; pl. 12
 spp., 64, 71
 Marine sedimentary deposits, correlation, 36
Markalius astroporus, 31
markleyana, *Planularia*, 65, 68, 76
 Martin, L. T., Lodo Formation, definition, 39, 40
 Martin, L. T., cited, 29, 39, 40
 Martinez Stage, 7
martinezensis, *Cibicides*, 11, 16
Cibicoides, 19
 Martini, E., cited, 38
martini, *Globigerinita*, 59
Martinottiella eocenica, 16, 64, 76; pls. 1, 11
patens, 69, 71, 76; pl. 11
mauricensis, *Cibicidina*, 17
mayi, *Lenticulina*, 69
Siphogenerina, 71, 72, 77; pl. 14
 McDougall, Kristin, cited, 68
mckannai, *Acarinina*, 19
 McKeel, D. R., cited, 28, 60
 Media Agua Creek, area, 22
 area, comparison, 33
 Kern County, Calif., 32
 type sections, benthic foraminiferal zones, 33
 reference section for lower Tertiary, California, 33
 section, San Joaquin Valley, 33
mediaguaensis, *Karrerella*, 13, 16, 19; pl. 1
 Megafossils, age assignments, 36
Melonis pompilioides, 69, 71, 76
 sp., 65
metastypus, *Discoasteroides*, 31
mezicana, *Globigerinatheka*, 59
Osangularia, 17; pl. 4
Vaginulinopsis, 28
kerni, *Vaginulinopsis*, 11, 17; pl. 2
nudicostata, *Vaginulinopsis*, 10, 17; pl. 2
mezicanus, *Bolivinoidea*, 65, 74; pl. 13
Proroporus, 74
 Mickey, M. B., cited, 28
miera, *Pseudohastigerina*, 59, 60; pl. 7
Micrantholithus altus, 56; pl. 9
basquensis, 56
entaster, 31
flos, 56, 61
vesper, 31
microcostata, *Bulimina*, 64, 68, 69, 74; pl. 14
 Microfossils, planktic, calcareous, 59
 planktic, San Lorenzo River section, 73
micrus, *Nonion*, 60
 Middle Eocene-upper Eocene boundary, 39, 52, 53, 73
 Middle Eocene-upper Eocene boundary, international
 usage, 59
 Midway Group fauna, 7, 12, 13, 19
midwayana, *Osangularia*, 17
midwayensis, *Cristellaria*, 76
Lenticulina, 10, 64, 67, 76; pl. 12
Robulus, 76
subcylindrica, *Chiloguembelina*, 28
minima, *Eponides*, 17
Globigerina, 59
angiporoides, 52, 59
minuta, *Plectofrondicularia*, 11, 17; pl. 2
Quinqueloculina, 64, 77; pl. 11
minutus, *Lanternithus*, 56, 61; pl. 9
miocenica, *Nonionella*, 65, 69, 70, 71, 76; pl. 16
Plectofrondicularia, 69, 76
Pullenia, 71, 77; pl. 17
directa, *Plectofrondicularia*, 71, 76; pl. 12
miriamae, *Uvigerina lodoensis*, 11, 16; pl. 3
mirus, *Discoaster*, 30, 38
mitreus, *Fasciculithus*, 31
mochlophorus, *Lophodolichus*, 31
 Mohnian Stage of Klempell, 1
 Molluscan fauna, living, 1
 Lodo Formation, 7
 Martinez Stage, 7
 Trinidad, 1
 Vaqueros Sandstone, 1
 Monterey Formation, 51
 Año Nuevo section, 51, 71
monumentensis, *Karrerella*, 17; pl. 1
 Moreno Shale, 7, 29
 Lodo Gulch, 39
moriformis, *Sphenolithus*, 31, 38, 51, 53, 56, 61
 Morin, R. W., cited, 28
Morozovella aequa, 19
aragonensis, 19
formosa, 42
 Zone, 42
gracillia, 42
subbotinae, 19, 42
velascoensis, 19
 Mount Diablo area, foraminifers, large, 41, 42
 locality descriptions, 48
 Mount Diablo, Calif., 2, 4, 41
multicostata, *Dietyomitra*, 28
Siphogenerina, 71, 72, 77; pl. 14
multilineata, *Dentalina*, 11
Plectofrondicularia packardii, 65, 68, 69, 76; pl. 13
multilobata, *Pullenia*, 71, 77
multiplora, *Discolithina*, 51, 61
Pontosphaera, 38
multiradiatus, *Discoaster*, 30
munda, *Globorotalia*, 51, 60; pl. 8
- N
naheolensis, *Fronicularia*, 11
nana, *Globorotalia*, 51, 60; pl. 8
Globorotalia opima, 60
 nannofossils, 22, 38, 60, 62, 74
 calcareous, 60
 California, 22
 Santa Cruz Mountains, Calif., 38
 distribution, 22
 Miocene, 74
 nannoplankton, age, 29, 33
 biostratigraphy, central Coast Ranges, Calif., 22
 calcareous, 13, 22, 38, 44
 chronology, Bukry, 31, 33
 data, Bulitian Stage, 36
 Cheneyan Stage, 36
 Narizian Stage, 36
 Penutian Stage, 36
 synthesis, 36
 Ulatian Stage, 36
 Ynezian Stage, 36
 Zemorrian Stage, 36
 Devils Den section, age, 28
 Paleogene, Santa Cruz Mountains, 38
 Nannoplankton zones, 1, 22
 Bear Creek area, 39
 Bukry, 22, 28, 29, 33, 44
 Butano Sandstone, 38, 39
 central Coast Ranges, California, 22
 Devils Den aqueduct section, 13, 22, 28, 29
 Gredal Shale Member, 13, 29
 Locatelli Formation, 38
 Lodo Formation, 29
 type, 29, 32
 Lodo Gulch area, 29
 Lompico area, 38
 Martini, 22
 Media Agua Creek area, 32
 Paleogene of California, summary, 36
 Panoche Formation, 29
 San Joaquin Valley, 33
 Santa Cruz Mountains, 38
 Smith Grade area, 38
 Twobar Shale Member, 39
Nannotetrina cristata, 38
fulgens, 31
quadrata, 29, 31
 Zone, 13, 29, 31, 33
 boundary, 29
 Narizian Stage, 28, 33, 36, 42, 53, 59, 62, 67, 68, 69, 70,
 72, 73
 age, 36
 boundary, 29, 59, 73
 of Mallory, 42
 Narizian Stage-Refugian Stage boundary, 59
nascens, *Lophodolichus*, 31
Nautilus calcar, 75
pompilioides, 76
navaroanna, *Ramulina*, 17
Neochiastozygus dubius, 39
Neococcolithes dubius, 31
protenus, 31
Neoeponides sp., pl. 15
neogammation, *Cyclicargolithus*, 60
 Nilsen, T. H., cited, 4
nobilis, *Discoaster*, 30
nodifer, *Discoaster*, 30, 56, 61; pl. 9
nodifera, *Siphogenerina*, 69, 70, 71, 77; pls. 10, 14, 15
Nodogenerina adolphina, 17

Nodogenerina—Continued

- advena*, 77
kressenbergensis, 11, 17
lepidula, 11, 17
rohri, 76
wegimanni, 77
Nodosarella advena, 16; pl. 4
atlantisae hispidula, 76; pl. 16
constricta, 16
ignota, 16
Nodosaria affinis, 11
arundinea, 10, 16
boffalorae, 76
coccosensis, 75
deliciae, 10, 16
ewaldi, 10, 16
gyrata, 10, 64; pl. 11
holserica, 69, 70, 76
latejugata, 10, 16, 28
lepidula, 77
longiscata, 64, 69, 71, 76
parezilis, 69, 71
sentifera, 71, 76; pl. 11
pyrula, 64, 71, 76
velascoensis, 17
(Glandulina) laevigata, 75
sp., 65, 71
spp., 64
nonaradiatus, *Discoaster*, 30
Nonion halkyardi, 65, 76
incisum kernensis, 76
micrus, 60
Nonionella ansata, 17
costifera, 69, 71, 76
incisa, 76
kernensis, 71, 76
miocenica, 65, 69, 70, 71, 76; pl. 16
nonionelloides, *Haplophragmoides*, 10
Nonionina costifera, 76
incisa, 76
quinqueloba, 77
nudicostatus, *Vaginulinopsis mexicana*, 17; pl. 2
nudorobusta, *Uvigerina garzaensis*, 69
Nummulites parvula, 46
striatoreticulata, 45
striatoreticulatus, 42, 45; pl. 6
trinitatis, 45
willcozi, 42, 45; pl. 6
sp., 42, 45
nuttalli, *Bifarina*, 11, 16, 28; pl. 2
Pleurostomella, 17
Nuttallides truempyi, 7, 10, 12, 13, 16, 19; pl. 4

O

- oamaruensis*, *Reticulofenestra*, 56, 61
obesa, *Cyclammina cancellata*, 75; pl. 10
impolita, *Uvigerina*, 71, 77
oblitua, *Helicosphaera*, 53, 61
obliquata, *Gyroidina orbicularis*, 17
obliquicameratus, *Haplophragmoides*, 19
obscurus, *Impiaster*, 38
octocamerata, *Gyroidina soldanii*, 10, 17
officinalis, *Globigerina*, 59
Oligocene, Europe, 1
Oligocene-Miocene boundary, 51
opelagicus, *Coccolithus*, 38
Operculina incerta, 74
Operculinoides willcozi, 45
opima nana, *Globorotalia*, 60
orbicularis obliquata, *Gyroidina*, 17
planata, *Gyroidina*, 11, 64, 69, 71, 75; pl. 17
Orridosalis umbonatus, 10, 16, 64, 69, 71, 76; pl. 17
ornator, *Dentalina spinosa*, 64, 75; pl. 11
ornatus, *Discoaster tani*, 31
Orphan Knoll, 12
orphanensis, *Bulimina*, 12, 13
orphanknolli, *Sphenolithus*, 38
Orthomorphina rohri, 65, 69, 71, 76; pls. 10, 12
Orthophragmina clarki, 47
flintensis, 47

Orthophragmina—Continued

- marginata*, 46
orthostylus, *Tribrachiatus*, 29, 31, 51, 53, 56, 61
Osangularia mexicana, 17; pl. 4
midwayana, 17
plummerae, 7
tenuicarinata, 11, 16; pl. 4
sp., 19
Osburn, E. D., cited, 44
Ostracode fauna, 12
ouachitaensis, *Globigerina*, 59
ovata, *Bulimina*, 10, 77
Praeglobbulimina, 64, 65, 71, 77

P

- pachyderma*, *Cibicides*, 17, 64, 68, 75
Truncatulina, 75
Pacific Ocean, 4
pacifica, *Cyclammina*, 65, 75
Globbulimina, 65, 69, 71, 75; pl. 14
packardi, *Plectofrondicularia*, 65, 68, 69, 76; pl. 13
multilineata, *Plectofrondicularia*, 65, 68, 69, 76; pl. 13
packardi, *Plectofrondicularia*, 76
Paleobathymetry, 12, 13
Paleogene, 12
Paleocene stages of Mallory, 1
paleocenica, *Pleurostomella*, 11, 17, 19
Paleogene, California, 4, 22, 38
California, summary, 36
depositional patterns, 4
faunas, Coast Ranges, 1, 22
provincial stages, California, 38
rocks, complete section, Devils Den area, 41
east of San Andreas fault, 2
stratigraphy, 41
west of San Andreas fault, 2
Santa Cruz Mountains, nannoplankton, 38
sequences, California, 5
displaced along secondary lateral faults, 4
displaced laterally along San Andreas, 4
foraminiferal, 5
stages, California, based on benthic foraminifers, 1
California, zonations based on planktic foraminifers and nannoplankton, 1
Paleogeography, central California, 4
Panoche Formation, Devils Den aqueduct section, 22
Devils Den area, 28, 29
age, 29
Media Agua Creek area, 33
parezilis, *Nodosaria*, 69, 71
sentifera, *Nodosaria*, 71, 76; pl. 11
partschiana, *Epistomina*, 10, 17
parvula, *Amphistegina*, 42, 46; pl. 6
Nummulites, 46
parvulus, *Eoconuloides*, 46
patagonica, *Subbotina*, 42
patens, *Clavulina*, 76
Martinottiella, 69, 71, 76; pl. 11
Pedinocyclus larvalis, 53, 56, 61
pelagicus, *Coccolithus*, 30, 38, 53
Pemma angulatum, 38, 56, 61
basquensis, 61
pennyi, *Ammodiscus*, 10, 16; pl. 1
penonensis, *Asterocyclina*, 47
Penutian Stage, 28, 29, 33, 36, 44
age, 29, 33, 36
Mallory, 44
type section, 33
perch nielsenasae, *Helicosphaera*, 53, 61
perdurum, *Lithostromation*, 38, 56
perplexa, *Silicosigmoidina*, 16
Pierce, R. L., cited, 1, 49, 58
pilulifer, *Reophax*, 64, 69; pl. 10
pilulifera, *Reophax*, 71, 77
pisciformis, *Bolivina*, 17
plana, *Discolithina*, 30, 31, 51, 53, 61
Pontosphaera, 38

- planata*, *Gyroidina orbicularis*, 11, 64, 69, 71, 75; pl. 17
Planktic, defined, 1
Planktic assemblages, San Lorenzo River section, 59
Planktic foraminifers, Devils Den aqueduct section, 13
Lodo Gulch area, 13
zonations, 1, 7, 13, 41
Plankton, calcareous, Año Nuevo area, 51
calcareous, discussion, 59
Eocene, 59
Santa Cruz Mountains, 49
Santa Lucia Range, 49
Planktonic, defined, 1
Planktonic foraminiferal zones, 1, 22
Año Nuevo section, 51, 74
Berggren, 22
Blow, 22
Butano Sandstone, 19
Church Creek area, 58
Devils Den aqueduct section, 13
Hardenbol and Berggren, 22
Kings Creek section, 52
Lodo Formation, 7, 12, 19
Lodo Gulch area, 5, 12, 19
Paleogene of California, summary, 36
Rices Mudstone Member, 52, 53
San Lorenzo Formation, 52, 53
San Lorenzo River section, 52, 59, 73
Santa Cruz Mountains, 49
Santa Lucia Range, 49
Twohar Shale Member, 52, 53
Planorotalites, 28
indiscriminata, 28
pseudomenardi, 19
Zone, 7
pseudoscutula, 59
Planularia markleyana, 65, 68, 76
tolmani, 69
truncana, 11, 17
Planulina haydoni, 75
Plectina cubensis, 13, 17
garzaensis, 64, 76; pl. 11
Plectofrondicularia californica, 71, 76
garzaensis, 17
kerni, 10; pl. 2
minuta, 11, 17; pl. 2
miocenica, 69, 76
directa, 71, 76; pl. 12
packardi, 65, 68, 69, 76; pl. 13
multilineata, 65, 68, 69, 76; pl. 13
packardi, 76
stratigraphic range, 73
vaughani, 65, 69, 71, 76; pl. 13
whitei, 10; pl. 2
spp., 64
plectopons, *Zygodiscus*, 31
Pleurostomella acuta, 11, 16
alazanensis cubensis, 17
altamans, 64, 68, 77; pl. 16
gredalensis, 11
nuttalli, 17
paleocenica, 11, 17, 19
plousios, *Amphipyndax*, 28
plummerae, *Osangularia*, 7
Stilostomella, 10
Point of Rocks, 36
Point of Rocks Sandstone, Kreyenhagen Formation, Devils Den aqueduct section, 5, 22, 29, 36
Kreyenhagen Formation, Devils Den aqueduct section, age, 29
Devils Den aqueduct section, nannoplankton, 29
samples, 5
time-stratigraphic relations, 33
Media Agua Creek area, 33, 36
Media Agua Creek section, 33, 36
age, 33, 36
correlation, 36
nannoplankton, 33
San Joaquin Valley, 33
time-stratigraphic relations, 33

pompilioides, *Melonis*, 69, 71, 76
Nautilus, 76
Pontosphaera multipora, 38
plana, 38
 Poore, R. Z., cited, 29, 33, 52, 63
praebuloides, *Globigerina*, 51, 52, 59
praecursorius, *Cibicides*, 11
Praeglobobulimina ovata, 64, 65, 71, 77
pupoides, 64, 71, 77
praestainforthi, *Globigerinita*, 51
praeturritilina, *Globigerina*, 59; pl. 8
Globigerina turritilina, 59
predistentus, *Sphenolithus*, 38, 53, 56, 61
primus, *Eponides*, 7, 11, 16
principiensis, *Dorothia*, 17
Tritaxilina, 13, 16
Proporocyclina flintensis, 47
 (*Proporocyclina*) *clarki*, *Pseudophragmina*, 47; pl. 6
flintensis, *Pseudophragmina*, 42, 46; pl. 6
teres, *Pseudophragmina*, 47
Proroporus mexicanus, 74
protenus, *Neococcolithus*, 31
protoannula, *Cyclococcolithina*, 60
pseudochapmani, *Globorotalia*, 28
pseudocontinua, *Globorotalia*, 60
pseudocultrata, *Lenticulina*, 10
pseudocultratus, *Lenticulina*, 71, 76; pl. 12
Robulus, 76
pseudogammation, *Cyclococcolithina*, 30
Pseudoglandulina gallowayi, 77
Pseudohastigerina lillisi, 59, 60; pl. 7
micra, 59, 60; pl. 7
pseudomenardii, *Planorotalites*, 19
Pseudonodosaria gallowayi, 69, 70, 71, 77; pl. 12
inflata, 64, 69, 71, 77
Pseudoparrella danvillensis, 64
Pseudophragmina clarki, 42
psila, 28
teres, 42
 (*Proporocyclina*) *clarki*, 47; pl. 6
flintensis, 42, 46, 47; pl. 6
teres, 47
 (*Pseudophragmina*) *flintensis*, 47
 sp., pl. 6
 (*Pseudophragmina*) *flintensis*, *Pseudophragmina*, 47
pseudoradians, *Sphenolithus*, 31, 56, 61; pl. 9
pseudorotulata, *Lenticulina*, 65, 71, 76
pseudorotulatus, *Robulus*, 76
pseudoscutula, *Planorotalites*, 59
pseudoumbilica, *Reticulofenestra*, 31
pseudoungerianus evolutus, *Cibicides*, 65, 69, 71, 75;
 pl. 16
pseudovenezuelana, *Globigerina*, 59
pseudovortex, *Lenticulina*, 10, 16, 64, 68, 71, 76; pl. 12
Robulus, 76
pseudovuellerstorfi, *Cibicoides*, 16
psila, *Pseudophragmina*, 28
pulcher, *Transversopontis*, 31
pulcheroides, *Discolithus*, 61
Transversopontis, 56, 61
Pullenia eocenica, 10, 16
lillisi, 60
miocenica, 71, 77; pl. 17
multilobata, 71, 77
quinqueloba, 64, 77; pl. 17
angusta, 17
reussi, 16
salisburyi, 69, 77
punctulata, *Daktylethra*, 56, 60; pl. 9
pupoides, *Bulimina*, 16, 77
pupoides, *Praeglobobulimina*, 64, 71, 77
pyrula, *Bulimina*, 16
Nodosaria, 64, 71, 76
 Q. R
quadrata, *Nannotetrina*, 29, 31
quadrulata, *Dentalina*, 69, 71, 72, 75; pl. 11
quinqueloba, *Nonionina*, 77
Pullenia, 64, 77; pl. 17
angusta, *Pullenia*, 17

Quinqueloculina imperialis, 65, 77
minuta, 64, 77; pl. 11
radians, *Sphenolithus*, 31, 38, 51, 53, 56, 61
Radiolarians, Cretaceous, 28
Ramulina navaroanna, 17
raphanus transversa, *Siphogenerina*, 77
recta, *Helicosphaera*, 53, 61
Rectoglandulina sp., 10
recurvus, *Isthmolithus*, 31, 53, 61
 Reference list, taxa, 74
 Reference section, Paleogene stratigraphy of
 California, 5
 References cited, 78
 Refugian Stage, 53, 58, 59, 68, 69, 70, 73
 boundaries, 53, 69, 73
 California, 68, 69
 Church Creek Formation, 58
 faunal assemblage, 69, 70, 73
 foraminifera, marine strata, California, 58
 Oregon, 68
 type area, California, 73
 Washington, 68
 Refugian faunas, basal, McDougall, 68
 San Lorenzo River section, 73
 Refugian-Zemmorrian boundary, 53, 69, 73
regina, *Anomalinoidea*, 16
 Regional setting, 2
 Relizian Stage, 52, 62, 72, 74
 boundary, 52
 faunal assemblages, 72, 74
reniformis, *Lophodolichus*, 31, 56, 61
Reophaz pilulifer, 64, 69; pl. 10
pilulifera, 71, 77
 spp., 71
reticulata, *Reticulofenestra*, 56, 61; pl. 9
Reticulofenestra, 61
dictyoda, 31, 38
hillae, 31
oamaruensis, 56, 61
pseudoumbilica, 31
reticulata, 56, 61; pl. 9
samodurovi, 31
umbilica, 31, 38, 53, 56, 61; pl. 9
 Zone, 53
 spp., 51, 53, 56
reussi, *Pullenia*, 16
Rhabdammina, 7, 12, 13, 19
eocenica, 10, 13, 16, 19, 64, 77; pl. 10
Rhabdosphaera, 61
crebra, 31
inflata, 29, 31
 Zone, 31
tenuis, 56, 61
 sp., 38
 spp., 56
 Rices Mudstone Member, San Lorenzo Formation, age,
 52
 San Lorenzo Formation, coccoliths, 51, 53
 foraminifers, benthic, 68
 planktic, 52
 Kings Creek section, 52
 San Lorenzo River section, 52, 68
 age, 68, 69
 Zayante Creek section, 52
 zone assignment, 68
 zone P14-P15 boundary, 52, 53
richardi, *Spiroplectammina*, 16, 19, 64, 77; pls. 1, 10
Robulus barbati, 75
chehalisensis, 75
chiranus, 69, 76
coaledensis, 76
cultratus, 76
inornata, 76
insuetus, 76
midwayensis, 76
pseudocultratus, 76
pseudorotulatus, 76
pseudovortex, 76
simplex, 76
welchi, 76
robusta, *Baggina*, 71, 72, 74; pl. 15

rohri, *Nodogenerina*, 76
Orthomorphina, 65, 69, 71, 76; pls. 10, 12
Truncorotaloides, 52, 59
Rosalina araucana, 77
rosetta, *Lenticulina*, 11
Rotulina umbonata, 76
Rotularia (Spirogyllus) tinajasensis, 28
 (*Rotularia*), *Spirogyllus*, 41
rotulata, *Lenticulina*, 10
rubiginosus, *Anomalinoidea*, 19
Rzehakina, 19
epigona-lata, 19
 S
saipanensis, *Discoaster*, 30, 38, 53, 56, 61; pl. 9
salinasensis, *Anomalina*, 71, 74; pl. 17
 Salinian block, 2
salisburyi, *Pullenia*, 69, 77
samodurovi, *Reticulofenestra*, 31
 Sample materials, 5
 numbers, key, 67
 Samples, Church Creek Formation, 60
 Israelsky collection, 5, 7
 Lodo Gulch, Brabb, 7, 10, 11, 39
 Paleogene, from four sections, 5
 Santa Lucia Range, 60
 San Andreas fault, 4
 San Joaquin Valley, faunal assemblages, 72
 rock units, time-transgressive, 33
 west side, 22
 San Jose area, foraminifers, large, 42
 locality descriptions, 48
 San Lorenzo Formation, Kings Creek section, 59
 Rices Mudstone Member, 51, 52, 53, 68, 69
 glauconite bed, 68
 San Lorenzo River section, 52, 59, 63
 Twobar Shale Member, 38, 39, 52, 53, 59, 63, 67, 68,
 69
 San Lorenzo River section, 52, 59, 60, 73
 age, 52, 67
Amphimorphina jenkinsi Zone, 67, 73
 biostratigraphy, 53
Bulimina corrugata Zone, 73
 Cañada de Santa Anita, 73
 deep water facies, 68
 faunal assemblages, 70
 Lincoln Creek Formation, 68
 lithostratigraphy, 53
 microfossils, planktic, 59, 73
 paleoecologic facies, 73
 Refugian Stage, 53, 73
 Santa Cruz Mountains, 49, 63
 type area, 73
 San Lorenzo syncline, 70
sanctaerucis, *Stilostomella*, 71, 77
 Santa Cruz Mountains, 13
 area, foraminifers, benthic, 61
 California, 2, 4, 5
 Paleogene, 38
 biostratigraphy, 38
 nannofossils, calcareous, 38
 Lompico area, 19
 nannoplankton, Paleogene, 38
 plankton, calcareous, 49
 San Lorenzo River section, 63
 Smith Grade-Empire Grade area, 13
 Zayante Creek section, 49, 70
 Santa Lucia Range, Church Creek area, 49, 58, 59
 plankton, calcareous, 49
 samples, 60
Saracenaria hantkeni, 65
schencki, 65, 68, 69, 71, 73, 77; pl. 12
 Saucian Stage, 52, 59, 71, 72, 74
 base, 71
 faunal assemblages, 72, 74
 Miocene, 74
 Saucian Stage-Relizian Stage boundary, 52
 Saucian Stage-Zemmorrian Stage boundary, 52, 59, 72,
 73, 74
saundersi, *Cristellaria*, 77

- saundersi—Continued
Vaginulinopsis, 19, 64, 77; pl. 12
sazea, *Thoracosphaera*, 38
scabrata, *Bolivina*, 64, 68, 74; pl. 13
schaubi, *Fasciculithus*, 31
Schenck, H. G., cited, 68
schencki, *Caucasina*, 22
Saracenaria, 65, 68, 69, 71, 73, 77; pl. 12
Sigmomorphina, 65, 68, 73, 77
Schmidt, R. R., cited, 7
Schoellhamer, J. E., cited, 39, 40
scrippsae, *Dictyococites*, 30, 51, 53, 56, 60
sculptilis lacinata, *Bulimina*, 65, 68, 74; pl. 14
Sea-level change, eustatic, Eocene, 21
Sedimentary sequence, Sierra Nevada region, Eocene, 41
semicostata lacrima, *Bulimina*, 17
seminulum, *Helicopontosphaera*, 31
Helicosphaera, 38, 56
semitriata, *Lagena*, 65, 71, 75; pl. 11
senilis, *Globigerina*, 59
sentifera, *Nodosaria paxillilis*, 71, 76; pl. 11
Serpula sulcata, 75
serraculoides, *Bramletteius*, 56, 60; pl. 9
shivelyi, *Textularia*, 69, 70, 77
Sierra Nevada, sedimentary sequence, Paleogene, 4, 41
sigmoides, *Zygodiscus*, 38
Sigmomorphina schencki, 65, 68, 73, 77
Silicosigmoilina, 7, 19
californica, 10, 17, 19; pl. 1
perplexa, 16
sp., 7
simiensis, *Cyclammina*, 11, 13, 16, 19; pl. 1
simplex, *Cristellaria*, 76
Lenticulina, 69, 71, 76
Robulus, 76
Siphogenerina eleganta, 74
hughesi, 71, 72, 77
mayi, 71, 72, 77; pl. 14
multicostata, 71, 72, 77; pl. 14
nodifera, 69, 70, 71, 77; pls. 10, 14, 15
raphanus transversa, 77
transversa, 71, 72, 77
Siphonina jacksonensis, 11, 17; pl. 4
wilcozensis, 11, 17, 28; pl. 4
sp., 17
Siphonodosaria gracilis, 17
sischoae, *Marginulina*, 11
Smith Grade-Empire Grade area, Santa Cruz Mountains, 5, 13
soldadoensis, *Acarinina*, 42
soldanii, *Gyroldina*, 65, 71, 75
octocamerata, *Gyroldina*, 10, 17
solitus, *Chiasmolithus*, 30, 56, 60
soluta, *Dentalina*, 65
sparsicostata, *Uvigerina*, 69, 70, 71, 77; pl. 15
Sphaerogypsina sp., pl. 6
Sphaeroidina bulloides, 69, 71, 77; pl. 13
Sphenolithus, 61
anarrhogus, 31
belemnus, 51, 61
ciperoensis Zone, 51, 52, 53, 74
Año Nuevo section, 51, 74
conicus, 51, 61
dissimilis, 51, 53, 61
distentus Zone, 36, 52, 53, 73, 74
nannofossils, 74
furcatolithoides, 31, 56, 61; pl. 9
heteromorphus, 51, 61; pl. 9
Zone, 51, 74
moriformis, 31, 38, 51, 53, 56, 61
orphanoknolli, 38
predistentus, 38, 53, 56, 61
pseudoradians, 31, 56, 61; pl. 9
radians, 31, 38, 51, 53, 56, 61
spiniger, 31, 56, 61; pl. 9
spp., 53, 56
spiniger, *Sphenolithus*, 31, 56, 61; pl. 9
spinosa, *Dentalina*, 10, 16
ornator, *Dentalina*, 64, 75; pl. 11
Spirogyphus (*Rotularia*), 41
(*Spirogyphus*) *tinajasensis*, *Rotularia*, 28
Spiroloculina lamposa, 11, 16
tezana, 65, 77
wilcozensis, 77
Spiroplectammina directa, 16, 65, 77; pls. 1, 10
richardi, 16, 19, 64, 77; pls. 1, 10
tejonensis, 11, 17
sp., 7, 69
Spiroplectoides directa, 77
spiropunctata, *Cibicides*, 68
spiropunctatus, *Cibicides*, 11, 22, 64, 75; pl. 16
Cibicidoides, 16
squamosa hexagona, *Entosolenia*, 75
Stages, time transgressive, 1, 62
Stainforthia sp., 64
staurion, *Coccolithus*, 56, 60
Cruciplacolithus, 30
Stichocassidulina thalmani, 64, 68, 77; pl. 13
Stichomitra asymbatos, 28
Stilostomella adolphina, 64, 71, 77
advena, 64, 69, 71, 77; pl. 13
lepidula, 65, 69, 71, 77
plummerae, 10
sanctaerucis, 71, 77
wegimanni, 71, 77; pl. 13
sp., 16, 65, 69, 71
Stratigraphic data, synthesis, 36
Stratigraphic range, *Bolivina marginata*, 69
Uvigerina jacksonensis, 69
Stratigraphy, Paleogene rocks, 41
striatoreticulata, *Camerina*, 45
Nummulites, 45
striatoreticulatus, *Nummulites*, 42, 45; pl. 6
strictus, *Discoaster*, 30
subangulata, *Gyroldinoides*, 16
Subbotina boweri, 12
frontosa, 7, 12
inaequispira, 7, 12
linaperta, 7, 12, 19
patagonica, 42
triangularis, 19
triloculinoides, 19
turgida, 19
velascoensis, 19
sp., 19
subbotinae, *Morozovella*, 19, 42
subbullata, *Marginulina*, 10, 16, 64, 69, 76; pl. 12
subconica, *eggerella*, 64, 75; pl. 10
subcylindrica, *Chiloguembelina midwayensis*, 28
subdistichus, *Coccolithus*, 30
subfusiformis, *Buliminella*, 69, 71, 74
sublodoensis, *Discoaster*, 29, 31, 33
subpertusa, *Ericsonia*, 38
subrecta, *Marginulina*, 69, 76
substriata, *Lagena*, 16
substrigata, *Dentalina*, 10
sulcata, *Lagena*, 69, 75
Serpula, 75
Sullivan, F. R., cited, 33, 63, 68, 69
Summary, benthic foraminiferal stratigraphy, Paleogene, 19
faunal assemblages, correlation, 72
paleobathymetry, central ranges of California, 19
Paleogene, California, 36
suteri, *Globorotaloides*, 51, 52, 59
Sveadal area, foraminifers, large, 44
Sveadal area, locality descriptions, 48
Systematic paleontology, 45
- T
- tani*, *Discoaster*, 31, 56, 61
ornatus, *Discoaster*, 31
Taxa, reference list, 74
Taxonomic notes, foraminifera, planktic, 59
tejonensis, *Spiroplectammina*, 11, 17
Temblor Formation, foraminifers, large, 44
San Jose area, 44
tenuicarinata, *Osangularia*, 11, 16; pl. 4
tenuis, *Cruciplacolithus*, 56, 60
Rhabdosphaera, 56, 61
tenuissima, *Fronicularia*, 69, 75
teres, *Pseudophragmina*, 42
Pseudophragmina (*Proporocyclina*), 47
terryi, *Lenticulina*, 10, 16
Tertiary provincial stages, relations to nannoplankton zones, 22
Tertiary stages, California, validity, 1
Tessellatolithus, 61
sp., 51, 53
tezana, *Spiroloculina*, 65, 77
Textularia shivelyi, 69, 70, 77
sp., Fairchild and others, 69, 77
thalmani, *Stichocassidulina*, 64, 68, 77; pl. 13
Thermocline, Paleogene, 21
theta, *Lenticulina*, 10, 17
Thoracosphaera, 61
sazea, 38
sp., 56
spp., 38, 56
Throckmorton, C. K., cited, 1, 22, 39, 49, 63, 71
tinajasensis, *Rotularia* (*Spirogyphus*), 28
tolmani, *Planularia*, 69
Toweius craticulus, 31
eminens, 31
Trachyleberidea, 12
transversa, *Siphogenerina*, 71, 72, 77
Siphogenerina raphanus, 77
Transversopontis pulcher, 31
pulcheroides, 56, 61
triangularis, *Gaudryina*, 69, 75; pl. 10
Subbotina, 19
triangulata, *Verneulina*, 10, 16, 28; pl. 1
Tribrachiatus orthostylus, 29, 31, 51, 53, 56, 61
Zone, 13, 28, 29, 31, 33
boundary, 19
/Discoaster lodoensis Zone, undifferentiated, 31
tribrachiatulus, *Discoaster*, 61
Trifarina, 7
advena californica, 10, 17, 28; pl. 3
hannai, 65, 77
wilcozensis, 65, 77; pl. 15
trihedra, *Bulimina*, 17
triloculinoides, *Subbotina*, 19
trinitatensis, *Bulimina*, 10, 12, 16; pl. 3
Nummulites, 45
tripartita, *Globigerina*, 51, 59
Triquetrorhabdulus carinatus, 51, 61
Zone, Año Nuevo section, 51, 74
Tritaxilina, 7, 13
colei, 11, 13, 16; pl. 1
pricipiensis, 13, 16
Trochammina, 7
globigeriniformis, 16
spp., 16, 71
Trochamminoides contortus, 16
Trochoaster, 61
sp., 51
truempyi, *Nuttallides*, 7, 10, 12, 13, 16, 19; pl. 4
truncana, *Planularia*, 11, 17
truncanella, *Bulimina*, 17
Truncatulina americana, 74
floridana, 75
pachyderma, 75
Truncorotaloides collectea, 59
rohri, 52, 59
tubulifera, *Angulogerina abbreviata*, 17
Tumey Hills, Fresno County, Calif., 5, 39
tumeyensis, *Valvulinaria*, 65, 68, 77
turbinata, *Lenticulina*, 11, 17
turgida, *Subbotina*, 19
Turritilina abbreviata, pl. 2
brevispira, 10, 16
turritilina praeturritilina, *Globigerina*, 59
Two-bar Shale Member, San Lorenzo Formation, Bear Creek area, 38, 39
San Lorenzo Formation, Bear Creek area, age, 39
boundary, Eocene, 39
coccoliths, 52
foraminifers, benthic, 68
planktic, 52, 53, 59

Twobar Shale Member—Continued
 San Lorenzo Formation—Continued
 Kings Creek section, 52, 59
 age, 52
 nannofossils, 38
 San Lorenzo River area, 38
 age, 39, 67
 San Lorenzo River section, 52, 59, 63, 67, 68,
 69
 age, 52
 zone assignments, 52, 59, 68
tympaiformis, *Fasciculithus*, 31, 38
 Type sections, benthic foraminiferal zones, 33

U

ulatisensis, *Lenticulina*, 16, 28; pl. 2
 Ulatisian-Narizian Stage, boundary, 29
 Ulatisian Stage, 28, 29, 33, 36, 42, 67, 68
 age, 29, 33, 36
 of Mallory, 42
umbilica, *Reticulofenestra*, 31, 38, 53, 56, 61; pl. 9
umbonata, *Anomalina*, 64
 Rotalina, 76
umbonatus, *Eponides*, 76
 Oridorsalis, 10, 16, 64, 69, 71, 76; pl. 17
unicavus, *Catapsydrax*, 51
 U.S. Committee on Stratigraphy, 1
utlisindex, *Globigerina*, 59
Uvigerina alabamensis, 11
 aubariana, 65, 69, 70, 71, 77; pl. 14
 churchi, 64, 68, 77; pl. 14
 cocoaensis, 68, 69, 73, 77
 first occurrence, 68
 species group, 65, 68, 69, 73
 stratigraphic range, 73
 elongata, 7, 11, 17, 64, 77; pls. 3, 14
 gallowayi, 69, 70, 71, 77; pl. 14
 Zone, 70, 73
 Zemmorrian Stage, 58, 73
garzaensis, 65, 68, 77; pl. 14
 nudorobusta, 69
 hispidocostata, 77
 jacksonensis, 69
 stratigraphic range, 69
lodoensis, 28
 miriamae, 11, 16; pl. 3
vicksburgensis Zone, 69
 sp., 71
Uvigerinella obesa impolita, 71, 77
 sparsicostata, 69, 70, 71, 77; pl. 15
 Zone, 71, 73
 Uvigerinids, *elongata* group, 12

V, W

Vaginulinopsis asperuliformis, 11, 22; pl. 2

Vaginulinopsis—Continued

echinata, 16
kelleyi, 10, 16
mexicana, 28
 kerni, 11, 17; pl. 2
 nudicostata, 10, 17; pl. 2
saundersi, 19, 64, 77; pl. 12
 sp., 17
Valvulineria, 12
 allomorphinoides, 11
 araucana, 71, 77; pl. 15
 californica, 71, 72, 77
 childsi, 10, 16
 indiscriminata, 28
 tumeyensis, 65, 68, 77
 Zone, lower Refugian, 69
 sp., of Mallory, 17
 sp., 11
 Vaqueros Formation, Año Nuevo Section, 51, 71
 coccoliths, 51
 foraminifers, assemblages, 71, 72
 benthic, 71, 72
 planktic, 51
 stage assignments, 72
 Vaqueros Sandstone, 1
 coccoliths, 52
 San Lorenzo River section, 63
 Zayante Creek section, 70
variabilis, *Discoaster*, 51, 61
vaughani, *Plectofrondicularia*, 65, 69, 71, 76; pl. 13
 Vedder, J. G., cited, 1
velascoensis, *Morozovella*, 19
 Nodosaria, 17
 Subbotina, 19
Verneuilina, 19
 triangulata, 10, 16, 28; pl. 1
 sp., of Fairchild and others, 69, 71, 77
 sp., pl. 10
 Vertebrate fauna, California, 1
vesper, *Micrantholithus*, 31
vicksburgensis, *Bifarina*, 10
Virgulina bramlettei, 75
 californiensis, 75
 dibollensis, 75
 wilcozensis, 77
vortex, *Lenticulina*, 10, 16
vulgaris, *Lagena*, 65, 71, 75
Vulvulina curta, 10, 16
 Wagonwheel Formation, Devils Den area, 30
 Warren, A. D., cited, 41, 44
washingtonensis, *Karreriella*, 65, 68, 73, 75; pl. 11
 Weaver, C. E., cited, 1
 Weaver, D. W., cited, 22
weaveri, *Lenticulina*, 10
wegimanni, *Nodogenerina*, 77
 Stilostomella, 71, 77; pl. 13
welchi, *Lenticulina*, 17, 65, 68, 69, 76; pl. 12
 Robulus, 76

welleri, *Anomalinoidea*, 10, 16, 19
wellsi, *Eocumuloides*, 42, 46; pl. 6
wemmelenensis, *Discoaster*, 31
 White, R. T., cited, 39, 40
whitei, *Bulimina*, 11, 16; pl. 3
 Cibicidoides, 10, 16, 19; pl. 5
 Plectofrondicularia, 10; pl. 2
wilcozensis, *Acarinina*, 19
 Angulogerina, 28; pl. 3
 californica, *Alabamina*, 16, 22
 Dentalina, 11
 Siphonina, 11, 17, 28; pl. 4
 Spiroloculina, 77
 Trifarina, 65, 77; pl. 15
 Virgulina, 77
willcoxi, *Camerina*, 45
 Nummulites, 42, 45; pl. 6
 Operculinoides, 45
wilsoni, *Globorotaloides*, 59
woodi, *Globigerina*, 51
 Worm tube, 28, 41

Y, Z

Ynezian Stage, 28, 29, 33, 36
 age, 29, 36
 type section, 33
 Ynezian-Bulitian Stage boundary, 29, 33
 Zayante Creek, 70
 Zayante Creek section, 52, 59, 70
 coccolith assemblages, 52
 faunal assemblages, 74
 foraminifers, benthic, 69
 planktic, 52
 Santa Cruz Mountains, 49, 70
Zeauvigerina, sp., 17
 Zemmorrian Stage, 36, 38, 52, 53, 58, 59, 69, 70, 71, 72,
 73
 age, 36, 38, 70
 boundary, 70, 71, 73
 criteria, 73
 faunal assemblages, 69, 70, 71
 Uvigerina gallowayi Zone, 58, 73
 Zemmorrian Stage-Refugian Stage boundary, 53, 69, 73
 Zemmorrian Stage-Saucesian Stage boundary, 52, 59,
 72, 73, 74
zetina indirecta, *Fursenkoina*, 11, 17
 Zone assignments, primary indicators, 59
 Zone P14-P15 boundary, 73
Zygodiscus adamas, 31
 plectopons, 31
 sigmoides, 38
Zygodiscus dubius, 56, 61; pl. 9
Zygrhablithus, 61
 bijugatus, 31, 38, 56, 61
 sp., 51, 53


PLATES 1-17

**[Contact photographs of the plates in this report are available, at cost,
from U.S. Geological Survey Library, Federal Center, Denver, Colorado 80225]**

PLATE 1

[All samples from either the aqueduct section in the Devils Den area or from the Lodo Gulch section. All figures × 65]

- FIGURE 1. *Ammodiscus* cf. *A. pennyi* Cushman and Jarvis
Aqueduct section, sample 76CB 1281H
- 2, 3, 4. *Cyclammmina incisa* (Stache)
Aqueduct section, sample 76CB 1281
5. *Cyclammmina simiensis* Berry
Aqueduct section, sample 76CB 1281
6. *Cyclammmina* cf. *C. clarki* (Hanna)
Aqueduct section, sample 76CB 1291
7. *Spiroplectammmina richardi* Martin
Lodo Gulch section, sample 76CB 1231B
- 8, 9. *Spiroplectammmina directa* (Cushman and Siegfus)
Aqueduct section, sample 76CB 1281M
10. *Verneuilina triangulata* Cook
Lodo Gulch section, sample 76CB 1231A
11. *Gaudryina coalingensis* Cushman and Hanna
Lodo Gulch section, sample 76CB 1231A
12. *Gaudryina* cf. *G. laevigata* Franke
Aqueduct section, sample 76CB 1281
13. *Clavulinoides californicus* Mallory
Aqueduct section, sample 76CB 1281
14. *Dorothia germanica* Cushman
Aqueduct section, sample 76CB 1281
15. *Tritaxilina colei* Cushman and Siegfus
Aqueduct section, sample 76CB 1281
16. *Karrieriella monumentensis* Mallory
Aqueduct section, sample 76CB 1281
17. *Karrieriella mediaguaensis* Mallory
Aqueduct section, sample 76CB 1281
18. *Martinottiella eocenica* Cushman and Bermudez
Aqueduct section, sample 76CB 1281
19. *Silicosigmoilina californica* Cushman and Church
Aqueduct section, sample 76CB 1281L




BENTHIC FORAMINIFERS FROM THE CALIFORNIA COAST RANGES

PLATE 2

[All samples from either the aqueduct section in the Devils Den area or from the Lodo Gulch section]

- FIGURE 1. *Lenticulina ulatisensis* (Boyd)
 Aqueduct section, sample 76CB 1273, × 65
- 2, 3. *Vaginulinopsis mexicana* var. *kerni* (Cook)
 Lodo Gulch section, samples 76CB 1221G, 76 CB 1231A, × 100
- 4, 5. *Vaginulinopsis asperuliformis* (Nuttall)
 Lodo Gulch section, sample 76CB 1231B, × 100
6. *Vaginulinopsis mexicana* var. *nudicostatus* (Cushman and Hanna)
 Lodo Gulch section, sample 76CB 1231A, × 15
7. *Amphimorphina ignota* Cushman and Siegfus
 Aqueduct section, sample 76CB 1281D, × 65
8. *Plectofrondicularia kerni* Cook
 Lodo Gulch section, sample 76CB 1231D, × 65
9. *Plectofrondicularia whitei* Martin
 Lodo Gulch section, sample 76CB 1231, × 65
10. *Plectofrondicularia minuta* Sullivan
 Aqueduct section, sample 76CB 1281, × 65
- 11, 12. *Buliminella grata* var. *convoluta* Mallory
 Aqueduct section, sample 76CB 1281, × 65
13. *Buliminella bradburyi* (Martin)
 Aqueduct section, sample 76CB 1281, × 100
14. *Turritilina abbreviata* Ten Dam
 Lodo Gulch section, sample 76CB 1221G, × 65
- 15, 16, 17. *Aragonia aragonensis* (Nuttall)
 Lodo Gulch section, samples 76CB 1221E, × 100
 76CB 1221G, × 200
 76CB 1221G, × 235
18. *Bifarina eleganta* (Plummer)
 Lodo Gulch section, sample 76CB 1221H, × 65
19. *Bifarina nuttalli* Cushman and Siegfus
 Aqueduct section, sample 76CB 1633, × 100



BENTHIC FORAMINIFERS FROM THE CALIFORNIA COAST RANGES

PLATE 3

[All samples from either the aqueduct section in the Devils Dens area or from the Lodo Gulch section]

FIGURES 1-5. *Bulimina callahani* Galloway and Morrey

Lodo Gulch section sample 76CB 1231D, 1-4; $\times 200$

Aqueduct section, sample 76CB 1281D, 5; $\times 120$

6. *Bulimina curtissima* Cushman and Siegfus 1221G, $\times 200$

7. *Bulimina impendens* Parker and Bermudez

Lodo Gulch section, sample 76CB 1231D, $\times 200$

8. *Bulimina macilenta* Cushman and Parker

Lodo Gulch section, sample 76CB 1231B, $\times 65$

9, 10. *Bulimina trinitatensis* Cushman and Jarvis

Aqueduct section, sample 76CB 1281, 9; $\times 65$, 10; $\times 235$

11, 12. *Bulimina whitei* Martin

Aqueduct section, sample 76CB 1273, $\times 100$

13. *Bolivina huneri* Howe

Aqueduct section, sample 76CB 1281, $\times 135$

14. *Angulogerina* cf. *A. wilcoxensis* Cushman and Ponton

Aqueduct section, sample 76CB 1281C, $\times 100$

15. *Trifarina advena* var. *californica* Mallory

Aqueduct section, sample 76CB 1281D, $\times 100$

16, 17. *Uvigerina elongata* Cole

Lodo Gulch section, sample 76CB 1231, 76CB 1231C, $\times 200$

18, 19, 20. *Uvigerina lodoensis* var. *miriamae* Mallory


Lodo Gulch section, sample 76CB 1221G, 18; $\times 200$

76CB 1221G, 19; $\times 135$

sample 76CB 1221C, 20; $\times 135$

21. *Loxostomoides applinae* (Plummer)

Lodo Gulch section, sample 76CB 1231D, $\times 65$




BENTHIC FORAMINIFERS FROM THE CALIFORNIA COAST RANGES

PLATE 4

[All samples are from either the aqueduct section in the Devils Den area or from the Lodo Gulch section]

- FIGURE 1. *Nodosarella advena* Cushman and Siegfus
Aqueduct section, sample 76CB 1281E, $\times 65$
- 2, 3, 4. *Nuttallides truempyi* (Nuttall)
Aqueduct section, sample 76CB 1281, $\times 65$
5. *Baggatella californica* Mallory
Aqueduct section, sample 76CB 1281, $\times 100$
- 6, 7. *Siphonina* cf. *jacksonensis* Cushman
Aqueduct section, sample 76CB 1281D, $\times 135$
- 8, 9, 10. *Siphonina wilcoxensis* Cushman
Aqueduct section, sample 76CB 1281D, $\times 100$
- 11, 12. *Osangularia tenuicarinata* (Cushman and Siegfus)
Aqueduct section, sample 76CB 1281, $\times 100$
- 13, 14. *Osangularia mexicana* (Cole)
Aqueduct section, sample 76CB 1281, $\times 165$
- 15, 16, 17. *Globocassidulina globosa* Hantken
Lodo Gulch section, sample 76CB 1231B, 15, 16; $\times 100$, 17; $\times 665$




BENTHIC FORAMINIFERS FROM THE CALIFORNIA COAST RANGES

PLATE 5

[All samples are from either the aqueduct section in the Devils Den area or from the Lodo Gulch section. All figures × 65]

- FIGURES 1, 2, 3. *Anomalinoides acutus* (Plummer)
Lodo Gulch section, sample 76CB 1231C
- 4, 5, 6. *Anomalinoides crassiseptus* (Cushman and Siegfus)
Aqueduct section, sample 76CB 1281H
- 7, 8, 9. *Anomalinoides aragonensis* (Cole)
Lodo Gulch section, sample 76CB 1231G
- 10, 11, 12. *Anomalinoides keeni* (Martin)
Lodo Gulch section, sample 76CB 1231B
- 13, 14, 15. *Anomalinoides garzaensis* (Cushman and Siegfus)
Aqueduct section, sample 76CB 1281
- 16, 17, 18. *Cibicidoides beatus* (Martin)
Lodo Gulch section, sample 76CB 1231C
- 19, 20, 21. *Cibicidoides grimsdalei* (Nuttall)
Aqueduct section, sample 76CB 1281J
- 22, 23, 24. *Cibicidoides fortunatus* (Martin)
Aqueduct section, sample 76CB 1284
- 25, 26, 27. *Cibicidoides whitei* (Martin)
Aqueduct section, sample 76CB 1281J




BENTHIC FORAMINIFERS FROM THE CALIFORNIA COAST RANGES

PLATE 6

[All figures × 20]

- FIGURES 1, 2, 14. *Amphistegina parvula* (Cushman) (p. 46)
1, Equatorial section; 2, Subaxial section, locality 76CB1633 (Devils Den)
14, Axial section, locality 71CB971B (San Jose)
- 3, 13. *Eoconuloides lopeztrigoi* (D. K. Palmer) (p. 46)
3, Axial section, locality 76CB1634 (Devils Den)
13, Axial section, 71CB971B (San Jose)
4. *Eoconuloides wellsi* Cole and Bermudez (p. 46)
Subequatorial section, 76CB1634 (Devils Den)
5. *Pseudophragmina* sp.
Axial section, 76CB1641 (Devils Den)
- 6, 7, 10. *Nummulites willcoxi* Heilprin (p. 45)
6, 7, Subaxial section; 10, Equatorial section, 71CB983C (San Jose)
- 8, 16. *Eofabiana grahami* Kupper (p. 47)
8, Locality 71CB983C (San Jose);
16, Osbun 49-6 (Sveadal)
9. *Nummulites striatoreticulatus* L. Rutten (p. 45)
Axial section, 71CB983C (San Jose)
11. *Asterocyclina aster* (Woodring) (p. 47)
Subequatorial section, 71CB983C (San Jose)
12. *Pseudophragmina (Proporocyclina) flintensis* (Cushman) (p. 46)
Axial section
15. *Sphaerogypsina* sp. Osbun 49-6 (Sveadal)
17. *Pseudophragmina (Proporocyclina) clarki* (Cushman) (p. 47)
Axial section (A); Locality Osbun 1-1B (Sveadal)




LARGE FORAMINIFERS FROM CALIFORNIA

PLATE 7

[Scale bar varies]

- FIGURES 1, 2. *Chiloguembelina cubensis* (Palmer). 1, Side view, sample Mf1350. 2, Edge view of fig. 1. Scale bar=30 μ m for both. (p. 59)
- 3, 6. *Pseudohastigerina lillisi* (Church). 3, Side view, sample Mf1351. 6, Edge view, sample Mf1351. Scale bar=100 μ m for both. (p. 60)
- 4, 5. *Pseudohastigerina micra* (Cole). 4, Side view, sample Mf1351. 5, Edge view, sample Mf1351. Scale bar=30 μ m for both. (p. 60)
- 7-9. *Gümbelitra columbiana* Howe. All specimens from sample Mf4746. 7, Oblique spiral view. 8, Oblique umbilical view. Scale bar=30 μ m for all. (p. 60)




PLANKTIC FORAMINIFERS FROM SAN LORENZO FORMATION

PLATE 8

[Scale bar varies]

- FIGURES 1-4. *Globorotalia nana* Bolli. All specimens from the Vaqueros(?) Formation, Año Nuevo section. 1, Umbilical view, sample Mf4665. 2, Side view, sample Mf4664. 4, Spiral view, sample Mf4665. Scale bar=30 μm for all. (p. 60)
- 5, 6. *Globorotalia* sp. aff. *G. munda* Jenkins. Both specimens from sample Mf4665 in the Vaqueros(?) Formation, Año Nuevo section. 5, Umbilical view. 6, Side view. Scale bar=30 μm for both. (p. 60)
- 7, 8. *Globigerina praeturrilina* Blow and Banner. 7, Oblique umbilical view, sample Mf3305 from the Twobar Shale Member of the San Lorenzo Formation, San Lorenzo River section. 8, Oblique side view of fig. 7. Scale bar=100 μm for both. (p. 59)




PLANKTIC FORAMINIFERS FROM AÑO NUEVO AND SAN LORENZO RIVER SECTIONS

PLATE 9

[All figures are photomicrographs at the same magnification; scale bar equals 5 μ m. B F, bright field; XP, cross-polarized light]

- FIGURES 1, 2. *Bramletteius serraculoides* Gartner. Sample Mf3301 from the Butano Sandstone in the San Lorenzo River section. 1, XP. 2, BF.
- 3, 4. *Coccolithus formosus* Kamptner. Sample Mf3306 from the Twobar Shale Member in the San Lorenzo River section. 3, XP. 4, BF.
5. *Daktylethra punctulata* Gartner. Sample Mf3306 from the Twobar Shale Member in the San Lorenzo River section; XP.
6. *Discoaster barbadiensis* Tan. Sample Mf3301 from the Butano Sandstone in the San Lorenzo River section; BF.
7. *Discoaster deflandrei* Bramlette and Riedel. Sample Mf4668 from the Vaqueros(?) Formation in the Año Nuevo section; BF.
8. *Discoaster nodifer* (Bramlette and Riedel). Sample Mf3301 from the Butano Sandstone in the San Lorenzo River section; BF.
9. *Discoaster saipanensis* Bramlette and Riedel. Sample Mf3306 from the Twobar Shale Member in the San Lorenzo River section; BF.
- 10, 11. *Dictyococcites bisectus* (Hay, Mohler, and Wade). Sample Mf4664 from the Vaqueros(?) Formation in the Año Nuevo section. 10, BF. 11, XP.
- 12, 13. *Lanternithus minutus* Stradner. Sample Mf3306 from the Twobar Shale Member in the San Lorenzo River section. 12, BF. 13, XP.
14. *Micrantholithus altus* Bybell and Gartner. Sample Mf3306 from the Twobar Shale Member in the San Lorenzo River section; BF.
15. *Reticulofenestra reticulata* (Gartner and Smith). Sample Mf3306 from the Twobar Shale Member in the San Lorenzo River section; XP.
16. *Reticulofenestra umbilica* (Levin). Sample Mf3301 from the Butano Sandstone in the San Lorenzo River section; XP.
17. *Sphenolithus furcatolithoides* Locker, below; *Sphenolithus spiniger* Bukry, above. Sample Mf3301 from the Butano Sandstone in the San Lorenzo River section; XP.
- 18, 19. *Sphenolithus heteromorphus* Deflandre. Sample Mf4668 from the Vaqueros(?) Formation in the Año Nuevo section; XP. 18, Basal spines. 19, Apical and basal spines.
- 20, 21. *Sphenolithus pseudoradians* Bramlette and Wilcoxon. Sample Mf3301 from the Butano Sandstone in the San Lorenzo River section. 20, BF. 21, XP.
22. *Zygodolithus dubius* Deflandre. Sample Mf3306 from the Twobar Shale Member in the San Lorenzo River section; BF.




1

2

3

4

5




6

7

8

9




10

11

12

13

14



15

16

17

18

19

20

21




22

CALCAREOUS NANNOFOSSILS FROM THE SANTA CRUZ MOUNTAINS

PLATE 10


- FIGURE
1. *Rhabdammina eocenica* Cushman and Hanna, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 300 μ m. USNM no. 262082
 2. *Reophax pilulifer* Brady, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 300 μ m. USNM no. 262083
 3. *Haplophragmoides deflata* Sullivan, Mf1352 (field no. 76CB1354), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262084
 4. *Haplophragmoides* sp. #1, Mf1352 (field no. 76CB1354), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262085
 5. *Haplophragmoides* sp., Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262086
 6. *Cyclammmina cancellata* Brady *obesa* Cushman and Laiming, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 1 mm. USNM no. 262087
 7. *Bolivina marginata* Cushman, *Siphogenerina nodifera* Cushman and Kleinpell, and *Orthomorphina rohri* (Cushman and Stainforth) attached to the test of *Cyclammmina cancellata* Brady *obesa* Cushman and Laiming, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 300 μ m.
 8. *Spiroplectammina richardi* Martin, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262088
 9. *Spiroplectammina directa* (Cushman and Siegfus), Mf1360 (field no. EB256A), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262089
 10. *Verneuilina* sp., Mf4683 (field no. 76CB1603), Zayante Creek section. Bar equals 100 μ m. USNM no. 262090
 11. *Verneuilina* sp., Mf4683 (field no. 76CB1603), Zayante Creek section. Bar equals 30 μ m.
 12. *Gaudryina gracilis* Cushman and Laiming, Mf4682 (field no. 76CB1602), Zayante Creek section. Bar equals 100 μ m. USNM no. 262092
 13. *Gaudryina gracilis* Cushman and Laiming, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262093
 14. *Gaudryina triangularis* Cushman, Mf4682 (field no. 76CB1602), Zayante Creek section. Bar equals 100 μ m. USNM no. 262094
 15. *Dorothia cubana* (Cushman and Bermudez), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262095
 16. *Eggerella elongata* Blaisdell, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262096
 17. *Eggerella subconica* Parr, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262097



BENTHIC FORAMINIFERA FROM THE SANTA CRUZ MOUNTAINS

PLATE 11


- FIGURE
1. *Karreriella elongata* Mallory, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262098
 2. *Karreriella washingtonensis* Rau, Mf1360 (field no. EB256A3), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262099
 3. *Martinottiella eocenica* Cushman and Bermudez, Mf3304 (field no. 76CB1451, San Lorenzo River section. Bar equals 100 μ m. USNM no. 262100
 4. *Martinottiella patens* (Cushman and Laiming), Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 300 μ m. USNM no. 262101
 5. *Plectina garzaensis* Cushman and Siegfus, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262102
 6. *Plectina garzaensis* Cushman and Siegfus, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 300 μ m. USNM no. 262103
 - 7, 8. *Quinqueloculina minuta* Beck, Mf3304 (field no. 76CB1451), San Lorenzo River Section. Bar equals 100 μ m. USNM no. 262104
 9. *Nodosaria gyrata* Mallory, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262105
 10. *Nodosaria parexilis sentifera* Cushman and Parker, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 100 μ m. USNM no. 262106
 11. *Dentalina cooperensis* Cushman, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262107
 12. *Dentalina quadrulata* Cushman and Laiming, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262108
 13. *Dentalina* sp., Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262109
 14. *Dentalina spinosa ornatior* Smith, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 300 μ m. USNM no. 262110
 15. *Lagena acuticosta* Reuss, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 30 μ m. USNM no. 262111
 16. *Lagena semistriata* Williamson, Mf2700 (field no. EB395D), San Lorenzo River section. Bar equals 30 μ m. USNM no. 262112
 17. *Lenticulina barbati* (Cushman and Hobson), Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 300 μ m. USNM no. 262113
 18. *Lenticulina calcar* (Linne), Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 100 μ m. USNM no. 262114
 19. *Lenticulina* cf. *L. clypeiformis* d'Orbigny, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 300 μ m. USNM no. 262115
 20. *Lenticulina insuetus* (Cushman and Stainforth), Mf1350 (field no. 76CB1352), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262116



BENTHIC FORAMINIFERS FROM THE SANTA CRUZ MOUNTAINS

PLATE 12


- FIGURE
1. *Lenticulina insuetus* (Cushman and Stainforth), Mf1356 (field no. EB256B), San Lorenzo River section. Bar equals 300 μ m. USNM no. 262117
 2. *Lenticulina midwayensis* (Plummer), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 300 μ m. USNM no. 262118
 3. *Lenticulina pseudocultratus* (Cole), Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 2621109
 4. *Lenticulina pseudovortex* (Cole), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262120
 5. *Lenticulina* sp., Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262121
 6. *Lenticulina* sp., Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 300 μ m. USNM no. 262122
 7. *Lenticulina* spp., Mf304 (field no. 76CB1451), San Lorenzo River section. Bar equals 300 μ m. USNM no. 262123
 8. *Lenticulina welchi* (Church), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262124
 9. ?*Marginulina* sp., Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262125
 10. *Marginulina subbullata* Hantken, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262126
 11. *Orthomorphina rohri* (Cushman and Stainforth), Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 300 μ m. USNM no. 262127
 12. *Pseudonodosaria gallowayi* (Cushman), Mf4682 (field no. 76CB1602), Zayante Creek section. Bar equals 100 μ m. USNM no. 262128
 13. *Saracenaria schencki* Cushman and Hobson, Mf1360 (field no. EB256A3), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262129
 14. *Vaginulinopsis saundersi* (Hanna and Hanna), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 300 μ m. USNM no. 262130
 15. *Plectofrondicularia miocenica directa* Cushman and Laiming, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 300 μ m. USNM no. 262131



BENTHIC FORAMINIFERS FROM THE SANTA CRUZ MOUNTAINS

PLATE 13


- FIGURE
1. *Plectofrondicularia packardi* Cushman and Schenck, Mf1360 (field no. EB256A), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262132
 2. *Plectofrondicularia packardi multilineata* Cushman and Simonson, Mf1360 (field no. EB256A), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262133
 3. *Plectofrondicularia vauhani* Cushman, Mf1360 (field no. EB256A), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262134
 4. *Amphimorphina becki* Mallory, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262135
 5. *Buliminella elegantissima* (d'Orbigny), Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 30 μ m. USNM no. 262136
 6. *Sphaeroidina bulloides* d'Orbigny, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262137
 7. *Sphaeroidina bulloides* d'Orbigny, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262138
 8. *Bolivina kleinpelli* Beck, Mf2700 (field no. EB395D), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262139
 9. *Bolivina marginata* Cushman, Mf2701 (field no. EB675), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262140
 10. *Bolivina marginata* Cushman, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262141
 11. *Bolivina marginata* Cushman, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 100 μ m. USNM no. 262142
 12. *Bolivina scabrata* Cushman and Bermudez, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 30 μ m. USNM no. 262143
 13. *Bolivina* sp., Mf4661 (field no. 76CB1451), Año Nuevo section. Bar equals 100 μ m. USNM no. 262144
 14. *Bolivinoidea mexicanus* (Cole), Mf1351 (field no. 76CB1353), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262145
 15. *Cassidulinoides californiensis* Bramlette, Mf2700 (field no. EB395D), San Lorenzo River section. Bar equals 30 μ m. USNM no. 262146
 16. *Stichocassidulina* cf. *S. thalmani* Stone, Mallory, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262147
 17. *Stilostomella advena* (Cushman and Laiming), Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 100 μ m. USNM no. 262148
 18. *Stilostomella wegmanni* (Cole), Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 300 μ m. USNM no. 262149
 19. *Bulimina carneroensis* Cushman and Kleinpell, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 100 μ m. USNM no. 262150



BENTHIC FORAMINIFERS FROM THE SANTA CRUZ MOUNTAINS

PLATE 14


- FIGURE
1. *Bulimina carneroensis* Cushman and Kleinpell, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262151
 2. *Bulimina corrugata* Cushman and Siegfus, Mf1357 (field no. EB256K), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262152
 3. *Bulimina corrugata* Cushman and Siegfus, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 30 μ m. USNM no. 262153
 4. *Bulimina curtissima* Cushman and Siegfus, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262154
 5. *Bulimina inflata alligata* Cushman and Laiming, Mf2700 (field no. EB395D), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262155
 6. *Bulimina microcostata* Cushman and Parker, Mf1531 (field no. 76CB1353), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262156
 7. *Bulimina sculptilis lacinata* Cushman and Parker, Mf1356 (field no. EB256B), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262157
 8. *Bulimina sculptilis lacinata* Cushman and Parker, Mf1356 (field no. EB256B), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262158
 9. *Globobulimina pacifica* Cushman, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262159
 10. *Uvigerina auberiana* d'Orbigny, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262161
 11. *Uvigerina auberiana* d'Orbigny, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262161.
 12. *Uvigerina churchi* Cushman and Siegfus, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262162
 13. *Uvigerina elongata* Cole, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262163
 14. *Uvigerina gallowayi* Cushman, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 100 μ m. USNM no. 262164
 15. *Uvigerina garzaensis* Cushman and Siegfus, Mf1356 (field no. EB256B), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262165
 16. *Siphogenerina mayi* Cushman and Parker, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262166
 17. *Siphogenerina multicostata* Cushman and Jarvis, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262167
 18. *Siphogenerina nodifera* Cushman and Kleinpell, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 300 μ m. USNM no. 262168



BENTHIC FORAMINIFERS FROM THE SANTA CRUZ MOUNTAINS

PLATE 15


- FIGURE
1. *Siphogenerina nodifera* Cushman and Kleinpell, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 300 μm . USNM no. 262169
 2. *Siphogenerina nodifera* Cushman and Kleinpell, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 300 μm . USNM no. 262170
 3. *Siphogenerina nodifera* Cushman and Kleinpell, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 300 μm . USNM no. 262171
 4. *Siphogenerina nodifera* Cushman and Kleinpell, Mf4682 (field no. 76CB1602), Zayante Creek section. Bar equals 300 μm . USNM no. 262172
 5. *Trifarina wilcoxensis* (Cushman and Ponton), Mf1531 (field no. 75CB1353), San Lorenzo River section. Bar equals 100 μm . USNM no. 262173
 6. *Uvigerinella sparsicostata* Cushman and Laiming, Mf4683 (field no. 76CB1603), Zayante Creek section. Bar equals 100 μm . USNM no. 262174
 7. *Baggina robusta* Kleinpell, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μm . USNM no. 262175
 8. *Valvulineria araucana* (d'Orbigny), Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μm . USNM no. 262176
 9. *Bifarina eleganta* (Plummer), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μm . USNM no. 262177
 - 10, 11, 12. ?*Neoeponides* sp., Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μm . USNM no. 262178
 13. *Cibicides floridanus* (Cushman), Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μm . USNM no. 262179
 - 14, 15. *Cibicides americanus crassiseptus* Cushman and Laiming, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 100 μm . USNM no. 262180
 16. *Cibicides haydoni* (Cushman and Schenck), Mf1360 (field no. EB2456A) San Lorenzo River section. Bar equals 100 μm . USNM no. 262181



BENTHIC FORAMINIFERS FROM THE SANTA CRUZ MOUNTAINS.

PLATE 16


- FIGURE
1. *Cibicides pseudoungerianus evolutus* Cushman and Hobson, Mf2700 (field no. EB395D), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262182
 - 2, 3. *Cibicides spiropunctatus* Galloway and Morrey, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262183
 4. *Pleurostomella alternans* Schwager, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 30 μ m. USNM no. 262184
 5. *Nodosarella atlantisae hispidula* (Cushman), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262185
 6. *Nodosarella atlantisae hispidula* (Cushman), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262186
 7. *Fursenkoina bramlettei* (Galloway and Morrey), Mf4667 (field no. 76CB1541), Año Nuevo section. Bar equals 100 μ m. USNM no. 262187
 8. *Fursenkoina bramletti* (Galloway and Morrey), Mf1531 (field no. 75CB1353), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262188
 9. *Fursenkoina californiensis* (Cushman), Mf3304 (field no. 75CB1451), San Lorenzo River section. Bar equals 30 μ m. USNM no. 262189
 10. *Fursenkoina dibollensis* (Cushman and Applin), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262190
 11. *Cassidulina crassipunctata* Cushman and Hobson, Mf2700 (field no. EB395D), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262191
 12. *Cassidulina crassipunctata* Cushman and Hobson, Mf1360 (field no. EB256A), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262192
 13. *Cassidulina diversa* Cushman and Stone, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262193
 14. *Nonionella miocenica* Cushman, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 100 μ m. USNM no. 262194
 15. *Nonionella miocenica* Cushman, Mf2700 (field no. EB395D), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262195



BENTHIC FORAMINIFERS FROM THE SANTA CRUZ MOUNTAINS

PLATE 17

- FIGURE 1, 2. *Pullenia quinqueloba* (Reuss), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262196
3. *Pullenia miocenica* Kleinpell, Mf4664 (field no. 76CB1531), Año Nuevo section. Bar equals 100 μ m. USNM no. 262197
4. *Gyroidina orbicularis planata* Cushman, Mf2700 (field no. EB395D), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262198
- 5, 6. *Oridorsalis umbonatus* (Reuss), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262199
7. *Anomalina californiensis* Cushman and Hobson, Mf4665 (field no. 76CB1532), Año Nuevo section. Bar equals 100 μ m. USNM no. 262200
- 8, 9. *Anomalina garzaensis* Cushman and Siegfus, Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262201
10. *Anomalina salinasensis* Kleinpell, Mf4661 (field no. 76CB1541), Año Nuevo section. Bar equals 100 μ m. USNM no. 262202
- 11, 12. *Boldia hodgei* (Cushman and Schenck), Mf3304 (field no. 76CB1451), San Lorenzo River section. Bar equals 100 μ m. USNM no. 262203



BENTHIC FORAMINIFERS FROM THE SANTA CRUZ MOUNTAINS

